

ANTIOXIDANT AND ANTIBACTERIAL ACTIVITIES OF *TRIANTHEMA DECANDRA* LINN*SUKANTHA T.A¹, SHUBASHINI K.SRIPATHI², RAVINDRAN N.T.³ AND BALASHANMUGAM, P⁴

¹Department of Chemistry, K.S. Rangasamy College of Technology, Tiruchengode 637215, Tamil Nadu, India, ²Department of Chemistry, Avinashilingam Institute of Home Science and Higher Education for Women, Coimbatore 641 043, Tamil Nadu, India, ³Department of Biochemistry, Adhiparasakthi College of Arts and Science, G.B. Nagar, Kalavai 632506, Tamil Nadu, India, ⁴CAS in Botany, University of Madras, Guindy Campus, Chennai 600 025, Tamil Nadu, India. Email: tasukantha@gmail.com.

Received: 6 Sep 2011, Revised and Accepted: 6 Dec 2011

ABSTRACT

Trianthema decandra (Vellai sharuni in Tamil and Gadabani in Hindi) is a prostrate weed widely distributed in India. Its root is traditionally used in the treatment of hepatitis and asthma and for wound-healing. The antioxidant and antibacterial potential of *Trianthema decandra* was assessed and the results revealed significant activity in the ethyl acetate and methanolic extracts of roots and leaves of the plant.

Keywords: *Trianthema decandra*, Antioxidant, Antibacterial, Phenolics, DPPH scavenging.

INTRODUCTION

Medicinal plants are gifts of nature to cure limitless number of diseases in human beings¹. The abundance of plants on the earth's surfaces has led to an increasing interest in the investigation of different extracts obtained from traditional medicinal plants as potential sources of new therapeutic agents². Antioxidants are a group of compounds that facilitate survival of plants and may promote the health of humans who consume a variety of plant foods³. In plants, the term 'antioxidant' refers to a wide range of phenolic compounds that vary from simple phenolic acids to highly polymerized compounds such as tannins. Phenolic compounds or polyphenols are categorized into 15 main classes with more than 8,000 identified compounds. The largest category is the flavonoid group, comprising 13 classes with over 5,000 compounds⁴. In plants, polyphenols are important for structural supports, as antiherbivorous substances, attracting pollinators, for protection from ultraviolet radiation and for wound repair⁵. The human body also synthesizes endogenous antioxidants such as superoxide dismutases, glutathione peroxidases, alpha-tocopherol and melatonin to counteract cellular damage by active oxygen and free radicals⁶. Many studies suggest that endogenous antioxidants or exogenous antioxidants from diet can function as free radical scavengers and improve human health. In this regard, consumption of a variety of plant foods may provide additional health benefits^{6,7}.

The use of plant extracts and phytochemicals with known antimicrobial properties can be of great significance in therapeutic treatments. In the last few years, a number of studies have been conducted to prove such efficiency⁸⁻¹⁰. Many plants have been used because of their antimicrobial traits, which is due to compounds synthesized as secondary metabolites by the plant. These products are known by their active substances, for example, the phenolic compounds which are part of the essential oils¹¹, as well as tannins¹².

Trianthema decandra (Family: *Aizoaceae*) is a prostrate, glabrous, succulent and annual weed found almost throughout India, in cultivated and waste land. The genus *Trianthema* consists of 20 species, but only a few have been phytochemically investigated. It is commonly known as *gadabani* (Hindi) and *vellai sharuni* (Tamil)¹³. *Trianthema decandra* has been used in various parts of Asia, Africa, Australia and South America for curing various diseases. In some African countries, the plant is in popular use for treating skin diseases, asthma, hepatitis, wounds, fever and toothaches. In India, it is used for the treatment of ophthalmic¹⁴. The root applied to the eye cures corneal ulcers, itching, dimness of sight and night blindness. In orchitis, the root is ground in milk and administered¹⁵. The bitter roots are used for curing bacterial infections and it is also given in combination with ginger as a cathartic. The leaves contain huge amounts of vitamin C, and used to treat edema. The juice of leaves is used to treat the black quarter. The decoction of the herb is used as a vermifuge and is useful in rheumatitis. It is also used as an antidote to alcoholic poison.

The present work was carried out to evaluate the antioxidant and antimicrobial activities of the extracts of *Trianthema decandra* roots and leaves.

MATERIALS AND METHODS

Plant materials and extraction

The plant *Trianthema decandra* was collected from Namakkal district, Tamil Nadu. The plant was taxonomically identified and authenticated by the Botanical survey of India, Coimbatore (Tamil Nadu) and a voucher specimen was deposited for future reference. Around 250 g of the roots and leaves (separately) were dried in shade, pulverized by a mechanical grinder and passed through a 40-mesh sieve to get a fine powder and stored in an airtight container. The dried powder (25 g) was extracted with petroleum ether (60-80 °C), wax removed and then extracted with ethyl acetate, 80% methanol and water sequentially in a soxhlet apparatus. The solvents from various extracts were then concentrated in rotary evaporator at reduced pressure below 40 °C.

Chemicals

1,1-diphenyl-2-picryl-hydrazyl (DPPH), butylated hydroxy toluene (BHT) and catechol were purchased from Sigma, St. Louis, MO, USA. All other chemicals and reagents were of analytical grade.

Bacterial strains

To study the antibacterial activity of various extracts of *Trianthema decandra* roots and leaves, the strains of bacteria were obtained from Microbiology Laboratory at PSG Hospitals, Coimbatore. The selected bacteria included *Staphylococcus epidermis*, *Escherichia coli*, *Klebsiella pneumonia*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Pseudomonas aeruginosa*, *Pseudomonas putida* and *Proteus vulgaris*.

Antioxidant activity

Determination of total phenolic content

Total phenol content was determined by the method adapted from Singleton and Rossi¹⁶ with some modifications using the Folin-Ciocalteu reagent. 1 ml of the extract was mixed with 1 ml of Folin-Ciocalteu's phenol reagent (1:10). After 3 min, 1 ml of saturated sodium carbonate (35%) was added to the mixture and it was made up to 10 ml by adding deionised distilled water. The mixture was kept for 90 min at room temperature in the dark. The absorbance was measured at 725 nm against the blank. The total phenolic content is expressed as mg of Catechol equivalents (CE) per g of dry extract.

Determination of DPPH free radical scavenging activity

The scavenging effect of the extracts on DPPH radicals was determined according to the method adapted from Shimada *et al.*¹⁷. Various concentrations of sample (1.5 ml) were mixed with 3 ml of

200 μM DPPH solution. The mixture was shaken vigorously and allowed to stand for 40 min, and the absorbance was measured at 517 nm, using butylated hydroxy toluene (BHT) as the standard. The percentage of inhibition was calculated according to the formula: $[(A_{\text{control}} - A_{\text{sample}})/A_{\text{control}}] \times 100$. IC_{50} values were determined by linear regression.

Determination of Antibacterial activity by disc diffusion method

The *in vitro* antibacterial activity of the sample solutions was studied by disc diffusion method. Plates were prepared by pouring 20 ml of sterile nutrient agar (Hi-media) into sterile Petri dishes and were inoculated with a loopful broth culture of each organism. Sterile discs (6 mm dia) impregnated with 20 μl (1.5 mg/disc) various extracts dissolved in dimethyl sulphoxide (DMSO) were air-dried and placed on the agar plates. The plates were incubated at 37 $^{\circ}\text{C}$ for 24 h. Control studies with polymixin-B and rifampicin 30 μg were used as antibacterial standard drugs. The control experiments with solvent DMSO were done concurrently¹⁸. The results were recorded by measuring the zones of growth inhibition surrounding the disc. Clear inhibition zones around the discs indicate the presence of

antibacterial activity. All data regarding antibacterial activity are the average of triplicate analyses.

RESULTS AND DISCUSSION

Antioxidant activity

Total phenolic content

The total phenolic content of the various extracts of *Trianthema decandra* roots and leaves is given in Table 1. The ethyl acetate and methanolic extracts of both the roots and leaves showed higher levels of total phenolic contents than the petroleum ether and aqueous extracts. Phenols are very important plant constituents because of their scavenging ability due to their hydroxyl groups¹⁹. The imbalance between reactive oxygen species (ROS) and antioxidant defense mechanism leads to oxidative modifications in cellular membrane or intracellular molecules. Under pathological conditions, ROS are overproduced and results in lipid peroxidation and oxidative stress²⁰. Antioxidant-based drugs or formulations are used for the treatment of complex diseases like atherosclerosis, stroke, diabetes, Alzheimer's disease and cancer²¹.

Table 1: Total phenolic content ($\mu\text{g/ml}$) of the extracts of *Trianthema decandra* roots and leaves

Extract	Petroleum ether	Ethyl acetate	Methanol	Aqueous
<i>Trianthema decandra</i> roots	5.40 + 0.15	32.50 + 1.00	40.00 + 1.20	5.00 + 0.15
<i>Trianthema decandra</i> leaves	5.00 + 0.15	31.5 + 0.95	45.00 + 1.35	20.00 + 0.60

Values are expressed as mean + SD (n=3) as Catechol equivalents (CE).

DPPH free radical scavenging activity

The DPPH is considered to be a model of lipophilic radical. A chain reaction in lipophilic radicals was initiated by lipid autooxidation. The radical scavenging activity of *Trianthema decandra* roots and leaves extracts was determined from decrease in absorbance at 517 nm due to scavenging of stable DPPH free radical. The scavenging effects of *Trianthema decandra* roots and leaves extracts and BHT are

given in Table 2. The extracts had significant scavenging effects on the DPPH radical. The positive DPPH test suggests that the samples are free radical scavengers. The IC_{50} values of the extracts in the DPPH free radical scavenging assay are illustrated in the Fig 1. These results indicate that the *Trianthema decandra* roots and leaves extracts, particularly the ethyl acetate and methanolic extracts, exhibit the ability to quench the DPPH radical, suggesting that the extracts are good antioxidants with radical scavenging activity.

Table 2: DPPH free radical scavenging assay of *Trianthema decandra* roots and leaves

Extract	% Inhibition			
	Petroleum ether	Ethyl acetate	Methanol	Aqueous
<i>Trianthema decandra</i> roots (1 mg/ml)	0.95 + 0.03	92.79 + 2.80	54.95 + 1.65	12.06 + 0.35
<i>Trianthema decandra</i> leaves (1 mg/ml)	46.85 + 1.40	61.26 + 1.85	81.08 + 2.45	77.48 + 2.30

Values are expressed as mean + SD (n=3)




Fig. 1: IC_{50} values for DPPH free radical scavenging activity of *Trianthema decandra* roots and leaves

TDR - *Trianthema decandra* roots extracts

TDL - *Trianthema decandra* leaves extracts

PE - Petroleum ether extract

EA - Ethyl acetate extract

ME - Methanolic extract

AQ - Aqueous extract

Phenolic compounds may contribute directly to the antioxidative action²². The results indicate a strong relationship between total phenolic contents and radical scavenging activity, suggesting that phenolic compounds are responsible for the antioxidative properties of *Trianthema decandra* roots and leaves extracts. Phenolic compounds are reported to be effective hydrogen donors, which makes them good antioxidants^{23,24}.

Antibacterial activity

The antibacterial activities of various plants have been reported by many researchers²⁵. As the plants produce secondary metabolites in order to protect themselves from microorganisms, herbivores and insects, this antimicrobial effect is somehow expected from plants. Plants are important sources of potentially useful structures for the development of new chemotherapeutic agents. The first step towards this goal is the *in vitro* antibacterial activity assay²⁶. Many reports are available on the antibacterial properties of plants^{27, 28}.

In the present study, the methanolic extract of *Trianthema decandra* roots was found to be effective against *S. epidermis*, *K. pneumonia*, *S.*

aureus, *E. faecalis*, and *P. vulgaris*, while the ethyl acetate extract of roots was active against *K. pneumonia*, *S. aureus*, *P. aeruginosa* and *P. vulgaris* (Table 3). But petroleum ether extract of the roots was found to be effective against *K. pneumonia*, *S. aureus* and *P. vulgaris*, while aqueous extract was active only against *S. aureus*. Of the extracts of *Trianthema decandra* leaves, the ethyl acetate extract of *Trianthema decandra* showed inhibitory activity against all the bacteria tested. The methanolic extract of leaves was found to be active against all other organisms except *S. epidermis* and *E. coli*. While the aqueous extract of leaves showed mild activity against *P. aeruginosa*, the petroleum ether extract was found to be ineffective against any of the bacteria tested.

Table 3: Antibacterial activity of *Trianthema decandra* roots and leaves

Extracts	Zone of inhibition (mm dia)									
	Petroleum ether		Ethyl acetate		Methanol		Aqueous		Standards	
Organisms	TDR	TDL	TDR	TDL	TDR	TDL	TDR	TDL	PM	RF
<i>Staphylococcus epidermis</i>	-	-	-	25	10	-	-	-	30	18
<i>Escherichia coli</i>	-	-	-	25	-	-	-	-	28	11
<i>Klebsiella pneumonia</i>	10	-	14	20	20	22	-	-	28	30
<i>Staphylococcus aureus</i>	16	-	20	26	20	20	22	-	38	25
<i>Enterococcus faecalis</i>	-	-	-	20	14	30	-	-	35	30
<i>Pseudomonas aeruginosa</i>	-	-	8	30	-	20	-	10	10	18
<i>Pseudomonas putida</i>	-	-	-	22	-	8	-	-	36	10
<i>Proteus vulgaris</i>	18	-	20	20	20	8	-	-	24	4

Values are expressed as mean (n=3)

TDR - *Trianthema decandra* roots extracts

TDL - *Trianthema decandra* leaves extracts

PM - Polymyxin - B

RF - Rifampicin

The ethyl acetate and methanolic extracts of both the roots and leaves of *Trianthema decandra* showed good antibacterial activity against most of the tested bacteria, suggesting the presence of antibacterial agents in the plant.

Taking into account the total phenolic contents, DPPH free radical scavenging and antibacterial activities, the ethyl acetate and methanolic extracts of both the roots and leaves of *Trianthema decandra* showed higher levels compared to the petroleum ether and aqueous extracts. Antioxidants that retard the oxidation process may additionally exhibit antimicrobial activity²⁹. Thus there is correlation between the antibacterial activity and the antioxidant activity of the *Trianthema decandra* extracts. Our results suggested that both the roots and leaves of *Trianthema decandra* could be a promising source of natural antioxidants, as in ripe *Pepino furit*³⁰. Some medicinal plants are potent antioxidants and may be efficient as preventive agents in many diseases. The World Health Organization has also recognized the importance of traditional medicine and has created strategies, guidelines and standards for botanical medicines. Proven agro-industrial technologies need to be applied to the cultivation and processing of medicinal plants and the manufacture of herbal medicines³¹.

CONCLUSION

The present results reveal that the ethyl acetate and methanolic extracts of roots and leaves of *Trianthema decandra* possess antioxidant and antibacterial potential, by virtue of their phenolic contents, DPPH free radical scavenging and antibacterial activities. The results also revealed that the leaves have more antioxidant and antibacterial potential than the roots. The broad spectrum antibacterial activity of the above extracts may not be due to a single phytomolecule, but may be due to the presence of a number of bioactive metabolites.

ACKNOWLEDGEMENT

The authors are thankful to the management, K.S. Rangasamy College of Technology, Tiruchengode, India, for providing the fund and facilities.

REFERENCES

- Bushra Begum R, Ganga Devi T. Antibacterial activity of selected seaweeds from Kovalam southwest coast of India. Asian J Micorbiol Biotech Env Sci 2003; 5: 319-322.
- Bonjar GHS, Farrokhi PR. Anti-bacillus activity of some plants used in traditional medicine of Iran. Niger J Nat Prod Med 2004; 8: 34-39.
- Lampart-Szczapa E, Korczak J, Nogala-Kalucka M, Zawirska-Wojtasiak R. Antioxidant properties of lupin seed products. Food Chem 2003; 83: 279-285.
- Kris-Etherton PM, Hecker KD, Bonanome A, Coval SM, Binkoski AE, Hilpert KF, Griel AE, Etherton TD. Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. Am J Med 2002; 113: 71-88.
- Harborne JB. Phytochemical methods: a guide to modern techniques of plant analysis. 3rd ed. Bury, St. Edmunds, Suffolk: St. Edmundsbury press; 1998. p. 40.
- Mojzisova G, Kuchts M. Minireview: Dietary flavonoids and risk of coronary heart disease. Physiol Res 2001; 50: 529-536.
- Connor AM, Luby JJ, Tong CBS, Fin CE, Hancock JF. Genotypic and environmental variation in antioxidant activity, total phenolic content and anthocyanin content among blueberry cultivars. J Am Soc Hort Sci 2002; 127: 89-97.
- Ikram M, Inamul H. Screening of medicinal plants for antimicrobial activities. Fitoterapia 1984; 55: 62-64.
- Almagboul, AZ, Bashir AK, Farouk A, Salih AKM. Antimicrobial activity of certain Sudanese plants used in folklore medicine. Screening for antibacterial activity. Fitoterapia 1985; 56: 331-337.
- Izzo AA, Di Carlo G, Biscardi D, Fusco R, Macolo N, Boreli F, Capasso F, Fasulo MP, Autore G. Biological screening of Italian medicinal plants for antibacterial activity. Phtother Res 1995; 9: 281-286.
- Jansen AM, Cheffer JJC, Svendsen AB. Antimicrobial activity of essential oils: a 1976-1986 literature review. Aspects of test methods. Planta Med 1987; 40: 395-398.
- Scalbert A. Antimicrobial properties of tannins. Phytochem 1991; 30: 3875-3883.
- Kirtikar KR, Basu BD. Indian medicinal plants. Vol 2. Dehradun: International Book Distributors; pp. 1180-1181.
- Upadhyay OP, Kumar K, Tiwari KA. Ethno botanical study of skin treatment uses of medicinal plants of Bihar. Pharm Biol 1998; 36: 167-172.

15. Naovi SAH, Khan MSY, Vohora SB. Antibacterial, antifungal and anthelmintic investigations on Indian medicinal plants. *Fitoterapia* 1991; 62: 221.
16. Singleton VL, Rossi JA. Colorimetry of total phenolics with phosphotungstic acid reagents. *Am J Enol Vitic* 1965; 16: 144-158.
17. Shimada K, Fujikawa K, Yahara N, Nakamura T. Antioxidative properties of xanthan on the autooxidation of soybean oil in cyclodextrin emulsion. *J Agric Food Chem* 1992; 40: 945-948.
18. Umadevi S, Mohanta GP, Chelladurai V, Manna PK, Manavalan R. Antibacterial and antifungal activity of *Andrographis echinodes*. *J Nat Remedies* 2003; 53: 185-188.
19. Diplock AT. Will the 'good fairies' please prove to us that vitamin E lessens human degenerative diseases? *Free Radical Res.* 1997; 27: 511-532.
20. El-Habit, OHM, Saada HN, Azab KS, Abdel-Rahman M, El-Malah DF. The modifying effect of beta-carotene on gamma radiation-induced elevation of oxidative reactions and genotoxicity in male rats *Mutat Res* 2000; 466: 179-186.
21. Davies MJ. The oxidative environment and protein damage. *Biochim Biophys Acta* 2005; 1703: 93-109.
22. Hatano T, Edamatsu R, Hiramatsu M, Mori A, Fujita Y, Yashuara D. Effects of interaction of tannins with co-existing substances. VI. Effects of tannins and related polyphenols on superoxide anion radical and on DPPH radical. *Chem Pharm Bull* 1989; 37: 2016-2021.
23. Rice-Evans CA, Miller NJ, Bowell PG, Bramley PM, Pridhan JB. The relative antioxidant activity of plant derived polyphenolic flavonoids. *Free Radical Res* 1995; 22: 375-383.
24. Shahidi F, Naczk, M. Method of analysis and quantification of phenolic compounds. Lanchester, PA, USA: Technomic Publishing Company. 1995. pp. 287-293.
25. Cowan MM. Plant products as antimicrobial agents. *Clin Microbiol Rev* 1999; 22: 564-582.
26. Tona L, Kambu K, Ngimbi N, Cimanga K, Vlietinck AJ. Antiamoebic and phytochemical screening of some Congolese medicinal plants. *J Ethnopharmacol* 1998; 61: 57-65.
27. Samy RP, Ignacimuthu S. Antibacterial activity of some folklore medicinal plants used by tribes in Western Ghats in India. *J Ethnopharmacol* 2000; 69: 63-71.
28. Palombo EA, Semple SJ. Antibacterial activity of traditional medicinal plants. *J Ethnopharmacol* 2001; 77: 151-157.
29. Puupponen-Pimia R, Nohynek L, Meilier C, Kahkonen M, Heinonen M, Hopia A, Oksman-Caldentey KM. Antimicrobial properties of phenolic compounds from berries. *J Appl Microbiol* 2001; 90: 494-507.
30. Sudha G, Sangeetha Priya M, Indhu Shree R, Vadivukkarasai S. Antioxidant activity of ripe Pepino fruit (*Solanum muricatum* Aiton). *Int J Pharm Pharm Sci* 2011; 3: 257-261.
31. Rout SP, Choudary KA, Kar DM, Das L, Jain A. Plants in traditional medicinal system - future source of new drugs. *Int J Pharm Pharm Sci* 2009; 1:1-23.