



Bibliography

- Abruzzese, G. A., Cerrone, G. E., Gamez, J. M., Graffigna, M. N., Belli, S., Lioy, G., Mormandi, E., Otero, P., Levalle, O. A., & Motta, A. B. (2017). Lipid Accumulation Product (LAP) and Visceral Adiposity Index (VAI) as Markers of Insulin Resistance and Metabolic Associated Disturbances in Young Argentine Women with Polycystic Ovary Syndrome. *Hormone and metabolic research = Hormon- und Stoffwechselforschung = Hormones et metabolisme*, 49(1), 23–29. <https://doi.org/10.1055/s-0042-113463>
- Agrawal, H., Aggarwal, K., & Jain, A. (2019). Visceral adiposity index: Simple Tool for assessing cardiometabolic risk in women with polycystic ovary syndrome. *Indian Journal of Endocrinology and Metabolism*, 23(2), 232–237. <https://doi.org/10.4103/ijem.IJEM-559-18>
- Ahmed, N., Adam, S. I. M., Nawi, A. M., Hassan, M. R., & Ghazi, H. F. (2016). Abdominal obesity indicators: Waist circumference or waist-to-hip ratio in Malaysian adults population. *International Journal of Preventive Medicine*, 2016(June). <https://doi.org/10.4103/2008-7802.183654>
- Ahn, N., Baumeister, S. E., Amann, U., Rathmann, W., Peters, A., Huth, C., ... Meisinger, C. (2019). Visceral adiposity index (VAI), lipid accumulation product (LAP), and product of triglycerides and glucose (TyG) to discriminate prediabetes and diabetes. *Scientific Reports*, 9(1), 1–11. <https://doi.org/10.1038/s41598-019-46187-8>
- Ainsworth, B. E., Haskell, W. L., Herrmann, S. D., Meckes, N., Bassett, D. R., Tudor-Locke, C., ... Leon, A. S. (2011). 2011 compendium of physical activities: A second update of codes and MET values. *Medicine and Science in Sports and Exercise*, 43(8), 1575–1581. <https://doi.org/10.1249/MSS.0b013e31821ece12>
- Akdemir, A. O., Karabakan, M., Aktas, B. K., Bozkurt, A., Ozgur, E. G., Akdogan, N., & Yaris, M. (2019). Visceral adiposity index is useful for evaluating obesity effect on erectile dysfunction. *Andrologia*, 51(6), e13282.

- Akseer, N., Mehta, S., Wigle, J., Chera, R., Brickman, Z. J., Al-Gashm, S., Akter, R., Nessa, A., Husain, M. F., Wahed, F., Khatun, N., Yesmin, M., Nasreen, S., & Tajkia, T. (2020). Effect of Obesity on Fasting Blood Sugar. *Mymensingh medical journal: MMJ*, 26(1), 7–11.
- Akter, R., Nessa, A., Husain, M. F., Wahed, F., Khatun, N., Yesmin, M., Nasreen, S., & Tajkia, T. (2017). Effect of Obesity on Fasting Blood Sugar. *Mymensingh medical journal : MMJ*, 26(1), 7–11.
- Alexander, K. E., Ventura, E. E., Spruijt-Metz, D., Weigensberg, M. J., Goran, M. I., & Davis, J. N. (2009). Association of breakfast skipping with visceral fat and insulin indices in overweight Latino youth. *Obesity*, 17(8), 1528–1533. <https://doi.org/10.1038/oby.2009.127>
- Aljaffar, A. G. Y. (2018). A Study of Serum Lipid Profile among Obese and Non Obese Individuals: A Hospital based Study from Karbala, Iraq. *International Journal of Contemporary Medical Research [IJCMR]*, 5(4), 17–20. <https://doi.org/10.21276/ijcmr.2018.5.4.11>
- Alkhalaqi, A., Al-Naimi, F., Qassmi, R., Shi, Z., Ganji, V., Salih, R., & Bawadi, H. (2020). Visceral adiposity index is a better predictor of type 2 diabetes than body mass index in Qatari population. *Medicine*, 99(35).
- Amato, M. C., & Giordano, C. (2014). Visceral adiposity index: an indicator of adipose tissue dysfunction. *International journal of endocrinology*, 2014, 730827. <https://doi.org/10.1155/2014/730827>
- Amato, M. C., Giordano, C., Galia, M., Criscimanna, A., Vitabile, S., Midiri, M., & Galluzzo, A. (2010). Visceral adiposity index: A reliable indicator of visceral fat function associated with cardiometabolic risk. *Diabetes Care*, 33(4), 920–922. <https://doi.org/10.2337/dc09-1825>
- Anyanwu, G. E., Ekezie, J., Danborn, B., & Ugochukwu, A. I. (2010). *Impact of education on obesity and blood pressure in developing countries : A study on the Ibos of Nigeria*. 2(7). <https://doi.org/10.4297/najms.2010.2320>
- Aparicio, A., Rodríguez-Rodríguez, E. E., Aranceta-Bartrina, J., Gil, Á., González-Gross, M., Serra-Majem, L., ... Ortega, R. M. (2017). Differences in meal patterns and timing with regard to central obesity in the ANIBES

- (Anthropometric data, macronutrients and micronutrients intake, practice of physical activity, socioeconomic data and lifestyles in Spain') Study. *Public Health Nutrition*, 20(13), 2364–2373. <https://doi.org/10.1017/S1368980017000635>
- Arimura ST, Moura BM, Pimentel GD, Silva ME, Sousa MV. (2011). Waist circumference is better associated with high density lipoprotein (HDL-c) than with body mass index (BMI) in adults with metabolic syndrome. *Nutr Hosp*. Nov-Dec;26(6):1328-32. doi: 10.1590/S0212-16112011000600020. PMID: 22411379.
- Ashwell, M., & Gibson, S. (2014). *A proposal for a primary screening tool: 'Keep your waist circumference to less than half your height.'* 1–6.
- Ayundini, G., Astrella, C., Tahapary, D., & Soewondo, P. (2019). A systematic review on the association between lipid accumulation product index and type 2 diabetes mellitus. *Journal of the ASEAN Federation of Endocrine Societies*, 34(1), 16.
- Bacopoulou, F., Efthymiou, V., Landis, G., Rentoumis, A., & Chrousos, G. P. (2015). Waist circumference, waist-to-hip ratio and waist-to-height ratio reference percentiles for abdominal obesity among Greek adolescents. *BMC Pediatrics*, 15(1), 1–9. <https://doi.org/10.1186/s12887-015-0366-z>
- Banerjee, R., Ray, K., Bhattacharjee, S., Guha, S., Banerjee, I., & Nath, I. (2014). A Study of Insulin Resistance and its Clinico. Metabolic Associations Among Apparently Healthy Individuals Attending a Tertiary Care Hospital. *Annals of medical and health sciences research*, 4(5), 823-828.
- Bhosle, D. D. (2016). Homeostasis Model Assessment of Insulin Resistance (HOMA-IR) in the Diagnosis of Insulin Resistance and Prediabetes. *Journal of Medical Science And Clinical Research*, 04(09), 12705–12710. <https://doi.org/10.18535/jmscr/v4i9.65>
- Biyik, Z., & Guney, I. (2019). Lipid accumulation product and visceral adiposity index: two new indices to predict metabolic syndrome in chronic kidney disease. *European Review for Medical and Pharmacological Sciences*, 23(5), 2167–2173. https://doi.org/10.26355/eurrev_201903_17262

- Bonnie, R., Stroud, C., & Breiner, H. (2015). Young adults in the 21st century. In *Investing in the health and well-being of young adults*. Retrieved from https://www.ncbi.nlm.nih.gov/books/NBK284787/pdf/Bookshelf_NBK284787.pdfWorld Health Organisation, WHO, 2002
- Borrueal, S., Moltó, J. F., Alpañés, M., Fernández-Durán, E., Álvarez-Blasco, F., Luque-Ramírez, M., & Escobar-Morreale, H. F. (2014). Surrogate markers of visceral adiposity in young adults: Waist circumference and body mass index are more accurate than waist hip ratio, model of adipose distribution and visceral adiposity index. *PLoS ONE*, 9(12), 1–17. <https://doi.org/10.1371/journal.pone.0114112>
- Bozorgmanesh, M., Hadaegh, F., & Azizi, F. (2011). Predictive performance of the visceral adiposity index for a visceral adiposity-related risk: Type 2 Diabetes. *Lipids in Health and Disease*, 10(1), 88. <https://doi.org/10.1186/1476-511X-10-88>
- Bray, G. A. (2004). Medical consequences of obesity. *Journal of Clinical Endocrinology and Metabolism*, 89(6), 2583–2589. <https://doi.org/10.1210/jc.2004-0535>
- Bray, G. A., & Popkin, B. M. (1998). Dietary fat intake does affect obesity. *American Journal of Clinical Nutrition*, 68(6), 1157–1173. <https://doi.org/10.1093/ajcn/68.6.1157>
- Brisson, D., Perron, P., Guay, S. P., Gaudet, D., & Bouchard, L. (2010). The hypertriglyceridemic waist” phenotype and glucose intolerance in pregnancy. *Cmaj*, 182(15). <https://doi.org/10.1503/cmaj.100378>
- Browning, L. M., Hsieh, S. D., & Ashwell, M. (2010). A systematic review of waist-to-height ratio as a screening tool for the prediction of cardiovascular disease and diabetes: 05 could be a suitable global boundary value. *Nutrition Research Reviews*, 23(2), 247–269. <https://doi.org/10.1017/S0954422410000144>
- Byambasukh, O., Vinke, P., Kromhout, D., Navis, G., & Corpeleijn, E. (2021). Physical activity and 4-year changes in body weight in 52,498 non-obese people: the Lifelines cohort. *International Journal of Behavioral Nutrition and Physical Activity*, 18(1), 1–11. <https://doi.org/10.1186/s12966-021-01141-8>

- Cameron, A. J., Magliano, D. J., Shaw, J. E., Zimmet, P. Z., Carstensen, B., Alberti, K. G. M. M., ... Barr, E. L. M. (2012). *The influence of hip circumference on the relationship between abdominal obesity and mortality*. (January), 484–494. <https://doi.org/10.1093/ije/dyr198>
- Canepa, M., Strait, J. B., Milaneschi, Y., AlGhatrif, M., Ramachandran, R., Makrogiannis, S., ... Ferrucci, L. (2013). The relationship between visceral adiposity and left ventricular diastolic function: Results from the Baltimore Longitudinal Study of Aging. *Nutrition, Metabolism and Cardiovascular Diseases*, 23(12), 1263–1270. <https://doi.org/10.1016/j.numecd.2013.04.003>
- Cartolano, F. D. C., Pappiani, C., Freitas, M. C. P. D., Figueiredo, A. M., Carioca, A. A. F., & Damasceno, N. R. T. (2018). Is lipid accumulation product associated with an Atherogenic lipoprotein profile in Brazilian subjects?. *Arquivos brasileiros de cardiologia*, 110, 339-347.
- Castro, A. V., Kolka, C. M., Kim, S. P., & Bergman, R. N. (2014). Obesity, insulin resistance and comorbidities? Mechanisms of association. *Arquivos brasileiros de endocrinologia e metabologia*, 58(6), 600–609. <https://doi.org/10.1590/0004-2730000003223>
- Cefalu, W. T. (2001). Insulin resistance: cellular and clinical concepts. *Experimental biology and medicine*, 226(1), 13-26.
- Census of India. (2011). *District census handbook, SIKKIM. SERIES-12,*
- Chait, A., & den Hartigh, L. J. (2020). Adipose Tissue Distribution, Inflammation and Its Metabolic Consequences, Including Diabetes and Cardiovascular Disease. *Frontiers in Cardiovascular Medicine*, 7(February), 1–41. <https://doi.org/10.3389/fcvm.2020.00022>
- Chan, R. S. M., & Woo, J. (2010). Prevention of overweight and obesity: How effective is the current public health approach. *International Journal of Environmental Research and Public Health*, 7(3), 765–783. <https://doi.org/10.3390/ijerph7030765>
- Chandalia, M., Davila, H., Pan, W., Szuszkiewicz, M., Tuvdendorj, D., Livingston, E. H., & Abate, N. (2012). Adipose tissue dysfunction in humans: A potential role for the transmembrane protein ENPP1. *Journal of*

- Clinical Endocrinology and Metabolism*, 97(12), 4663–4672.
<https://doi.org/10.1210/jc.2012-2018>
- Chaffin, M., Hairston, K. G., Hanley, A. J., Tooze, J. A., Norris, J. M., Palmer, N. D., ... & Wagenknecht, L. E. (2021). Association of Visceral Adipose Tissue and Insulin Resistance with Incident Metabolic Syndrome Independent of Obesity Status: The IRAS Family Study. *Obesity*.
- Chaudhari, A., Gujarathi, S., & Bhatia, G. (2020). Comparison of blood glucose in obese and non-obese students in a medical college. *International Journal of Research in Medical Sciences*, 8(11), 3916. <https://doi.org/10.18203/2320-6012.ijrms20204877>
- Chen, H. Y., Chiu, Y. L., Chuang, Y. F., Hsu, S. P., Pai, M. F., Yang, J. Y., & Peng, Y. Sen. (2014). Visceral adiposity index and risks of cardiovascular events and mortality in prevalent hemodialysis patients. *Cardiovascular Diabetology*, 13(1), 1–9. <https://doi.org/10.1186/s12933-014-0136-5>
- Cheng, C. H., Ho, C. C., Yang, C. F., Huang, Y. C., Lai, C. H., & Liaw, Y. P. (2010). Waist-to-hip ratio is a better anthropometric index than body mass index for predicting the risk of type 2 diabetes in Taiwanese population. *Nutrition Research*, 30(9), 585–593. <https://doi.org/10.1016/j.nutres.2010.08.007>
- Chiang, J. K., & Koo, M. (2012). Lipid accumulation product: a simple and accurate index for predicting metabolic syndrome in Taiwanese people aged 50 and over. *BMC cardiovascular disorders*, 12(1), 1-6.
- Chille et.al. (2017) laboratory medicine, J. Bras. Patol. Med. Lab. 53 (2) Mar-Apr 2017 <https://doi.org/10.5935/1676-2444.20170012>
- Ching, Y. K., Chin, Y. S., Appukutty, M., Gan, W. Y., & Chan, Y. M. (2020). Comparisons of conventional and novel anthropometric obesity indices to predict metabolic syndrome among vegetarians in Malaysia. *Scientific Reports*, 10(1), 1–13. <https://doi.org/10.1038/s41598-020-78035-5>
- Costa da, M. A. P., Vasconcelos, A. G. G., & da Fonseca, M. de J. M. (2015). Prevalência de obesidade, excesso de peso e obesidade abdominal e associação com prática de atividade física em uma universidade federal. *Revista Brasileira de Epidemiologia*, 17(2), 421–436. <https://doi.org/10.1590/1809-4503201400020011ENG>

- Curtis, R. G., Olds, T., Ferguson, T., Frayse, F., Dumuid, D., Esterman, A., ... Maher, C. A. (2021). Changes in diet, activity, weight, and wellbeing of parents during COVID-19 lockdown. *PLoS ONE*, *16*(3 March), 1–14. <https://doi.org/10.1371/journal.pone.0248008>
- Davis, (2021); <https://www.medicinenet.com/adiposity/definition.htm>
- Dev. M, N. (2016). Is Lipid Accumulation Product (LAP), a better obesity index in diagnosing cardiovascular disease? *TC (Mg/DL)*, *2*(6), 12–14. Retrieved from <https://pdfs.semanticscholar.org/a6a8/627759f52ce7b82bb4a974f8003c8e845ad2.pdf>
- Dhawan, D., & Sharma, S. (2020). Abdominal obesity, adipokines and non-communicable diseases. *The Journal of steroid biochemistry and molecular biology*, 105737.
- Dong, H., Xu, Y., Zhang, X., & Tian, S. (2017). Visceral adiposity index is strongly associated with hyperuricemia independently of metabolic health and obesity phenotypes. *Scientific reports*, *7*(1), 1-13
- Donnelly, J. E., Honas, J. J., Smith, B. K., Mayo, M. S., Gibson, C. A., Sullivan, D. K., Lee, J., Herrmann, S. D., Lambourne, K., & Washburn, R. A. (2013). Aerobic exercise alone results in clinically significant weight loss for men and women: midwest exercise trial 2. *Obesity (Silver Spring, Md.)*, *21*(3), E219–E228. <https://doi.org/10.1002/oby.20145>
- Du, T., Yu, X., Zhang, J., & Sun, X. (2015). Lipid accumulation product and visceral adiposity index are effective markers for identifying the metabolically obese normal-weight phenotype. *Acta Diabetologica*, *52*(5), 855–863. <https://doi.org/10.1007/s00592-015-0715-2>
- Dua, S., Bhuker, M., Sharma, P., Dhall, M., & Kapoor, S. (2014). Body mass index relates to blood pressure among adults. *North American Journal of Medical Sciences*, Vol. 6, pp. 89–95. <https://doi.org/10.4103/1947-2714.127751>
- Dutta, M., Selvamani, Y., Singh, P., & Prashad, L. (2019). The double burden of malnutrition among adults in India: evidence from the National Family Health Survey-4 (2015-16). *Epidemiology and health*, *41*, e2019050. <https://doi.org/10.4178/epih.e2019050>.
- Eastham and R.D. (1985): *Biochemical values in clinical Medicine*, 7th Ed. Bristol. England. John Wright & Sons, Ltd.

- Easwaran, M. R., Sivasubramanian, P., & Kannan, G. (2019). *Prevalence of central obesity and its association with sociodemographic profile among young adults attending Outdoor Patient Department of Community Health Centre in Madurai, Tamil Nadu*. 8(12), 1023–1028. <https://doi.org/10.5455/ijmsph.2019.0719721092019>
- Ebbert, J. O., & Jensen, M. D. (2013). *Fat Depots, Free Fatty Acids, and Dyslipidemia*. (Cvd), 498–508. <https://doi.org/10.3390/nu5020498>
Changes in diet and lifestyle and long-term weight gain in women and men.
- EBCOG SCIENTIFIC COMMITTEE. (2014). Obesity and Women ' s Health. *Facts Views Vis OBGYN*, 6(2), 175–176.
- Egger, G., & Dixon, J. (2010). Inflammatory effects of nutritional stimuli: Further support for the need for a big picture approach to tackling obesity and chronic disease. *Obesity Reviews*, 11(2), 137–149. <https://doi.org/10.1111/j.1467-789X.2009.00644.x>
- Egger, G., & Dixon, J. (2014). Beyond obesity and lifestyle: A review of 21st century chronic disease determinants. *BioMed Research International*, 2014. <https://doi.org/10.1155/2014/731685>
- Egziabher Guebre, F., Alix, P. M., Koppe, L., Pelletier, C. C., Kalbacher, E., Fouque, D., & Soulage, C. O. (2013). Ectopic lipid accumulation: A potential cause for metabolic disturbances and a contributor to the alteration of kidney function. *Biochimie*, 95(11), 1971–1979. <https://doi.org/10.1016/j.biochi.2013.07.017>
- Elliot, C. A., & Hamlin, M. J. (2018). Combined diet and physical activity is better than diet or physical activity alone at improving health outcomes for patients in New Zealand's primary care intervention. *BMC public health*, 18(1), 230. <https://doi.org/10.1186/s12889-018-5152-z>
- Elmadfa, I. (2009). European Nutrition and Health Report 2009. Preface. *Forum of Nutrition*, 62.
- Engin, A. (2017). The definition and prevalence of obesity and metabolic syndrome. *Obesity and lipotoxicity*, 1-17

- Fahed, M., Abou Jaoudeh, M. G., Merhi, S., Mosleh, J. M. B., Ghadieh, R., Ghadieh, R., El Hayek Fares, J. E. (2020). Evaluation of risk factors for insulin resistance: A cross sectional study among employees at a private university in Lebanon. *BMC Endocrine Disorders*, 20(1), 1–14. <https://doi.org/10.1186/s12902-020-00558-9>
- Feinstein, B. L., Sabates, R., Anderson, T. M., Sorhaindo, A., & Hammond, C. (2006). 4 . *What are the effects of education on health ?* (December 2005), 171–354.
- Felisbino-Mendes, M. S., Cousin, E., Malta, D. C., Machado, Í. E., Ribeiro, A., Duncan, B. B., Schmidt, M. I., Silva, D., Glenn, S., Afshin, A., & Velasquez-Melendez, G. (2020). The burden of non-communicable diseases attributable to high BMI in Brazil, 1990-2017: findings from the Global Burden of Disease Study. *Population health metrics*, 18(Suppl 1), 18. <https://doi.org/10.1186/s12963-020-00219-y>
- Ferreira Nascimento, M. V., Rendo-Urteaga, T., Vilanova-Campelo, R. C., Carvalho, H. B., da Paz Oliveira, G., Paes Landim, M. B., & Torres-Leal, F. L. (2017). The lipid accumulation product is a powerful tool to predict metabolic syndrome in undiagnosed Brazilian adults. *Clinical Nutrition*, 36(6), 1693–1700. <https://doi.org/10.1016/j.clnu.2016.12.020>
- Fiorentino, T. V., Marini, M. A., Succurro, E., Andreozzi, F., & Sesti, G. (2019). Relationships of surrogate indexes of insulin resistance with insulin sensitivity assessed by euglycemic hyperinsulinemic clamp and subclinical vascular damage. *BMJ Open Diabetes Research and Care*, 7(1). <https://doi.org/10.1136/bmjdr-2019-000911>
- Fischer, K., Pick, J. A., Moewes, D., & Nöthlings, U. (2015). Qualitative aspects of diet affecting visceral and subcutaneous abdominal adipose tissue: A systematic review of observational and controlled intervention studies. *Nutrition Reviews*, 73(4), 191–215. <https://doi.org/10.1093/nutrit/nuu006>
- Fossati, P., & Prencipe, L. (1982). Serum triglycerides determined colorimetrically with an enzyme that produces hydrogen peroxide. *Clinical Chemistry*, 28(10), 2077–2080. <https://doi.org/10.1093/clinchem/28.10.2077>
- Frayn, K. N. (2000). Visceral fat and insulin resistance—causative or correlative?. *British Journal of Nutrition*, 83(S1), S71-S77.

- Freedland, E. S. (2004). Role of a critical visceral adipose tissue threshold (CVATT) in metabolic syndrome: Implications for controlling dietary carbohydrates: A review. *Nutrition and Metabolism*, 1, 1–24. <https://doi.org/10.1186/1743-7075-1-12>
- Fujioka, S., Matsuzawa, Y., Tokunaga, K., & Tarui, S. (1987). Contribution of intra-abdominal fat accumulation to the impairment of glucose and lipid metabolism in human obesity. *Metabolism*, 36(1), 54-59.
- Gan, W. Y., Boo, S., Seik, M. Y., & Khoo, H. E. (2018). Comparing the nutritional status of vegetarians and non-vegetarians from a Buddhist Organisation in Kuala Lumpur, Malaysia. *Malaysian Journal of Nutrition*, 24(1), 89–101.
- Ganguly, S., Ray, L., Kuruvila, S., Nanda, S. K., & Ravichandran, K. (2018). Lipid accumulation product index as visceral obesity indicator in psoriasis: a case–control study. *Indian journal of dermatology*, 63(2), 136.
- Global BMI Mortality Collaboration, Di Angelantonio, E., Bhupathiraju, S., Wormser, D., Gao, P., Kaptoge, S., Berrington de Gonzalez, A., Cairns, B. J., Huxley, R., Jackson, C., Joshy, G., Lewington, S., Manson, J. E., Murphy, N., Patel, A. V., Samet, J. M., Woodward, M., Zheng, W., Zhou, M., Bansal, N., ... Hu, F. B. (2016). Body-mass index and all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four continents. *Lancet (London, England)*, 388(10046), 776–786. [https://doi.org/10.1016/S0140-6736\(16\)30175-1](https://doi.org/10.1016/S0140-6736(16)30175-1)
- Goh, L. G. H., Dhaliwal, S. S., Welborn, T. A., Lee, A. H., & Della, P. R. (2014). Anthropometric measurements of general and central obesity and the prediction of cardiovascular disease risk in women: A cross-sectional study. *BMJ Open*, 4(2). <https://doi.org/10.1136/bmjopen-2013-004138>
- Goldani, H., Adami, F. S., Antunes, M. T., Rosa, L. H., Fassina, P., Grave, M. T. Q., & Dal Bosco, S. M. (2015). Aplicabilidad de la adiposidad visceral del índice VAI en la predicción de componentes del síndrome metabólico en ancianos. *Nutricion Hospitalaria*, 32(4), 1609–1615. <https://doi.org/10.3305/nh.2015.32.4.9589>
- Gonçalves, C. G., Glade, M. J., & Meguid, M. M. (2016). Metabolically healthy obese individuals: Key protective factors. *Nutrition*, 32(1), 14–20. <https://doi.org/10.1016/j.nut.2015.07.010>

- Gotter Ana, (2021), Danielle Hildreth, CPT; Updated on January 31, 2021 <https://www.healthline.com/health/visceral-fat#:~:text=It's%20the%20fat%20that%20you,1%20%2C%202%20%2C%203%20>).
- Goyal, G., Panag, K. M. S., & Kaur, N. (2014). Correlation of insulin resistance by various methods with fasting insulin in obese. *International Journal of Applied and Basic Medical Research*, 4(3), 41. <https://doi.org/10.4103/2229-516x.140733>
- Graham, H. (2007). *Unequal lives: health and socioeconomic inequalities*. McGraw-Hill Education (UK).
- Griffith, M. L., Younk, L. M., & Davis, S. N. (2010). Visceral adiposity, insulin resistance, and type 2 diabetes. *American Journal of Lifestyle Medicine*, 4(3), 230-243.
- Grover, S. A., Kaouache, M., Rempel, P., Joseph, L., Dawes, M., Lau, D. C., & Lowensteyn, I. (2015). Years of life lost and healthy life-years lost from diabetes and cardiovascular disease in overweight and obese people: a modelling study. *The lancet Diabetes & endocrinology*, 3(2), 114-122.
- Gudmundsdottir, H., Høiegggen, A., Stenehjem, A., Waldum, B., & Os, I. (2012). Hypertension in women: Latest findings and clinical implications. *Therapeutic Advances in Chronic Disease*, 3(3), 137–146. <https://doi.org/10.1177/2040622312438935>
- Guo, S. X., Zhang, X. H., Zhang, J. Y., He, J., Yan, Y. Z., Ma, J. L., ... Ding, Y. S. (2016). Visceral Adiposity and Anthropometric Indicators as Screening Tools of Metabolic Syndrome among Low Income Rural Adults in Xinjiang. *Scientific Reports*, 6(July), 1–8. <https://doi.org/10.1038/srep36091>
- Gupta R, Mehrishi S. (1997). Waist-hip ratio and blood pressure correlation in an urban Indian population. *J Indian Med Assoc*; 95:412-5
- Gutch, M., Kumar, S., Razi, S. M., Gupta, K., & Gupta, A. (2015). Assessment of insulin sensitivity/resistance. *Indian Journal of Endocrinology and Metabolism*, 19(1), 160–164. <https://doi.org/10.4103/2230-8210.146874>

- Harris B., "Public health, nutrition, and the decline of mortality: the McKeown thesis revisited," *Social History of Medicine*, vol. 17, no. 3, pp. 379–407, 2004.
- Hébert, J. R., Hurley, T. G., Steck, S. E., Miller, D. R., Tabung, F. K., Peterson, K. E., ... Frongillo, E. A. (2014). Considering the value of dietary assessment data in informing nutrition-related health policy. *Advances in Nutrition*, 5(4), 447–455. <https://doi.org/10.3945/an.114.006189>
- Heymsfield, S. B., & Wadden, T. A. (2017). Mechanisms, Pathophysiology, and Management of Obesity. *The New England journal of medicine*, 376(3), 254–266. <https://doi.org/10.1056/NEJMra1514009>
- Howel D (2012) Waist Circumference and Abdominal Obesity among Older Adults: Patterns, Prevalence and Trends. *PLoS ONE* 7(10): e48528. <https://doi.org/10.1371/journal.pone.0048528>
- Hsieh, C. J., Wang, P. W., & Chen, T. Y. (2014). The relationship between regional abdominal fat distribution and both insulin resistance and subclinical chronic inflammation in non-diabetic adults. *Diabetology and Metabolic Syndrome*, 6(1), 1–7. <https://doi.org/10.1186/1758-5996-6-49>
- Hsieh, S. D., & Muto, T. (2005). The superiority of waist-to-height ratio as an anthropometric index to evaluate clustering of coronary risk factors among non-obese men and women. *Preventive Medicine*, 40(2), 216–220. <https://doi.org/10.1016/j.ypmed.2004.05.025>
- Hsieh, S. D., Yoshinaga, H., & Muto, T. (2003). Waist-to-height ratio, a simple and practical index for assessing central fat distribution and metabolic risk in Japanese men and women. *International Journal of Obesity*, 27(5), 610–616. <https://doi.org/10.1038/sj.ijo.0802259>
- Huang, X., Wang, Q., Liu, T., Pei, T., Liu, D., Zhu, H., & Huang, W. (2019). Body fat indices as effective predictors of insulin resistance in obese/non-obese polycystic ovary syndrome women in the Southwest of China. *Endocrine*, 65(1), 81–85. <https://doi.org/10.1007/s12020-019-01912-1>
- Hwang, Y. C., Hayashi, T., Fujimoto, W. Y., Kahn, S. E., Leonetti, D. L., McNeely, M. J., & Boyko, E. J. (2015). Visceral abdominal fat accumulation predicts the conversion of metabolically healthy obese subjects to an unhealthy

- phenotype. *International Journal of Obesity*, 39(9), 1365–1370. <https://doi.org/10.1038/ijo.2015.75>
- Ilhan Anik, G., Yildizhan, B., & Pekin, T. (2019). The impact of lipid accumulation product (LAP) and visceral adiposity index (VAI) on clinical, hormonal and metabolic parameters in lean women with polycystic ovary syndrome. *Gynecological Endocrinology*, 35(3), 233–236. <https://doi.org/10.1080/09513590.2018.1519794>
- Indian Food Composition Table, RDA 2020 Longvah, Thingnganing & Ananthan, Rajendran & Bhaskar, K & Venkaiah, K. (2017). Indian Food Composition Tables.
- Indias Moreno, I., Oliva-Olivera, W., Omiste, A., Castellano-Castillo, D., Lhamyani, S., Camargo, A., & Tinahones, F. J. (2016). Adipose tissue infiltration in normal-weight subjects and its impact on metabolic function. *Translational Research*, 172, 6-17.e3. <https://doi.org/10.1016/j.trsl.2016.01.002>
- International Diabetes Federation, IDF, 2001
- International Diabetes Federation, IDF, 2006
- Ito S. (2019). High-intensity interval training for health benefits and care of cardiac diseases - The key to an efficient exercise protocol. *World journal of cardiology*, 11(7), 171–188. <https://doi.org/10.4330/wjc.v11.i7.171>
- Janghorbani, M., Amini, M., Rezvanian, H., Gouya, M. M., Delavari, A., Alikhani, S., & Mahdavi, A. (2008). Association of body mass index and abdominal obesity with marital status in adults. *Archives of Iranian medicine*, 11(3), 274–281.
- Janssen, I., Heymsfield, S. B., Allison, D. B., Kotler, D. P., & Ross, R. (2002). Body mass index and waist circumference independently contribute to the prediction of nonabdominal, abdominal subcutaneous, and visceral fat. *The American journal of clinical nutrition*, 75(4), 683–688. <https://doi.org/10.1093/ajcn/75.4.683>
- Johns, D. J., Hartmann-Boyce, J., Jebb, S. A., Aveyard, P., & Behavioural Weight Management Review Group (2014). Diet or exercise interventions vs combined behavioral weight management programs: a systematic review and meta-analysis of direct comparisons. *Journal of the Academy of*

- Nutrition and Dietetics*, 114(10), 1557–1568. <https://doi.org/10.1016/j.jand.2014.07.005>
- Joshi, B., Lakhan, T., Mukherji, S., Patil, A., & Unisa, S. (2018). Visceral adiposity index among young girls with PCOS and its association with phenotypes and metabolic risk. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*, 7(2), 513. <https://doi.org/10.18203/2320-1770.ijrcog20180164>
- Jung, I. S., Shin, C. M., Park, S. J., Park, Y. S., Yoon, H., Jo, H. J., ... & Lee, D. H. (2019). Association of visceral adiposity and insulin resistance with colorectal adenoma and colorectal cancer. *Intestinal research*, 17(3), 404.
- Kaba, Z., Khamisa, N., & Tshuma, N. (2017). Age-group differences in risk perceptions of non-communicable diseases among adults in Diepsloot township, Johannesburg, South Africa: A cross-sectional study based on the health belief model. *South African Medical Journal*, 107(9), 797–804. <https://doi.org/10.7196/SAMJ.2017.v107i9.12168>
- Kahn, H. S. (2005). The "lipid accumulation product" performs better than the body mass index for recognizing cardiovascular risk: a population-based comparison. *BMC cardiovascular disorders*, 5(1), 1-10.
- Kahn, H. S. (2006). Than BMI for Identifying Diabetes. *Cardiovascular and Metabolic Risk*, 29(1), 151–153.
- Kahn, H. S., & Valdez, R. (2003). Metabolic risks identified by the combination of enlarged waist and elevated triacylglycerol concentration. *The American journal of clinical nutrition*, 78(5), 928-934.
- Kasirajan, S. G. T., Dakshnamurthy, P., Nachiyar, I., Suthakaran, P. K., & Rajendran, K. (2017). Association of lipid accumulation product with Gestational Diabetes Mellitus: a hospital based study. *Annals of Woman and Child Health*, 3(2), A17–A21. <https://doi.org/10.21276/awch.1461>
- Kershaw, E. E., & Flier, J. S. (2004). Adipose tissue as an endocrine organ. *The Journal of Clinical Endocrinology & Metabolism*, 89(6), 2548-2556.
- Kesztyüs, D., Erhardt, J., Schönsteiner, D., & Kesztyüs, T. (2018). Therapeutic Treatment for Abdominal Obesity in Adults. *Deutsches Arzteblatt*

- international*, 115(29-30), 487–493
<https://doi.org/10.3238/arztebl.2018.0487>
- Khoury, M., Manhiot, C., Bs, C., & Mccrindle, B. W. (2013). *Role of the Waist / Height Ratio in the Cardiometabolic Risk Assessment of Children Classified by Body Mass Index*. 62(8). <https://doi.org/10.1016/j.jacc.2013.01.026>
- Kim, B., Choi, H. Y., Kim, W., Ahn, C., Lee, J., Kim, J. G., ... Moon, S. (2018). The cut-off values of surrogate measures for insulin resistance in the Korean population according to the Korean Genome and Epidemiology Study (KOGES). *PLoS ONE*, 13(11), 1–10. <https://doi.org/10.1371/journal.pone.0206994>
- Kim, Y., & Jee, H. (2017). Influences of body mass index and physical activity on hypertension and stroke in Korean adult males: 10-year longitudinal study. *Journal of exercise nutrition & biochemistry*, 21(2), 16–22. <https://doi.org/10.20463/jenb.2017.0003>
- Kothari, C. and Garg, G., 2019. *Research methodology Methods and Techniques*. 3rd ed. New Delhi: New Age International (P) Ltd., p.63.
- Kouli, G. M., Panagiotakos, D. B., Kyrou, I., Georgousopoulou, E. N., Chrysohoou, C., Tsigos, C., Tousoulis, D., & Pitsavos, C. (2017). Visceral adiposity index and 10-year cardiovascular disease incidence: The ATTICA study. *Nutrition, metabolism, and cardiovascular diseases : NMCD*, 27(10), 881–889. <https://doi.org/10.1016/j.numecd.2017.06.015>
- Kuk, J. L., Saunders, T. J., Davidson, L. E., & Ross, R. (2009). *Age-related changes in total and regional fat distribution*. 8, 339–348. <https://doi.org/10.1016/j.arr.2009.06.001>
- Kumpatla, S., Michael, C., & Viswanathan, V. (2011). Visceral Adiposity Index and pattern of dyslipidaemia at different stages of glucose intolerance – A study from India. *Diabetes and Metabolic Syndrome: Clinical Research and Reviews*, 5(4), 173–178. <https://doi.org/10.1016/j.dsx.2012.05.003>
- Lam, B. C. C., Koh, G. C. H., Chen, C., Wong, M. T. K., & Fallows, S. J. (2015). Comparison of Body Mass Index (BMI), Body Adiposity Index (BAI), Waist Circumference (WC), Waist-To-Hip Ratio (WHR) and Waist-To-Height Ratio (WHtR) as predictors of cardiovascular disease risk factors in an adult population in Singapore. *PLoS ONE*, 10(4), 1–15. <https://doi.org/10.1371/journal.pone.0122985>

- Lean, M. E. J., Han, T. S., & Morrison, C. E. (1995). Waist circumference as a measure for indicating need for weight management. *Bmj*, *311*(6998), 158. <https://doi.org/10.1136/bmj.311.6998.158>
- Lee, C. M. Y., Huxley, R. R., Wildman, R. P., & Woodward, M. (2008). Indices of abdominal obesity are better discriminators of cardiovascular risk factors than BMI: a meta-analysis. *Journal of Clinical Epidemiology*, *61*(7), 646–653. <https://doi.org/10.1016/j.jclinepi.2007.08.012>
- Lee, P. H., Macfarlane, D. J., Lam, T. H., & Stewart, S. M. (2011). Validity of the International Physical Activity Questionnaire Short Form (IPAQ-SF): a systematic review. *The international journal of behavioral nutrition and physical activity*, *8*, 115. <https://doi.org/10.1186/1479-5868-8-115>
- Leiter LA, Fitchett DH, Gupta REG, Mancini GBJ, McFarlane PA, Ross R, et al. Identification and management of cardiometabolic risk in Canada: A position paper by the cardiometabolic risk working group (Executive summary). *Can J Cardiol*. 2011;27(2):124–31.
- Liao, C., Gao, W., Cao, W., Lv, J., Yu, C., Wang, S., ... Li, L. (2021). Associations of metabolic/obesity phenotypes with insulin resistance and C-reactive protein: Results from the CNTR study. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, *14*, 1141–1151. <https://doi.org/10.2147/DMSO.S298499>
- Libman, I. M., Barinas-Mitchell, E., Bartucci, A., Chaves-Gnecco, D., Robertson, R., & Arslanian, S. (2010). Fasting and 2-hour plasma glucose and insulin: Relationship with risk factors for cardiovascular disease in overweight nondiabetic children. *Diabetes Care*, *33*(12), 2674–2676. <https://doi.org/10.2337/dc10-0085>
- Lietz Jabłonowska, B., Wrzosek, M., Włodarczyk, M., & Nowicka, G. (2017). New indexes of body fat distribution, visceral adiposity index, body adiposity index, waist-to-height ratio, and metabolic disturbances in the obese. *Kardiologia Polska*, *75*(11), 1185–1191. <https://doi.org/10.5603/KP.a2017.0149>
- Linder, S., Abu-Omar, K., Geidl, W., Messing, S., Sarshar, M., Reimers, A. K., & Ziemainz, H. (2021). Physical inactivity in healthy, obese, and diabetic adults in Germany: An analysis of related socio-demographic

- variables. *PLoS ONE*, 16(2 February), 1–14. <https://doi.org/10.1371/journal.pone.0246634>
- Ling, J. C. Y., Mohamed, M. N. A., Jalaludin, M. Y., Rampal, S., Zaharan, N. L., & Mohamed, Z. (2016). Determinants of High Fasting Insulin and Insulin Resistance among Overweight/Obese Adolescents. *Scientific Reports*, 6(October), 1–10. <https://doi.org/10.1038/srep36270>
- Liu, P. J., Ma, F., Lou, H. P., & Chen, Y. (2016). Visceral adiposity index is associated with pre-diabetes and type 2 diabetes mellitus in Chinese adults aged 20-50. *Annals of Nutrition and Metabolism*, 68(4), 235–243. <https://doi.org/10.1159/000446121>
- Lohmann, T. G., Roche, A. F., & Martorell, R. (1988). Anthropometric Standardization Reference Manual Human Kinetics Books. Champaign, Ill.
- Lomas, J., Anderson, G. M., Domnick-Pierre, K., Vayda, E., & Enkin, M. W. (1989). The New England Journal of Medicine Downloaded from nejm.org at PENN STATE UNIVERSITY on November 25, 2015. For personal use only. No other uses without permission. From the NEJM Archive. Copyright © 2010 Massachusetts Medical Society. All rights reserved. *N Engl J Med*, 321(19), 1306–1311.
- Longo, M., Zatterale, F., Naderi, J., Parrillo, L., Formisano, P., Raciti, G. A., Beguinot, F., & Miele, C. (2019). Adipose Tissue Dysfunction as Determinant of Obesity-Associated Metabolic Complications. *International journal of molecular sciences*, 20(9), 2358. <https://doi.org/10.3390/ijms20092358>
- Luhar, S., Alice, P., Mallinson, C., Clarke, L., & Kinra, S. (2018). *Trends in the socioeconomic patterning of overweight / obesity in India: a repeated cross-sectional study using nationally representative data*. 6–9. <https://doi.org/10.1136/bmjopen-2018-023935>
- Lukács, A., Horváth, E., Máté, Z., Szabó, A., Virág, K., Papp, M., ... Paulik, E. (2019). Abdominal obesity increases metabolic risk factors in non-obese adults: A Hungarian cross-sectional study. *BMC Public Health*, 19(1), 1–8. <https://doi.org/10.1186/s12889-019-7839-1>

- Luo, Y., Ma, X., Shen, Y., Hao, Y., Hu, Y., Xiao, Y., ... Jia, W. (2014). Positive relationship between serum low-density lipoprotein cholesterol levels and visceral fat in a Chinese nondiabetic population. *PLoS ONE*, 9(11), 1–7. <https://doi.org/10.1371/journal.pone.0112715>
- Lwow, F., Jedrzejuk, D., Milewicz, A., & Szmigiero, L. (2016). Lipid accumulation product (LAP) as a criterion for the identification of the healthy obesity phenotype in postmenopausal women. *Experimental Gerontology*, 82, 81–87. <https://doi.org/10.1016/j.exger.2016.06.007>
- Manini T. M. (2010). Energy expenditure and aging. *Ageing research reviews*, 9(1), 1–11. <https://doi.org/10.1016/j.arr.2009.08.002>
- Männistö, S., Harald, K., Kontto, J., Lahti-Koski, M., Kaartinen, N. E., Saarni, S. E., ... Jousilahti, P. (2014). Dietary and lifestyle characteristics associated with normal-weight obesity: The National FINRISK 2007 Study. *British Journal of Nutrition*, 111(5), 887–894. <https://doi.org/10.1017/S0007114513002742>
- Martinez, K. E., Tucker, L. A., Bailey, B. W., & LeCheminant, J. D. (2017). Expanded normal weight obesity and insulin resistance in US adults of the national health and nutrition examination survey. *Journal of Diabetes Research*, 2017. <https://doi.org/10.1155/2017/9502643>
- Matsuzawa, Y., Fujioka, S., Tokunaga, K., & Tarui, S. (1987). Classification of obesity with respect to morbidity. *Proceedings of the Society for Experimental Biology and Medicine*, 200(2), 197-201.
- Matthews, D. R., Hosker, J. P., Rudenski, A. S., Naylor, B. A., Treacher, D. F., & Turner, R. C. (1985). Homeostasis model assessment: insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*, 28(7), 412-419.
- Mazidi, M., Kengne, A. P., Katsiki, N., Mikhailidis, D. P., & Banach, M. (2018). Lipid accumulation product and triglycerides/glucose index are useful predictors of insulin resistance. *Journal of Diabetes and Its Complications*, 32(3), 266–270. <https://doi.org/10.1016/j.jdiacomp.2017.10.007>
- Merchant, A. T., Anand, S. S., Vuksan, V., Jacobs, R., Davis, B., Teo, K., & Yusuf, S.. (2005). Protein Intake Is Inversely Associated with Abdominal Obesity in

- a Multi-Ethnic Population. *The Journal of Nutrition*, 135(5), 1196–1201. <https://doi.org/10.1093/jn/135.5.1196>
- Miller, L. C., Joshi, N., Lohani, M., Rogers, B., Mahato, S., Ghosh, S., & Webb, P. (2017). Women's education level amplifies the effects of a livelihoods-based intervention on household wealth, child diet, and child growth in rural Nepal. *International Journal for Equity in Health*, 16(1), 1–17. <https://doi.org/10.1186/s12939-017-0681-0>
- Mirmiran, P., Bahadoran, Z., & Azizi, F. (2014). Lipid accumulation product is associated with insulin resistance, lipid peroxidation, and systemic inflammation in type 2 diabetic patients. *Endocrinology and Metabolism*, 29(4), 443–449. <https://doi.org/10.3803/EnM.2014.29.4.443>
- Misra, A. (2015). Ethnic-Specific Criteria for Classification of Body Mass Index: A Perspective for Asian Indians and American Diabetes Association Position Statement. *Diabetes Technology and Therapeutics*, 17(9), 667–671. <https://doi.org/10.1089/dia.2015.0007>
- Misra, A., & Khurana, L. (2011). Obesity-related non-communicable diseases: South Asians vs White Caucasians. *International Journal of Obesity*, 35(2), 167–187. <https://doi.org/10.1038/ijo.2010.135>
- Misra, A., & Shrivastava, U. (2013). Obesity and dyslipidemia in South Asians. *Nutrients*, 5(7), 2708–2733. <https://doi.org/10.3390/nu5072708>
- Mittal B. (2019). Subcutaneous adipose tissue & visceral adipose tissue. *The Indian journal of medical research*, 149(5), 571–573. https://doi.org/10.4103/ijmr.IJMR_1910_18
- Miyazaki, Y., & DeFronzo, R. A. (2009). Visceral fat dominant distribution in male type 2 diabetic patients is closely related to hepatic insulin resistance, irrespective of body type. *Cardiovascular diabetology*, 8, 44. <https://doi.org/10.1186/1475-2840-8-44>
- Mogre, V., Nyaba, R., & Aleyira, S. (2014). *Lifestyle Risk Factors of General and Abdominal Obesity in Students of the School of Medicine and Health Science of the University of Development Studies , Tamale , Ghana. 2014.*
- Mohan, V. R., & Muliylil, J. (2009). Mortality patterns and the effect of socioeconomic factors on mortality in rural Tamil Nadu, south India: a community-based cohort study. *Transactions of the Royal Society*

- of Tropical Medicine and Hygiene*, 103(8), 801–806.
<https://doi.org/10.1016/j.trstmh.2009.04.014>
- Montero Rodea, E. R., Evia-viscarra, M. L., & Apolinar-jiménez, E. (2014). *Waist-to-Height Ratio Is a Better Anthropometric Index than Waist Circumference and BMI in Predicting Metabolic Syndrome among Obese Mexican Adolescents*. 2014.
- Morshed Shahed, M., Haq, T., Fariduddin, M., Parveen, S., T.-E.-A., Mahbub, N., & Hasanat, M. A. (2021). The Association of Visceral Adiposity Index with Insulin Resistance in Adults with Prediabetes. *Open Journal of Endocrine and Metabolic Diseases*, 11(01), 12–23.
<https://doi.org/10.4236/ojemd.2021.111002>
- Moon, H. U., Ha, K. H., Han, S. J., Kim, H. J., & Kim, D. J. (2018). The association of adiponectin and visceral fat with insulin resistance and β -cell dysfunction. *Journal of Korean medical science*, 34(1).
- Mottillo, S., Filion, K. B., Genest, J., Joseph, L., Pilote, L., Poirier, P., Eisenberg, M. J. (2010). The metabolic syndrome and cardiovascular risk: A systematic review and meta-analysis. *Journal of the American College of Cardiology*, 56(14), 1113–1132. <https://doi.org/10.1016/j.jacc.2010.05.034>
- Mozaffarian D, Hao T, Rimm EB, Willett WC, Hu FB *N Engl J Med*. 2011 Jun 23; 364(25):2392-404.
- Munusamy, V., George, M., Jena, A., Sridhar, A., & Elumalai, D. V. (2015). Comparison of Visceral Adiposity Index with Other Indices of Adiposity in Patients with Acute Myocardial Infarction. *Pharmacology, Toxicology and Biomedical Reports*, 1(2), 55–61. <https://doi.org/10.5530/ptb.1.2.4>
- Murakami, K., & Livingstone, M. B. E. (2015). Eating frequency is positively associated with overweight and central obesity in US adults. *Journal of Nutrition*, 145(12), 2715–2724. <https://doi.org/10.3945/jn.115.219808>
- Mutsert, R., Gast, K., Widya, R., de Koning, E., Jazet, I., Lamb, H., ... & den Heijer, M. (2018). Associations of abdominal subcutaneous and visceral fat with insulin resistance and secretion differ between men and women: the Netherlands epidemiology of obesity study. *Metabolic syndrome and related disorders*, 16(1), 54-63.

- Nascimento, J. X. P. T., Chein, M. B. D. C., De Sousa, R. M. L., Ferreira, A. D. S., Navarro, P. A., & Brito, L. M. O. (2015). Importance of lipid accumulation product index as a marker of CVD risk in PCOS women. *Lipids in Health and Disease*, *14*(1), 1–8. <https://doi.org/10.1186/s12944-015-0061-y>
- Nauck, M., Warnick, G. R., & Rifai, N. (2002). Methods for measurement of LDL-cholesterol: A critical assessment of direct measurement by homogeneous assays versus calculation. *Clinical Chemistry*, *48*(2), 236–254. <https://doi.org/10.1093/clinchem/48.2.236>
- Nayak Ramdas, V. K., Nayak, K. R., Vidyasagar, S., & P, R. (2020). Predictive performance of traditional and novel lipid combined anthropometric indices to identify prediabetes. *Diabetes and Metabolic Syndrome: Clinical Research and Reviews*, *14*(5), 1265–1272. <https://doi.org/10.1016/j.dsx.2020.06.045>
- Neeland, I. J., Ayers, C. R., Rohatgi, A. K., Turer, A. T., Berry, J. D., Das, S. R., Vega, G. L., Khera, A., McGuire, D. K., Grundy, S. M., & de Lemos, J. A. (2013). Associations of visceral and abdominal subcutaneous adipose tissue with markers of cardiac and metabolic risk in obese adults. *Obesity (Silver Spring, Md.)*, *21*(9), E439–E447. <https://doi.org/10.1002/oby.20135>
- Neeland, I. J., Turer, A. T., Ayers, C. R., Powell-Wiley, T. M., Vega, G. L., Farzaneh-Far, R., ... & de Lemos, J. A. (2012). Dysfunctional adiposity and the risk of prediabetes and type 2 diabetes in obese adults. *Jama*, *308*(11), 1150-1159.
- Nho, J.-H. (2017). Lifestyle Intervention for Obese Women. *Journal of Lifestyle Medicine*, *7*(2), 51–54. <https://doi.org/10.15280/jlm.2017.7.2.51>
- Nishida, C., Barba, C., Cavalli-Sforza, T., Cutter, J., Deurenberg, P., Darnton-Hill, I., ... Zimmet, P. (2004). Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *The Lancet*, *363*(9403), 157–163. [https://doi.org/10.1016/S0140-6736\(03\)15268-3](https://doi.org/10.1016/S0140-6736(03)15268-3)
- Nusrianto, R., Ayundini, G., Kristanti, M., Astrella, C., Amalina, N., Riyadina, W., ... & Soewondo, P. (2019). Visceral adiposity index and lipid accumulation product as a predictor of type 2 diabetes mellitus: the Bogor cohort study of

- non-communicable diseases risk factors. *Diabetes research and clinical practice*, 155, 107798.
- Nusselder, W. J., Franco, O. H., Peeters, A., & Mackenbach, J. P. (2009). Living healthier for longer: comparative effects of three heart-healthy behaviors on life expectancy with and without cardiovascular disease. *BMC Public Health*, 9(1), 1-9.
- O'Doherty, M. G., Cairns, K., O'Neill, V., Lamrock, F., Jørgensen, T., Brenner, H., Schöttker, B., Wilsgaard, T., Siganos, G., Kuulasmaa, K., Boffetta, P., Trichopoulou, A., & Kee, F. (2016). Effect of major lifestyle risk factors, independent and jointly, on life expectancy with and without cardiovascular disease: results from the Consortium on Health and Ageing Network of Cohorts in Europe and the United States (CHANCES). *European journal of epidemiology*, 31(5), 455–468. <https://doi.org/10.1007/s10654-015-0112-8>.
- Oh, J. Y., Sung, Y. A., & Lee, H. J. (2013). The lipid accumulation product as a useful index for identifying abnormal glucose regulation in young Korean women. *Diabetic Medicine*, 30(4), 436-442.
- Oh, S. K., Cho, A. R., Kwon, Y. J., Lee, H. S., & Lee, J. W. (2018). Derivation and validation of a new visceral adiposity index for predicting visceral obesity and cardiometabolic risk in a Korean population. *PLoS One*, 13(9), e0203787.
- Okafor, C. I., Fasanmade, O., Ofoegbu, E., & Ohwovoriole, A. E. (2011). *Original Article Comparison of the performance of two measures of central adiposity among apparently healthy Nigerians using the receiver operating characteristic analysis*. 15(4). <https://doi.org/10.4103/2230-8210.85588>
- Oli, N., Vaidya, A., & Thapa, G.. (2013). Behavioural Risk Factors of Noncommunicable Diseases among Nepalese Urban Poor: A Descriptive Study from a Slum Area of Kathmandu. *Epidemiology Research International*, 2013, 1–13. <https://doi.org/10.1155/2013/329156>
- Olivera, S., Muralidhara, H. B., Venkatesh, K., Guna, V. K., Gopalakrishna, K., & Kumar, Y. (2016). Potential applications of cellulose and chitosan nanoparticles/composites in wastewater treatment: a review. *Carbohydrate polymers*, 153, 600-618.
- Parente., Duncan, B. B., ... & Velasquez-Melendez, G. (2020). The burden of non-communicable diseases attributable to high BMI in Brazil, 1990–2017:

- Findings from the Global Burden of Disease Study. *Population Health Metrics*, 18(1), 1-13.
- Pajunen, P., Kotronen, A., Korpi-Hyövälti, E., Keinänen-Kiukaanniemi, S., Oksa, H., Niskanen, L., ... Peltonen, M. (2011). Metabolically healthy and unhealthy obesity phenotypes in the general population: The FIN-D2D Survey. *BMC Public Health*, 11. <https://doi.org/10.1186/1471-2458-11-754>
- Pan, M., Tu, R., Gu, J., Li, R., Liu, X., Chen, R., ... Hou, J. (2021). Associations of Socioeconomic Status and Physical Activity With Obesity Measures in Rural Chinese Adults. 8(January), 1–10. <https://doi.org/10.3389/fpubh.2020.594874>
- Paneni, F. (2014). Role of oxidative stress in endothelial insulin resistance. *World Journal of Diabetes*, 6(2), 326. <https://doi.org/10.4239/wjd.v6.i2.326>
- Paniagua, J. A. (2016). Nutrition, insulin resistance and dysfunctional adipose tissue determine the different components of metabolic syndrome. *World Journal of Diabetes*, 7(19), 483. <https://doi.org/10.4239/wjd.v7.i19.483>
- Parente, E. B., Mutter, S., Harjutsalo, V., Ahola, A. J., Forsblom, C., & Groop, P. H. (2020). Waist-height ratio and waist are the best estimators of visceral fat in type 1 diabetes. *Scientific Reports*, 10(1), 1–10. <https://doi.org/10.1038/s41598-020-75667-5>
- Park, Y. W., Zhu, S., Palaniappan, L., Heshka, S., Carnethon, M. R., & Heymsfield, S. B. (2003). The metabolic syndrome: prevalence and associated risk factor findings in the US population from the Third National Health and Nutrition Examination Survey, 1988-1994. *Archives of internal medicine*, 163(4), 427–436. <https://doi.org/10.1001/archinte.163.4.427>
- Parveen, S., Ayub, T. E., Haq, T., Nahar, N., Manbub, N., Islam, F., ... & Aziz, M. (2020). Association of visceral adiposity index with insulin resistance in adults with diabetes mellitus. *IMC Journal of Medical Science*, 14(1), 5-12.
- Pasdar, Y., Moradi, S., Moludi, J., Saiedi, S., Moradinazar, M., Hamzeh, B., ... Najafi, F. (2020). Waist-to-height ratio is a better discriminator of cardiovascular disease than other anthropometric indicators in Kurdish adults. *Scientific Reports*, 10(1), 1–10. <https://doi.org/10.1038/s41598-020-73224-8>

- Patel, P., & Abate, N. (2013). Body fat distribution and insulin resistance. *Nutrients*, 5(6), 2019-2027.
- Patel, S. A., Shivashankar, R., Ali, M. K., Anjana, R. M., Deepa, M., Kapoor, D., Kondal, D., Rautela, G., Mohan, V., Narayan, K. M., Kadir, M. M., Fatmi, Z., Prabhakaran, D., Tandon, N., & CARRS Investigators (2016). Is the "South Asian Phenotype" Unique to South Asians?: Comparing Cardiometabolic Risk Factors in the CARRS and NHANES Studies. *Global heart*, 11(1), 89–96.e3. <https://doi.org/10.1016/j.gheart.2015.12.010s>
- Pathak, K. Y., Mohanan, A., Acharya, S., Mandavia, D., & Jadhav, H. R. (2018). Exploring visceral adiposity index as a predictor of visceral adiposity dysfunction and evaluating its performance in predicting hepatic insulin resistance in Indian type 2 diabetics. *International Journal of Pharmacy and Pharmaceutical Sciences*, 8(8), 297–301.
- Petersen, M. C., & Shulman, G. I. (2018). Mechanisms of insulin action and insulin resistance. *Physiological reviews*, 98(4), 2133-2223.
- Powell, K. E., Paluch, A. E., & Blair, S. N. (2011). Physical activity for health: What kind? how much? how intense? on top of what? *Annual Review of Public Health*, 32, 349–365. <https://doi.org/10.1146/annurev-publhealth-031210-101151>
- Pradeepa, R., Anjana, R. M., Joshi, S. R., Bhansali, A., Deepa, M., Joshi, P. P., ... Das, A. K. (2015). Prevalence of generalized & abdominal obesity in urban & rural India- the ICMR-INDIAB study (Phase-I) [ICMR-INDIAB-3]. *Indian Journal of Medical Research*, 142(AUGUST), 139–150. <https://doi.org/10.4103/0971-5916.164234>
- Prasad, D. S., Kabir, Z., Dash, A. L., & Das, B. C. (2012). Prevalence and risk factors for metabolic syndrome in Asian Indians: A community study from urban Eastern India. *Journal of Cardiovascular Disease Research*, 3(3), 204–211. <https://doi.org/10.4103/0975-3583.98895>
- Prochaska, J. J., & Prochaska, J. O. (2011). A Review of Multiple Health Behavior Change Interventions for Primary Prevention. *American journal of lifestyle medicine*, 5(3), 10.1177/1559827610391883. <https://doi.org/10.1177/1559827610391883>

- Pujilestari, C. U., Nyström, L., Norberg, M., Weinehall, L., & Hakimi, M. (2017). *Socioeconomic inequality in abdominal obesity among older people in Purworejo District , Central Java , Indonesia – a decomposition analysis approach*. 16. <https://doi.org/10.1186/s12939-017-0708-6>
- Ramya, M. R., Kamath, V., & S., H. (2017). Assessment of physical activity level among obese and non-obese individuals using global physical activity questionnaire. *International Journal Of Community Medicine And Public Health*, 4(10), 3786. <https://doi.org/10.18203/2394-6040.ijcmph20174251>
- Randrianarisoa, E., Lehn-Stefan, A., Hieronimus, A., Rietig, R., Fritsche, A., Machann, J., Balletshofer, B., Häring, H. U., Stefan, N., & Rittig, K. (2019). Visceral Adiposity Index as an Independent Marker of Subclinical Atherosclerosis in Individuals Prone to Diabetes Mellitus. *Journal of atherosclerosis and thrombosis*, 26(9), 821–834. <https://doi.org/10.5551/jat.47274>
- Range, D., Time, T., Life, S., Linked, E., Assay, I., & Complex, S. (2017). Insulin Human ELISA Kit. <https://www.thermofisher.com/elisa/product/insulin-human-elisa-kit/KAQ1251>, 0–12.
- Ravussin, E., & Smith, S. R. (2002). Increased fat intake, impaired fat oxidation, and failure of fat cell proliferation result in ectopic fat storage, insulin resistance, and type 2 diabetes mellitus. *Annals of the New York Academy of Sciences*, 967(1), 363-378.
- Ray, L., Ravichandran, K., & Nanda, S. K. (2018). Comparison of lipi accumulation product index with body mass index and waist circumference as a predictor of metabolic syndrome in Indian population. *Metabolic syndrome and related disorders*, 16(5), 240-245.
- Reaven, G. (2004). The metabolic syndrome or the insulin resistance syndrome? Different names, different concepts, and different goals. *Endocrinology and Metabolism Clinics*, 33(2), 283-303.
- Rifai N, Bachorik PS, Albers JJ. Lipids, lipoproteins and apolipoproteins. In: Burtis CA, Ashwood ER, editors. *Tietz Textbook of Clinical Chemistry*. 3rd ed. Philadelphia: W.B Saunders Company; 1999. p. 809-61.

- Rivero Bacelis, A. P., Vázquez-Rodríguez, A., Carrasco-Legleu, C. E., De León, L. G., & Reza-López, S. A. (2020). Assessment of Physical Activity in Adults: A Review of Validated Questionnaires From a Nutritionist's Point of View. *Evaluation & the Health Professions*, 43(4), 235–254. <https://doi.org/10.1177/0163278720928500>
- Rodríguez Velásquez, C. M., Velásquez-Villa, M., Gómez-Ocampo, L., & Bermúdez-Cardona, J. (2014). Abdominal obesity and low physical activity are associated with insulin resistance in overweight adolescents: a cross-sectional study. *BMC pediatrics*, 14, 258. <https://doi.org/10.1186/1471-2431-14-258>
- Roger Shrimpton; Claudia Rokx,. (2012). *The Double Burden of Malnutrition : A Review of Global Evidence. Health, Nutrition and Population Discussion Paper; World Bank, Washington, DC. © World Bank. <https://openknowledge.worldbank.org/handle/10986/27417> License: CC BY 3.0 IGO.*” a useful index for identifying abnormal glucose regulation in young Korean women. *Diabetic Medicine*, 30(4), 436–442. <https://doi.org/10.1111/dme.12052>
- Ross, R. (2020). Statement Waist circumference as a vital sign in clinical practice : a Consensus Statement from the IAS and ICCR Working Group on Visceral Obesity. *Nature Reviews Endocrinology*, 16(March), 177–189. <https://doi.org/10.1038/s41574-019-0310-7>
- Rugge, B., Balslem, H., & Sehgal, R. (2011). Screening and Subclinical Hypothyroidism or Hyperthyroidism. *Agency for Healthcare Research and Quality U.S. Department of Health and Human Services*, (24).
- Ryo, M., Kishida, K., Nakamura, T., Yoshizumi, T., Funahashi, T., & Shimomura, I. (2014). Clinical significance of visceral adiposity assessed by computed tomography: a Japanese perspective. *World journal of radiology*, 6(7), 409.
- Sahakyan, K. R., Somers, V. K., Rodriguez-Escudero, J. P., Hodge, D. O., Carter, R. E., Sochor, O., Coutinho, T., Jensen, M. D., Roger, V. L., Singh, P., & Lopez-Jimenez, F. (2015). Normal-Weight Central Obesity: Implications for Total and Cardiovascular Mortality. *Annals of internal medicine*, 163(11), 827–835. <https://doi.org/10.7326/M14-2525>

- Salazar, J., Bermúdez, V., Fuenmayor, J., Nava, M., Ortega, Á., Duran, P., ... Rojas, J. (2021). Lipid Accumulation Product Is More Related to Insulin Resistance than the Visceral Adiposity Index in the Maracaibo City Population, Venezuela. *Journal of Obesity*, 2021. <https://doi.org/10.1155/2021/5514901>
- Samuel, P., Antonisamy, B., Raghupathy, P., Richard, J., & Fall, C. H. (2012). Socio-economic status and cardiovascular risk factors in rural and urban areas of Vellore, Tamilnadu, South India. *International journal of epidemiology*, 41(5), 1315–1327. <https://doi.org/10.1093/ije/dys001>
- Sarvottam, K., Ranjan, P., & Yadav, U. (2020). Age group and gender-wise comparison of obesity indices in subjects of Varanasi. 64(2), 109–117. <https://doi.org/10.25259/IJPP>
- Sandeep S, Gokulakrishnan K, Velmurugan K, Deepa M, Mohan V. Visceral & subcutaneous abdominal fat in relation to insulin resistance & metabolic syndrome in non-diabetic south Indians. *Indian J Med Res*. 2010;131(5):629–35
- Sawant, I. S., Wadkar, P. N., Ghule, S. B., Rajguru, Y. R., Salunkhe, V. P., & Sawant, S. D. (2017). Enhanced biological control of powdery mildew in vineyards by integrating a strain of *Trichoderma afroharzianum* with sulphur. *Biological Control*, 114, 133–143. <https://doi.org/10.1016/j.biocontrol.2017.08.011>
- Schwingshackl, L., Missbach, B., & Hoffmann, G. (2016). An umbrella review of garlic intake and risk of cardiovascular disease. *Phytomedicine : international journal of phytotherapy and phytopharmacology*, 23(11), 1127–1133. <https://doi.org/10.1016/j.phymed.2015.10.015>
- Seidell, J. C., Pérusse, L., Després, J. P., & Bouchard, C. (2001). Waist and hip circumferences have independent and opposite effects on cardiovascular disease risk factors: The Quebec Family Study. *American Journal of Clinical Nutrition*, 74(3), 315–321. <https://doi.org/10.1093/ajcn/74.3.315>
- Shahavandi, M., Djafari, F., Shahinfar, H., Davarzani, S., Babaei, N., Ebaditabar, M., ... Shab-Bidar, S. (2020). The association of plant-based dietary patterns with visceral adiposity, lipid accumulation product, and triglyceride-

- glucose index in Iranian adults. *Complementary Therapies in Medicine*, 53(2), 145–156. <https://doi.org/10.1016/j.ctim.2020.102531>
- Shahraki, M., Shahraki, T., & Ansari, H. (2008). The effects of socio-economic status on BMI, waist:hip ratio and waist circumference in a group of Iranian women. *Public Health Nutrition*, 11(7), 757–761. <https://doi.org/10.1017/S1368980007001577>
- Sharaye, K. O., & Eboh, E. I. (2020). The Locus of Fat Over-Accumulation as Predictor of Cardiometabolic Risks among Non Obese Normotensive Adults. *Journal of Advances in Medicine and Medical Research*, 32(19), 27–35. <https://doi.org/10.9734/jammr/2020/v32i1930665>
- Shields, M., Tremblay, M. S., Connor Gorber, S., & Janssen, I. (2012). Abdominal obesity and cardiovascular disease risk factors within body mass index categories. *Health reports*, 23(2), 7–15.
- Shuster, A., Patlas, M., Pinthus, J. H., & Mourtzakis, M. (2012). The clinical importance of visceral adiposity: A critical review of methods for visceral adipose tissue analysis. *British Journal of Radiology*, 85(1009), 1–10. <https://doi.org/10.1259/bjr/38447238>
- Singh, P. N., Arthur, K. N., Orlich, M. J., James, W., Purty, A., Job, J. S., ... Sabaté, J. (2014). Global epidemiology of obesity, vegetarian dietary patterns, and noncommunicable disease in Asian Indians. *American Journal of Clinical Nutrition*, 100(SUPPL. 1), 359–364. <https://doi.org/10.3945/ajcn.113.071571>
- Song, J., Zhao, Y., Nie, S., Chen, X., Wu, X., & Mi, J. (2018). The effect of lipid accumulation product and its interaction with other factors on hypertension risk in Chinese Han population: a cross-sectional study. *PLoS One*, 13(6), e0198105.
- Sorkin, J. D., Muller, D. C., & Andres, R. (1999). Longitudinal change in height of men and women: implications for interpretation of the body mass index: the Baltimore Longitudinal Study of Aging. *American journal of epidemiology*, 150(9), 969-977.
- Stenholm, S., Head, J., Aalto, V., Kivimäki, M., Kawachi, I., Zins, M., & Vahtera, J. (2017). Body mass index as a predictor of healthy and disease-free life

- expectancy between ages 50 and 75: a multicohort study. *International journal of obesity*, 41(5), 769-775.
- Štěpánek, L., Horáková, D., Štěpánek, L., Cibičková, L., Karásek, D., Vaverková, H., Nakládalová, M., Juríčková, L., & Kollárová, H. (2019). Associations Between Homeostasis Model Assessment (HOMA) and Routinely Examined Parameters in Individuals With Metabolic Syndrome. *Physiological research*, 68(6), 921–930. <https://doi.org/10.33549/physiolres.934141>
- Sun, K., Lin, D., Feng, Q., Li, F., Qi, Y., Feng, W., ... & Liu, D. (2019). Assessment of adiposity distribution and its association with diabetes and insulin resistance: a population-based study. *Diabetology & metabolic syndrome*, 11(1), 1-10.
- Sun, Y., Liu, B., Snetselaar, L. G., Wallace, R. B., Caan, B. J., Rohan, T. E., ... Bao, W. (2019). Association of Normal-Weight Central Obesity with All-Cause and Cause-Specific Mortality among Postmenopausal Women. *JAMA Network Open*, 2(7), 1–13. <https://doi.org/10.1001/jamanetworkopen.2019.7337>
- Suryadinata, R. V., Wirjatmadi, B., Adriani, M., & Lorensia, A. (2020). Effect of age and weight on physical activity. *Journal of Public Health Research*, 9(2), 187–190. <https://doi.org/10.4081/jphr.2020.1840>
- Tehard, B., Saris, W. H. M., Astrup, A., Martinez, J. A., Taylor, M. A., Barbe, P., ... Oppert, J. M. (2005). Comparison of two physical activity questionnaires in obese subjects: The NUGENOB study. *Medicine and Science in Sports and Exercise*, 37(9), 1535–1541. <https://doi.org/10.1249/01.mss.0000177464.68521.3b>
- Tran, N. T., Blizzard, C. L., Luong, K. N., Van Truong, N. Le, Tran, B. Q., Otahal, P., ... Callisaya, M. (2018). The importance of waist circumference and body mass index in cross-sectional relationships with risk of cardiovascular disease in Vietnam. *PLoS ONE*, 13(5), 1–13. <https://doi.org/10.1371/journal.pone.0198202>

- Trinder, P. (1969). A Simple Turbidimetric Method for the Determination of Serum Cholesterol. *Annals of Clinical Biochemistry*, 6(5), 165–166. <https://doi.org/10.1177/000456326900600505>
- Trinder, P. (1969): Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor. *Annals of Clinical Biochemistry*, 6, 24-27
- Tutunchi, H., Ebrahimi-Mameghani, M., Ostadrahimi, A., & Asghari-Jafarabadi, M. (2020). What are the optimal cut-off points of anthropometric indices for prediction of overweight and obesity? Predictive validity of waist circumference, waist-to-hip and waist-to-height ratios. *Health promotion perspectives*, 10(2), 142–147. <https://doi.org/10.34172/hpp.2020.23>
- Unger, R. H. (2003). Lipid overload and overflow: Metabolic trauma and the metabolic syndrome. *Trends in Endocrinology and Metabolism*, 14(9), 398–403. <https://doi.org/10.1016/j.tem.2003.09.008>
- Vassilatou, E., Lafoyianni, S., Vassiliadi, D. A., Ioannidis, D., Paschou, S. A., Mizamtsidi, M., ... Vryonidou, A. (2018). Visceral adiposity index for the diagnosis of nonalcoholic fatty liver disease in premenopausal women with and without polycystic ovary syndrome. *Maturitas*, 116(June), 1–7. <https://doi.org/10.1016/j.maturitas.2018.06.013>
- Verboven, K., Wouters, K., Gaens, K., Hansen, D., Bijnen, M., Wetzels, S., Jocken, J. W. (2018). Abdominal subcutaneous and visceral adipocyte size, lipolysis and inflammation relate to insulin resistance in male obese humans. *Scientific Reports*, 8(1), 1–8. <https://doi.org/10.1038/s41598-018-22962-x>
- Veum, V. L., Laupsa-Borge, J., Eng, Ø., Rostrup, E., Larsen, T. H., Nordrehaug, J. E., ... Mellgren, G. (2017). Visceral adiposity and metabolic syndrome after very high-fat and low-fat isocaloric diets: A randomized controlled trial. *American Journal of Clinical Nutrition*, 105(1), 85–99. <https://doi.org/10.3945/ajcn.115.123463>
- Vizzuso, S., Del Torto, A., Dilillo, D., Calcaterra, V., Di Profio, E., Leone, A., & Verduci, E. (2021). Visceral adiposity index (VAI) in children and adolescents with obesity: No association with daily energy intake but promising tool to identify metabolic syndrome (MetS). *Nutrients*, 13(2), 413.

- Vlassoff C. (2007). Gender differences in determinants and consequences of health and illness. *Journal of health, population, and nutrition*, 25(1), 47–61.
- Vogel P, Stein A, Marcadenti A. (2016) Índice De Adiposidade Visceral E Prognóstico Em Pacientes Com Insuficiência Cardíaca Isquêmica. *Sao Paulo Med J.* ;134(3):211–8.
- Wakabayashi, I., & Daimon, T. (2014). A strong association between lipid accumulation product and diabetes mellitus in Japanese women and men. *Journal of Atherosclerosis and Thrombosis*, 21(3), 282–288. <https://doi.org/10.5551/jat.20628>
- Wallace, T. M., & Matthews, D. R. (2002). The assessment of insulin resistance in man. *Diabetic medicine*, 19(7), 527-534.
- Wallace, T. M., Levy, J. C., & Matthews, D. R. (2004). *Use and Abuse of HOMA Modeling. Diabetes Care*, 27(6), 1487– 1495. doi:10.2337/diacare.27.6.1487
- Warner, E. T., Wolin, K. Y., Duncan, D. T., Heil, D. P., Askew, S., & Bennett, G. G. (2012). Differential accuracy of physical activity self-report by body mass index. *American Journal of Health Behavior*, 36(2), 168–178. <https://doi.org/10.5993/AJHB.36.2.3>
- WebMD, (2021),; <https://www.webmd.com/diet/what-is-visceral-fat#1>
- Weiss, E. P., Jordan, R. C., Frese, E. M., Albert, S. G., & Villareal, D. T. (2017). Effects of Weight Loss on Lean Mass, Strength, Bone, and Aerobic Capacity. *Medicine and science in sports and exercise*, 49(1), 206–217. <https://doi.org/10.1249/MSS.0000000000001074>
- White, Z., Hakim, C. H., Theret, M., Yang, N. N., Rossi, F., Cox, D., ... Bernatchez, P. (2020). High prevalence of plasma lipid abnormalities in human and canine Duchenne and Becker muscular dystrophies depicts a new type of primary genetic dyslipidemia. *Journal of Clinical Lipidology*, 14(4), 459-469.e0. <https://doi.org/10.1016/j.jacl.2020.05.098>
- World Health Organization, 2013
- World Health Organization, 2008
- World Health Organization, 2003

- Wilcox, G. (2005). Insulin and insulin resistance. *Clinical biochemist reviews*, 26(2), 19.
- Winter, Y., Sankowski, R., & Back, T. (2013). Genetic determinants of obesity and related vascular diseases. *Vitamins & Hormones*, 91, 29-48.
- Wong, M. C. S., Huang, J., Wang, J., Chan, P. S. F., Lok, V., Chen, X., Zheng, Z. J. (2020). Global, regional and time-trend prevalence of central obesity: a systematic review and meta-analysis of 13.2 million subjects. *European Journal of Epidemiology*, 35(7), 673–683. <https://doi.org/10.1007/s10654-020-00650-3>
- Wu, Y., Li, H., Tao, X., Fan, Y., Gao, Q., & Yang, J. (2021). Optimised anthropometric indices as predictive screening tools for metabolic syndrome in adults: A cross-sectional study. *BMJ Open*, 11(1), 7–10. <https://doi.org/10.1136/bmjopen-2020-043952>
- Xia, C., Li, R., Zhang, S., Gong, L., Ren, W., Wang, Z., & Li, Q. (2012). Lipid accumulation product is a powerful index for recognizing insulin resistance in non-diabetic individuals. *European Journal of Clinical Nutrition*, 66(9), 1035–1038. <https://doi.org/10.1038/ejcn.2012.83>
- Xu, H., Li, X., Adams, H., Kubena, K., & Guo, S. (2018). Etiology of metabolic syndrome and dietary intervention. *International Journal of Molecular Sciences*, 20(1), 1–19. <https://doi.org/10.3390/ijms20010128>
- Xu, X., Zhao, Y., Zhao, Z., Zhu, S., Liu, X., Zhou, C., & Zou, H. (2016). Correlation of visceral adiposity index with chronic kidney disease in the People's Republic of China: to rediscover the new clinical potential of an old indicator for visceral obesity. *Therapeutics and clinical risk management*, 12, 489.
- Yang, Y. J., Kim, M. K., Hwang, S. H., Ahn, Y., Shim, J. E., & Kim, D. H. (2010). Relative validities of 3-day food records and the food frequency questionnaire. *Nutrition Research and Practice*, 4(2), 142. <https://doi.org/10.4162/nrp.2010.4.2.142>
- Zhang, X., Shu, X. O., Li, H., Yang, G., Xiang, Y. B., Cai, Q., Ji, B. T., Gao, Y. T., & Zheng, W. (2013). Visceral adiposity and risk of coronary heart disease

in relatively lean Chinese adults. *International journal of cardiology*, 168(3), 2141–2145. <https://doi.org/10.1016/j.ijcard.2013.01.275>

Zeng, Q., He, Y., Dong, S., Zhao, X., Chen, Z., Song, Z., ... Wang, Y. (2014). Optimal cut-off values of BMI, waist circumference and waist:height ratio for defining obesity in Chinese adults. *British Journal of Nutrition*, 112(10), 1735–1744. <https://doi.org/10.1017/S0007114514002657>