



BIBLIOGRAPHY

BIBLIOGRAPHY

Abell, L.L., Levy, B.B., Brodie, B.B. and Kendall, F.E. (1952), A simplified method for the estimation of total cholesterol in serum and demonstration of its specificity, *J. Biol. Chem.*, 195:357.

Adewole, S.O. and Martins, E.A.C. (2006), Morphological changes and hypoglycemic effects of *Annona muricata* Linn. (*Annonaceae*) leaf aqueous extract on pancreatic β -cells of streptozotocin treated diabetic rats, *Afr. J. Biomed. Res.*, 9:173-187.

Agarwal, S. (2007), Revolution in antioxidant and antioxidant therapy – Glutathione, *The Antiseptic*, 3:45-47.

Ahmad, I., Adeghate, E., Cummings, E., Sharma, A.K. and Singh, J. (2004), Beneficial effects and mechanism of action of *Momordica* fruit juice in the treatment of streptozotocin-induced diabetes mellitus in rats, *Mol. Cell. Biochem.*, 261:63-70.

Ahmed, R.S., Suke, S.G., Seth, V., Jain, A. and Bhattacharya, S.N. (2006), Impact of oral vitamin E supplementation on oxidative stress and lipid peroxidation in patients with polymorphous light eruption, *Indian J. Med. Res.*, 123:781-787.

Akilandeswari, S., Manimaran, S., Valarmathi, R., Sundari, S.K.K. and Loganathan, V. (2001), Phytochemical observation of leaf *Justicia tranquebariensis* L.F., *Ancient Science of Life*, 20:60-61.

Akinmoladun, C.A., Ibukun, E.O., Afor, E., Obuotor, A. and Farombi, E.O. (2007), Phytochemical constituent and antioxidant activity of extract from the leaves of *Ocimum gratissimum*, Sci. Res. and Essays, 2:163-166.

American Diabetes Association, (2002), Screening for diabetic retinopathy, Diabetes Care, 25:S90-S93.

American Diabetes Association, (2008a), Diagnosis and classification of diabetes mellitus, Diabetes Care, 31:S55-S60.

American Diabetes Association, (2008b), Diabetes-related complications, glycemic control and falls in older adults, Diabetes Care, 31:S61-S65.

Ananthi, J., Prakasam, A. and Pugalendi, K.V. (2003), Antihyperglycemic activity of *Eclipta alba* leaf on alloxan-induced diabetic rats, Yale J. Biol. Med., 76:97-102.

Anwar, M.M. and Meki, A.R.M.A. (2003), Oxidative stress in streptozotocin-induced diabetic rats: effects of garlic oil and melatonin, Comparative Biochemistry and Physiology, Molecular and Integrative Physiology, 135:539-547.

Anwer, T., Sharma, M., Pillai, K.K. and Igbal, M. (2008), Effect of *Withania somnifera* on insulin sensitivity in non-insulin dependent diabetes mellitus rats, BCPT., 102:498-503.

Armbruster, D.A. (1987), Fructosamine: structure, analysis and clinical usefulness, Clin. Chem., 33:2153-2163.

- Aslan, M., Orhan, D.D., Orhan, N., Sezik, E. and Yesilada, E. (2007), A study of antidiabetic and antioxidant effects of *Helichrysum graveolens* Capitulum in streptozotocin-induced diabetic rats, J. Medi. Food., 10:396-400.
- Babu, V., Gangadevi, T. and Subramonium, A. (2003), Antidiabetic activity of ethanol extract of *Cassia kleinii* leaf in streptozotocin-induced diabetic rats and isolation of an active fraction and toxicity evaluation of the extract, Indian J. Pharmacol., 35:290-296.
- Babu, P.S., Prabuseenivasan, S. and Ignacimuthu, S. (2007), Cinnamaldehyde- A potential antidiabetic agent, Phytomed., 14:15-22.
- Baker, H., Frank, O., Angelis, B. and Feingold, S. (1951), Plasma tocopherol in man at various times after ingesting free or acetylated tocopherol, Nutr. Rep. Inv., 21:531-536.
- Balogopal, N.P., Kamalamma, N., Patel, G.T. and Misra, R. (2008), A community-based diabetes prevention and management education program in a rural village in India, Diabetes Care, 31:1097-1104.
- Barik, R., Jain, S., Qwatra, D., Joshi, A., Tripathi, G.S. and Goyal, R. (2008), Antidiabetic activity of aqueous root extract of *Ichnocarpus frutescens* in streptozotocin-nicotinamide induced type-II diabetes in rats, Indian J. Pharmacol., 40:19-22.
- Basch, E., Ulbricht, C., Kuo, G., Szapary, P. and Smith, M. (2003), Therapeutic applications of fenugreek, BNET, 8:1-10.
- Baskar, R., Rajeswari, V. and Kumar, T.S. (2007), *in vitro* antioxidant studies in leaves of *Annona* species, Indian J. Exp. Biol., 45:480-485.

Bayfield, R.F. and Cole, E.R. (1980), Colorimetric estimation of vitamin A with trichloroacetic acid, *Meth. Enzymol.*, 67:189-203.

Bean, M.F., Antoun, M., Abramson, D., Chang, C.J., Mc Laughlin, J.L. and Cassady, J.M. (1985), Cucurbitacin B and isocucurbitacin B cytotoxic components of *Helicteres isora*, *J. Nat. Prod.*, 48:500-503.

Begum, S., Hassan, S.I., Siddiqui, B.S., Shaheen, F., Ghayur, M.N. and Gilani, A.H. (2002), Triterpenoids from the leaves of *Psidium guajava*, *Phytochem.*, 61:399-403.

Benze, W. and Schmid, H. (1954), *Experimentia*, 10:12.

Beutler, E. (1984), Red cell metabolism - A manual of biochemical methods, 3rd edition, Grune and Stratton Oriando, pp.11-20.

Bhatia, A., Arora, S., Nagpal, A., Singh, B. and Ahuja, P.S. (2007), Evaluation of *in vitro* antimutagenic activity of “seabuckthorn” (*Hippophae rhamnoides* Linn.) in *Ames* assay, *JCCM*, 2:428-434.

Bhavapriya, V., Kalpana, S., Govindasamy, S. and Apparanamtham, T. (2001), Biochemical studies on hypoglycemic effect of Aavirai kudineer: A herbal formulation in alloxan-diabetic rats, *Indian J. Exp. Biol.*, 39:925-928.

Bnouham, M., Ziyyat, A., Mekhfi, H., Tahri, A. and Legssyer, A. (2006), Medicinal plants with potential antidiabetic activity - A review of ten years of herbal medicine research (1990-2000), *Intl. J. Diabetes and Metabolism*, 14:1-25.

Bolkent, S., Sacan, O., Karatug, A. and Yanardag, R. (2008), The effects of vitamin B6 on the liver of diabetic rats: A morphological and biochemical study, *IUFS J. Biol.*, 67:1-7.

Bonnycastle, L.L., Willer, C.J., Conneely, K.N., Jackson, A.U., Burrill, C.P., Watanabe, R.M., Chines, P.S., Narisu, N., Scott, L.J., Enloe, S.T., Swift, A.J., Duren, W.L., Stringham, H.M., Erdos, M.R., Riebow, N.L., Buchanan, T.A., Valle, T.T., Tuomilehto, J., Bergman, R.N., Mohlke, K.L., Boehnke, M. and Collins, F.S. (2006), Common variants in maturity-onset diabetes of the young genes contribute to risk of type 2 diabetes in Finns, *Diabetes*, 55:534-540.

Bouhdid, S., Skali, S.N., Idaomar, M., Zhiri, A., Baudoux, D., Amensour, M. and Abrini, J. (2008), Antibacterial and antioxidant activities of *Origanum compactum* essential oil, *Afr. J. Biotechnol.*, 7:1563-1570.

Brandstrup, N., Kirk, J.E. and Bruni, C. (1957), Determination of hexokinase in tissues, *J. Gerontol.*, 12:166-171.

Brownlee, M. (2001), Biochemistry and molecular cell biology of diabetic complications, *Nature*, 414:813-820.

Buchanan, A.T. and Xiang, A.H. (2005), Gestational diabetes mellitus, *J. Clin. Invest.*, 115:485-491.

Bukan, N., Sancak, B., Yavuz, O., Koca, C., Tutkan, F., Ozcelikay, T.A. and Altan, N. (2003), Lipid peroxidation and scavenging enzyme levels in the liver of streptozotocin-induced diabetic rats, *ITBB* 40:447-450.

Buricova, L. and Reblova, Z. (2008), Czech medicinal plants as possible sources of antioxidants, J. Food Sci., 26:132-138.

Burke, J.P., Williams, K., Narayan, K.M.V., Leibson, C., Haffner, S.M. and Stern, M.P. (2003), A population perspective on diabetes prevention: whom should we target for preventing weight gain? Diabetes care, 26:1999-2004.

Cameron, G.R., Milton, R.F. and Allen, J.W. (1943), Estimation of flavonoids, Lancet, 179.

Cardium Study, (2007), Dyslipidemia, A pharmacor service, A decision resources publication, 4.

Catala, A. (2008), Lipid peroxidation of membrane phospholipids generates hydroxy-alkenals and oxidized phospholipids active in physiological and/or pathological conditions, Chemistry and Physics of Lipids, 157:1-11.

Catchpole, B., Kennedy, L.J., Davison, L.J. and Ollier, W.E.R. (2008), Canine diabetes mellitus: from phenotype to genotype, J. Small Anim. Pract., 49:4-10.

CDC Division of Media Relations, (2008), New resource for preventing diabetes in African – Americans, 404: 639-3286.

Chakrabarti, R.R.K., Vikramadithyan, R., Mullangi, V.M., Sharma, H., Jagadheshan, Y., Rao, N., Sairam, P. and Rajagopalan, R. (2002), Antidiabetic and hypolipidemic activity of *Helicteres isora* in animal models, J. Ethnopharmacol., 81:343-349.

Chakraborty, D., Mandal, S.M., Chakraborty, J., Bhattacharya, P.K., Bandyopadhyay, A., Mitra, A. and Gupta, K. (2007), Antimicrobial activity of leaf extract of *Basilicum polystachyon* (L) Moench, Indian J. Exp. Biol., 45:744-748.

Chen, X., Cui, L., Duan, X., Ma, B. and Zhong, D. (2006), Pharmacokinetics and metabolism of flavonoid scutellarein in humans after a single oral administration, Drug Metab. Dispos., 34:1345-1352.

Chen-Scarabelli, C. and Scarabelli, M.T. (2006), Suboptimal glycemic control, independently of QT interval duration, is associated with increased risk of ventricular arrhythmias in a high-risk population, Pacing Clin. Electrophysiol., 29:9-14.

Cho, E.J., Yokeozawa, T., Rhyu, T.D. and Park, C.J. (2003), The inhibitory effects of 12 medicinal plants and their component compounds on lipid peroxidation, The Am. J. Chinese Med., 31:907-917.

Chopra, R.N., Nayar, S.L. and Chopra, I.C. (1956), Glossary of Indian medicinal plants, 1st ed., NISCAIR, CSIR, New Delhi, India, pp. 131.

Clark, E.P. and Collip, J.B. (1925), A study of the Tisdall method for the determination of blood serum calcium with a suggested modification, Biochem., 63:461-463.

Coghlan, M. and Leighton, B. (2008), Glucokinase activators in diabetes management, Expert Opin. Investig. Drugs, 17:145-67.

Costa, J., Borges, M., David, C. and Carneiro. A.V. (2006), Efficacy of lipid lowering drug treatment for diabetic and non-diabetic patients: meta-analysis of randomised controlled trials, *BMJ*, 332:1115-1124.

Culling, C.F.A. (1979), *Handbook of histopathological and histochemical techniques*, 3rd edition, New York, Butterworth and Co. Ltd., p. 7.

Czupryniak, L., Wolska, S.M., Pawlowski, M. and Loba, J. (2006), Elevated systolic blood pressure is present in almost all individuals with newly diagnosed diabetes, *J. Hum. Hypertens.*, 20:231-233.

Dhevi, R., Gayathri, K., Shabil, M.M., Subashini, U., Dubey, P.G.V., Rajamanickam, G.V. and Chitra, M. (2008), A preliminary biochemical screening of *Helicteres isora* stem bark in carbon tetrachloride induced toxicity in rats, *Bulg. J. Vet. Med.*, 11:235-242.

Dieye, A.M., Sarr, A., Diop, S.N., Ndiaye, M., Sy, G.Y., Diarra, M., Gaffary, R.I., Sy, A.N. and Faye, B. (2008), Medicinal plants and the treatment of diabetes in Senegal: survey with patients, *Fundam. Clin. Pharmacol.*, 22:211-216.

Diwan, A.G., Pradhan, A.B., Lingojar, D., Krishna, K.K., Singh, P. and Almelkar, S.I. (2006), Serum zinc, chromium and magnesium levels in type 2 diabetes, *Intl. J. Diab.*, 26:122-123.

Djomeni, P.D.D., Tedong, L., Asongalem, E.A., Dimo, T., Sokeng, S.D. and Kamtchouing, P. (2006), Hypoglycaemic and antidiabetic effect of root extracts of *Ceiba pentandra* in normal and diabetic rats, *Afr. J. Trad. Cam.*, 3:129-136.

Drabkin, D.L. and Austin, J.M. (1932), Spectrophotometric constants for common haemoglobin derivatives in human, dog and rabbit blood, J. Biol. Chem., 98:719-733.

Eddouks, M. and Maghrani, M. (2004), Phlorizin-like effect of *Fraxinus excelsior* in normal and diabetic rats, J. Ethnopharmacol., 94:149-154.

Eddouks, M., Lemhadri, A. and Michel, J.B. (2004), Caraway and Caper: Potential antihyperglycemic plants in diabetic rats, J. Ethnopharmacol., 94 : 143 - 148

Edem, M.S., Emeribe, A.O. and Akpotuzor, J.O. (2008), Haemorrheologic and fibrinolytic activities in diabetics resident in Calabar, Cross-River State, Nigeria, Afr. J. Biotechnol., 7:1400-1403.

Edeoga, H.O., Okwu, D.E. and Mbaebie, B.O. (2003), Phytochemical constituents of some Nigerian medicinal plants, Afr. J. Biotechnol., 4:685-688.

Elizabeth, K. and Rao, M.N.A. (1990), Oxygen radical scavenging activity of curcumin, Intl. J. Pharm., 58:237-240.

Evans, P. and Halliwell, B. (2001), Micronutrients: oxidant/antioxidant status, Brit. J. Nutr., 85:S67-S74.

Fabricant, D.S. and Farnsworth, N.R. (2001), The value of plants used in traditional medicine for drug discovery, EHP, 109:69-75.

Fadzelly, A.B.M., Asmah, R. and Fauziah, O. (2006), Effect of *Strobilanthes crispus* tea aqueous extracts on glucose and lipid profile in normal and streptozotocin-induced hyperglycemic rats, *Plant Food Hum. Nutr.*, 61:6-11.

Fale, L.P., Borges, C., Madeira, A.J.P., Ascensao, L., Araujo, M.E.M., Florencio, H.M. and Serralheiro, M.L.M. (2008), Rosmarinic acid, scutellarein 4'-methyl ether 7-*O*-glucuronide and (16S)-coleon E are the main compounds responsible for the antiacetylcholinesterase and antioxidant activity in herbal tea of *Plectranthus barbatus* ("falso boldo"), *Food Chem.*, Article in press, doi:10.1016/j.foodchem.2008.10.015.

Fiske, C.H. and Subbarow, Y. (1925), The colorimetric determination of phosphorus, *J. Biol. Chem.*, 66:375-400.

Folch, J., Lees, M. and Solane, S.G.H. (1957), A simple method for isolation and purification of total lipids from animal tissues, *J. Biol. Chem.*, 26:497-509.

Fong, S.D., Alello, L., Gardner, W.T., King, L.G., Blankenship, G., Cavallerano, D.J., Ferris, L.F. and Klein, R. (2004), Retinopathy in diabetes, *Diabetes Care*, 27:S84-S87.

Gancedo, J.M. and Gancedo, C. (1971), Fructose-1,6-bisphosphatase, phosphofructokinase and glucose-6-phosphatase dehydrogenase from fermenting and non-fermenting yeasts, *Arch. Microbiol.*, 76:132-138.

Goldberg, I.J. (2004), Why does diabetes increase atherosclerosis? I don't know!, *J. Clin. Invest.*, 114:613-615.

Good, C.A., Krames, H. and Somogy, M. (1933), Chemical procedure for analysis of polysaccharides, Meth. Enzymol., 8:34.

Gopalan, C., Sastri, B.V.R. and Balasubramanian, S.C. (1993), Nutritive value of Indian foods, NIN, ICMR, Hyderabad, 48-49.

Green, L.C., Wagner, D.A., Glogowski, J., Skipper, P.L., Wishnok, J.S. and Tannenbaum, S.R. (1982), Analysis of nitrate, nitrite and (^{15}N) nitrite in biological fluids, Anal. Biochem., 126:131-138.

Greenblatt, D.J., Leigh-Pemberton, A.R. and Moltke, V.L.L. (2006), *in vitro* interactions of water-soluble garlic components with human cytochromes P450, J. Nutr., 136:806-809.

Gupta, S., Shukla, R., Prabhu, K.M., Agarwal, S., Rusla, U. and Murthy, P.S. (2003), Acute and chronic toxicity studies on partially purified hypoglycemic preparation from water extract of bark of *Ficus bengalensis*, Indian J. Clin. Biochem., 17:58-63.

Gupta, R., Bajpai, G.K., Johri, S. and Saxena, A.M. (2008), An overview of Indian novel traditional medicinal plants with antidiabetic potentials, AJTCAM, 5:1-7.

Haan, J.B., Witting, P.K., Stefanovic, N., Pete, J., Daskalakis, M., Kola, I., Stocker, R. and Smolich, J.J. (2006), Lack of the antioxidant glutathione peroxidase-1 does not increase atherosclerosis in C57BL/J6 mice fed a high-fat diet, J. Lipid Res., 47:1157-1167.

Habib, M.Y., Islam, M.S., Awal, M.A. and Khan, M.A. (2005), Herbal products: A novel approach for diabetic patients, *Pakistan J. Nut.*, 4:17-21.

Habig, W.H., Pabst, M.J. and Jakoby, W. (1974), The first enzymatic step in mercapturic acid IV formation, *J. Biol. Chem.*, 249:7130-7139.

Hales, C.N. and Randle, P.J. (1963), Immunoassay of insulin with insulin-antibody precipitate, *Biochem. J.*, 88:137-143.

Halliwell, B. and Gutteridge, J.M.C. (1999), Free radicals in biology and medicine, 3rd edition, Oxford University Press, Oxford, pp. 237-248.

Harnafi, H. and Amrani, S. (2008), Spectrophotometric methods for determination of plant polyphenols content and their antioxidant activity assessment: an overview, *Pharmacognosy*, 12:20-22.

Hayder, N., Skandrani, I., Kilani, S., Bouhlef, I., Abdelwahed, A., Ammar, R.B., Mahmoud, A., Ghedira, K. and Chekir-Ghedira, L. (2007), Antimutagenic activity of *Myrtus communis* L. using the *Salmonella* microsome assay, *J. S. Afr. Bot.*, 74:121-125.

Hedge, J.E. and Hofreiter, B.T. (1962), Carbohydrate chemistry, 17th eds. Academic press, New York, pp. 34-40.

Heever, D.V.E. and Venter, S.L. (2006), Nutritional and medicinal properties of *Cleome gynandra*, *ISHS Acta Hort.*, 752.

Helmstadter, A. (2008), *Syzygium cumini* (L.) skeels (*Myrtaceae*) against diabetes- 125 years of research, *Pharmazie*, 63:91-101.

Hemalatha, S., Wahi, A.K., Singh, P.N. and Chansouria, J.P.N. (2004), Hypoglycemic activity of *Withania coagulans* Dunal in streptozotocin-induced diabetic rats, *J. Ethnopharmacol.*, 93:261-264.

Hippisley-Cox, J. and Pringle, M. (2004), Prevalence, care and outcomes for patients with diet-controlled diabetes in general practice: crosssectional survey, *Lancet*, 363:423-428.

Hjelm, M. and de Verdier, C.H. (1963), A methodological study of the enzymatic determination of glucose in blood, *Scand. J. Clin. Lab. Invest.*, 15:415-428.

Hosoda, K., Wang, M., Liao, M., Chuang, C., Iha, M., Clevidence, B. and Yamamoto, S. (2003), Antihyperglycemic effect of Oolong tea in type 2 diabetes, *Diabetes care*, 26:1714-1718.

Hu, D., Jablonski, K.A., Sparling, Y.H., Robbins, D.C., Lee, E.T., Welty, T.K. and Howard, B.V. (2003), Accuracy of lipoprotein lipids and apoproteins in predicting CHD in diabetic American Indians: The strong heart study, *Ann. Epidemiol.*, 12:79-85.

IDI, (2008), Diabetic complications, 7:13.

Idris, I., Thomson, G.A. and Sharma C.J. (2006), Diabetes mellitus and stroke, *Intl. J. Clin. Pract.*, 60:48-56.

Iniaghe, O.M., Malomo, S.O., Adebayo, J.O. and Arise, R.O. (2008), Evaluation of the antioxidant and hepatoprotective properties of the methanolic extract of *Acalypha racemosa* leaf in carbon tetrachloride-treated rats, Afr. J. Biotechnol., 7:1716-1720.

Irshad, S. and Chaudhuri, P.S. (2002), Oxidant-antioxidant system: Role and significance in human body, Indian J. Exp. Biol., 40:1233-1239.

Jachak, S.M. (2002), Herbal drugs as antidiabetic: an overview, CRIPS, 3:9-13.

Jackson, M.C. (1973), Soil chemical analysis, Prentice Hall of India Pvt. Ltd., pp. 461-464.

Jonas, M., Reicher-Reiss, H., Boyko, V., Behar, S. and Grossman, E. (2003), Hospital and 1-year outcome after acute myocardial infarction in patients with diabetes mellitus and hypertension, J. Hum. Hypertens., 17:665-670.

Joshi, S.G. (2000), Text book of medicinal plants, IBH Publishing Co., New Delhi, pp.119.

Kakkar, P., Das, B. and Viswanathan, P.N. (1984), A modified spectrophotometric assay of SOD, Indian J. Biochem. Biophysics, 21:130-132.

Kamalakannan, N. and Prince, S.M.P. (2003), Effect of *Aegle marmelos* Correa (Bael) fruit extract on tissue antioxidants in streptozotocin diabetic rats, Indian J. Exp. Biol., 41:1285-1288.

Kameswararao, R., Guiri, R., Kesavulu, M.M. and Apparao, C.H. (2001), Effect of oral administration of bark extracts of *Pterocarpus santalinus* L. on blood glucose level in experimental animals, J. Ethnopharmacol., 74:69-74.

Kameswararao, B., Kesavulu, M.M. and Apparao, C.H. (2003), Evaluation of antidiabetic effect of *Momordica cymbalaria* fruit in alloxan-diabetic rats, Fitoterapia, 74:7-13.

Kamiya, K., Saiki, Y., Hama, T., Fujimoto, Y., Hanani, E., Umar, M. and Satake, T. (2001), Flavonoid glucuronides from *Helicteres isora*, Phytochem., 57:297-301.

Karthikeyan, J. and Rani, P. (2003), Enzymatic and non-enzymatic antioxidants in selected *Piper* species, Indian J. Exp. Biol., 41:135-140.

Karthikumar, S., Vigneswari, K. and Jegatheesan, K. (2007), Screening of antibacterial and antioxidant activities of leaves of *Eclipta prostrata* (L), SRE, 2:101-104.

Kesari, A.N., Gupta, R.K. and Watal, G. (2005), Hypoglycemic effect of *Murraya koenigii* on normal and alloxan-diabetic rabbits, J. Ethnopharmacol., 97:247-251.

Kessler, M., Ubeaud, G. and Jung, L. (2003), Anti and pro-oxidant activity of rutin and quercetin derivatives, J. Pharm. Pharmacol., 55:131-142.

Ketz, J. (2001), A review of oral antidiabetic agents, Pharmacotherapy Update, 4:3.

Khalaf, A.N., Shakya, K.A., Othman, A.A., Agbar, Z.E. and Farah, H. (2008), Antioxidant activity of some common plants, Turk. J. Biol., 32:51-55.

Khanum, F., Bawa, A.S. and Singh, B. (2005), *Rhodiola rosea* : a versatile adaptogen- Comprehensive review, Food Sci. Food Safety, 4:55-62.

Kim, J.H., Feng, C., Wang, X. and Rajapakse, N. (2006), Effect of methyl jasmonate on secondary metabolites of sweet basil (*Ocimum basilicum* L.), J. Agri. Food chem., 154:2327-32.

King, K.J. and Wootton, I.D.P. (1956), Micromethods in medical Biochemistry, 3rd edition, Churchill, London, p. 61.

Kirthikar, K.R. and Basu, B.D. (1995), Indian medicinal plants, 1st edition, International book distributors, Dehradun, India, pp. 371-372.

Kirthikar, K.R. and Basu, B.D. (2000), Illustrated Indian medicinal plants, Vol. 7, Sri Satguru publications, Delhi, India, p.2241.

Kochhar, A., Nagi, M. and Sachdev, R. (2006), Proximate composition, available carbohydrates, dietary fibre and anti nutritional factors of selected traditional medicinal plants, J. Hum. Ecol., 19:195-199.

Koida, H. and Oda, T. (1959), Pathological occurrence of glucose-6-phosphatase in liver disease, Clin. Chem. Acta, 4:554-561.

Kornberg, A. and Horecker, B.L. (1955), Methods of enzymology, Vol.1, Academic Press, New York, p. 323.

Kretchmer, O.A., Dahan, Y.G., Yadun, L.S., Gollop, R. and Hayyim, G. (2004), The salt stress signal transduction pathway that activates the GPx1 promoter is mediated by intracellular H₂O₂ different from the pathway induced by extracellular H₂O₂, *Plant Physiol.*, 135:1-12.

Krishna, G. and Ranjan, S.K. (1991), *Special analytical techniques in nutritional biochemistry*, Kalyani publishers, New Delhi, pp. 61-81.

Kumar, G., Banu, S.G., Murugesan, A.G. and Pandian, R.M. (2006a), Hypoglycaemic effect of *Helicteres isora* bark extracts in rats, *J. Ethnopharmacol.*, 107:304-307.

Kumar, G., Murugesan, A.G. and Pandian R.M. (2006b), Effect of *Helicteres isora* bark extract on blood glucose and hepatic enzymes in experimental diabetes, *Pharmazie*, 61:353-355.

Kumar, S.P.G., Arulselvan, P., Kumar, D.S. and Subramanian, S.P. (2006c), Antidiabetic activity of fruits of *Terminalia chebula* on streptozotocin-induced diabetic rats, *J. Health Sci.*, 52:283-291.

Kumar, G. and Murugesan, A.G. (2007), Influence of *Helicteres isora* bark extracts on plasma and tissue glycoprotein components in streptozotocin diabetic rats, *J. Clin. Diag. Res.*, 4:330-338.

Kumar, A.R., Subburathinam, K.M. and Prabakar, G. (2007a), Phytochemical screening of selected medicinal plants of *Asclepiadaceae* family, *Asian J. Microbiol. Biotech. Env. Sci.*, 9:177-180.

Kumar, G., Banu, S.G., Murugesan, A.G. and Pandian, R.M. (2007b), Antihyperglycaemic and antiperoxidative effect of *Helicteres isora* L. bark extracts in streptozotocin-induced diabetic rats, J. Appl. Biomed., 5:97-104.

Kumar, G. and Murugesan, A.G. (2008), Hypolipidaemic activity of *Helicteres isora* L. bark extracts in streptozotocin-induced diabetic rats, J. Ethnopharmacol., 116:161-166.

Kumar, G., Banu, S.G. and Murugesan, A.G. (2008), Effect of *Helicteres isora* bark extracts on heart antioxidant status and lipid peroxidation in streptozotocin diabetic rats, J. Appl. Biomed., 6:89-95.

Kumaran, A. and Karunakaran, R.J. (2007), Antioxidant activity of *Cassia auriculata* flowers, Fitoterapia, 78:46-47.

Lai, P.K. (2004), Antimicrobial and chemopreventive properties of herbs and spices, Curr. Med. Chem., 52:1451-1460.

Lavhale, M.S. and Mishra, S.H. (2007), Evaluation of free radical scavenging activity of *Butea monosperma* Linn., Indian J. Exp. Biol., 45:376-384.

Lee, S., Kim, J.S., Oh, S.H., Park, K.Y., Lee, H.G. and Kim, S.D. (2008), Antihyperglycemic effect of *Fomitopsis pinicola* extracts in streptozotocin-induced diabetic rats, J. Med. Food, 11:518-524.

Lemhadri, A., Zeggwagh, N.A., Maghrani, M., Jouad, H. and Eddouks, M. (2004), Anti-hyperglycaemic activity of the aqueous extract of *Origanum vulgare* growing wild in Tafilalet region, J. Ethnopharmacol., 92:251-256.

Lopes-Virella, M.F. (1977), Assay for lipid peroxides in animal tissues by TBARS, *Annal. Biochem.*, 95:351-358.

Lowey, A., Moore, S., Norris, C., Wright, D., Silcock, J. and Hammond, P. (2007), The cost-effectiveness of pharmacist-led treatment of cardiac risk in patients with type 2 diabetes, *Pharm. World Sci.*, 29:541-545.

Lowry, O.H., Rosenbrough, N.J., Farr, A.L. and Randall, R.J. (1951), Protein measurment with Folin phenol reagent, *J. Biol. Chem.*, 193:265.

Luck, H. (1974), In: *Methods in enzymatic analysis-2* (Ed. Bergmeyer), Academic Press, New York, p. 885.

Ludvik, B., Neuffer, B. and Pacini, G. (2004), Efficacy of *Ipomoea batatas* (Caiapo) on diabetes control in type 2 diabetic subjects treated with diet, *Diabetes care*, 27:436-440.

Luo, J.Z. and Luo, L. (2008), Ginseng on hyperglycemia: effects and mechanisms, *eCAM Advance Access*, 1-5.

Madhu, G.C., Dharma, C.P. and Devi, B.D. (2005), Effect of vitamin E supplementation on diabetes induced oxidative stress in experimental diabetes in rats, *Indian J. Exp. Biol.*, 43:177-180.

Mahdi, A.A., Chandra, A., Singh, R.K., Shukla, S., Mishra, L.C. and Ahmad, S. (2003), Effect of herbal hypoglycemic agents on oxidative stress and antioxidant status in diabetic rats, *IJCB*, 18:8-15.

Mair, R.D. and Hall, T. (1977), In: Organic Peroxides, Swem, D.C. and Witky (eds), Intersciences, Wikyced, New York, p. 532.

Malloy, H.T. and Evelyn, K.A. (1937), The determination of bilirubin with the photoelectric colorimeter, J. Biol. Chem., 119:481-490.

Manjunatha, H. and Srinivasan, K. (2007), Hypolipidemic and antioxidant effects of curcumin and capsaicin in high fat fed rats, Can. J. Physiol. Pharmacol., 85:588-596.

Maritim, A.C., Sanders, R.A. and Watkins, J.B. (2003), Diabetes oxidative stress and antioxidants: A review, J. Biochem. Mol. Toxicol., 17:24-38.

Maron, D.M. and Ames, B.N. (1983), Revised methods for the *Salmonella* mutagenicity test, Mutation Research, 113:173-215.

Maroo, J., Vasu, V.J. and Gupta, S. (2003), Dose dependent hypoglycemic effect of aqueous extract of *Enicostemma littorale* Blume in alloxan-induced diabetic rats, Phytomed., 10:196-199.

Mc Cann, J., Choi, E., Yamasaki, E. and Ames, B.N. (1975), Detection of carcinogens as mutagens in the *Salmonella* microsome test: assay of 300 chemicals, Proc. Natl. Acad. Sci., USA, 72:5135-5139.

Mensor, L.L., Menezes, F.S., Leitao, G.G., Reis, A.S., Dossantos, T., Coube, C.S. and Ceitao, S.G. (2001), Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method, Phytother. Res., 15:127-130.

Metin, G., Atukeren, P., Gumustas, K.M., Belce, A. and Kayserilioglu, A. (2002), The effect of vitamin E treatment on oxidative stress generated in trained rats, *Tohoku J. Exp. Med.*, 198:47-53.

Misra, H.P. and Fridovich, A. (1972), Assay of superoxide dismutase, *J. Biol. Chem.*, 247:3170-3171.

Mishra, J., Srivastava, R.K., Shukla, S.V. and Raghav, C.S. (2007), Antioxidants in aromatic and medicinal plants, *Science Tech Entrepreneur*, 1-16.

Mohan, V. (2007), Macrovascular component of diabetes atherosclerosis and insulin (CUPS-18), *Japi.*, 55.

Mondal, S.K., Chakraborty, G., Gupta, M. and Mazumder, U.K. (2005), Hepatoprotective activity of *Diospyros malabarica* bark in carbon tetrachloride intoxicated rats, *Eur. Bull. Drug Res.*, 13:25-30.

Mori, M.D., Baviera, M.A., Ramalho, O.T.L., Venramini, C.R., Brunetti, L.I. and Pepato, T.M. (2003), Temporal response pattern of biochemical analyses in experimental diabetes, *Biotechnol. Appl. Biochem.*, 38:183-191.

Moron, M.S., De Pierre, J.N. and Mannervik, V. (1979), Levels of glutathione, glutathione reductase, glutathione-S-transferase activities in rat lung and liver, *Biochem. Biophys. Acta.*, 582:67-68.

Mukherjee, P.K., Maiti, K., Mukherjee, K. and Houghton, P.J. (2006), Leads from Indian medicinal plants with hypoglycemic potentials, J. Ethnopharmacol., 106:1-28.

Mullen, B.A., Young, I.S., Free, H. and McCance, D.R. (2002), Ascorbic acid reduces blood pressure and arterial stiffness in type 2 diabetes, Hypertension, 40:804-809.

Murray, K.R., Granner, D.K. and Rodwell, W.V. (2006), Harper's illustrated Biochemistry, 27th edition, McGraw Hill, International edition, pp. 155-160, 173.

Mythilypriya, R., Shanthi, P. and Sachdanandam, P. (2007), Oral acute and subacute toxicity studies with Kalpaamrutha, a modified indigenous preparation on rats, J. Health Sci., 53:351-358.

Nadkarni, A.K. (2000), In India, Materia medica, Vol I, Popular Prakashan Ltd., Mumbai, India, p. 116.

Nalamolu, K.R., Krishna, M.B. and Nammi, S. (2004), Effect of chronic administration of *Boerhaavia diffusa* Linn. leaf extract on experimental diabetes in rats, TJPR, 3:305-309.

Narendhirakannan, R.T., Subramanian, S. and Kandaswamy, M. (2005), Free radical scavenging activity of *Cleome gynandra* L. leaves on adjuvant induced arthritis in rats, Mol. Cell. Biochem., 276:71-80.

Netelson, S. (1957), Microtechniques of clinical chemistry for the routine laboratory, C.C. Thomas, Springfield, Illinois, p. 381.

Nichans, W.J. Jr. and Samuelson, B. (1968), Formation of malondialdehyde from phospholipid arachidonate during microsomal lipid peroxidation, *Eur. J. Biochem.*, 6:126-130.

Odukaya, O.A., Agha, S.I.I., Segun, F.I., Sofidiya, M.O. and Ilori, O.O. (2007), Antioxidant activity of selected Nigerian green leafy vegetables, *Am. J. Food Tech.*, 2:169-175.

Ogbonnia, S., Adekunle, A.A., Bosa, M.K. and Enwuru, V.N. (2008), Evaluation of acute and subacute toxicity of *Alstonia congensis* Engler (*Apocynaceae*) bark and *Xylopia aethiopica* (Dunal) A. Rich (*Annonaceae*) fruit mixtures used in the treatment of diabetes, *Afr. J. Biotechnol.*, 7:701-705.

Okhawa, H., Ohisi, N. and Yagi, K. (1979), Assay for lipid peroxidation in animal tissues by thiobarbituric acid reactions, *Annal. Biochem.*, 95:351.

Olanlokun, J.O. (2008), Protective influence of vitamin E on the antioxidant defence system in the whole blood and liver of normal and alloxan-induced diabetic rats, *Indian J. Clin. Bio.*, 23:62-66.

Oleszek, W. and Stochmal, A. (2002), Triterpene saponins and flavonoids in the seeds of *Trifolium* species, *Phytochem.*, 61:165-170.

Omaye, S.T., Turbul, T.D. and Sauberlich, H.C. (1979), Selected method for the determination of ascorbic acid in animal cells, tissues and fluids, Mc. Cormic, D.B, Wright, D.L. (eds), *Methods of enzymology*, Academic Press, New York, pp. 3- 11.

Orzechowski, A. (2003), Justification for antioxidant preconditioning (or how to protect insulin mediated actions under oxidative stress), J. Bio. Sci., 28:39-49.

Oser, B.L. (1971), Food analysis, Hawks Physiological Chemistry, 14th Edn. Mc Graw Hill, New York, pp. 1094-1098.

Osinubi, A.A., Ajay, O.G. and Adesiyun, A.E. (2006), Evaluation of the antidiabetic effect of aqueous leaf extract of *Tapinanthus butungii* in male Sprague-Dawley rats, Med. J. Islamic World Acad. Sci., 16:41-47.

Owen, J.A., Iggo, B., Scandrett, F.J. and Stewart, C.P. (1954), The determination of creatinine in plasma or serum and in urine; a critical examination, Biochem. J., 58:426-437.

Padma, P. (2001), Effect of *Annona muricata* and *Polyalthia cerasoides* on brain neurotransmitters and enzyme monoamine oxidase following cold immobilization stress, J. Natural Remedies., 1:144-146.

Palanivel, R., Thangavel, M., Selvendran, K. and Sakthisekaran, D. (2001), Insulino mimetic effect of ammonium paratungstate on protein metabolism in streptozotocin-induced diabetic rats, Biomed., 21:23-30.

Parham, M., Amini, M., Aminorroya, A. and Heidarian, E. (2008), Effect of zinc supplementation on microalbuminuria in patients with type 2 diabetes: a double blind, randomized, placebo controlled, cross-over trial, Rev. Diabet. Stud., 5:102-109.

Pari, L. and Latha, M. (2002), Effect of *Cassia auriculata* flowers on blood sugar levels, serum and tissue lipids in streptozotocin-induced diabetic rats, Singapore Med. J., 43: 617-621.

Pari, L. and Latha, M. (2004), Effect of *Scorparia dulcis* (Sweet Broomweed) plant extract on plasma antioxidants in streptozotocin-induced experimental diabetes in male albino Wistar rats, Pharmazie, 59:557-560.

Pari, L. and Murugan, P. (2007), Antihyperlipidemic effect of curcumin and tetrahydrocurcumin in experimental type 2 diabetic rats, J. Med. Food, 29:881-889.

Pari, L. and Satheesh, M.A. (2004), Antidiabetic activity of *Boerhaavia diffusa* L. effect on hepatic key enzymes in experimental diabetes, J. Ethnopharmacol., 91:109-13.

Peter, R., Cox, A. and Evans, M. (2008), Management of diabetes in cardiovascular disease patients, Heart, 94:369-375.

Petersen, M. and Simmonds, M.S.J. (2003), Rosmarinic acid, Phytochem., 62:121-125.

Petrolonis, A.J., Yang, Q., Tummino, P.J., Fish, S.M., Pruck, A.E., Jain, S., Parsons, T.F., Dales, N.A., Ge, L., Langston, S.P., Schuller, A.G., Tartaglia, L.A., Chen, H. and Hong, S.B. (2004), Enzymatic characterization of the pancreatic islet-specific glucose-6-phosphatase related protein (IGRP), J. Biol. Chem., 279:13976-13683.

Pohocha, N. and Grampurohit, N.D. (2001), Antispasmodic activity of the fruits of *Helicteres isora* Linn., *Phytother. Res.*, 15:49-52.

Prajapati, N.D., Purohit, S.S., Sharma, A.K. and Kumara, T. (2003), A handbook of medicinal plants, Edition: Agrobios Publishers, Jodhpur, pp. 188,189.

Prakasam, A., Sethupathy, S. and Pugalendi, K.V. (2002), Antihyperglycaemic activity of *Casearia esculenta* root extract in blood glucose and plasma antioxidant status in streptozotocin-induced diabetic rats, *Pharmazie*, 57:758-760.

Proks, P., Reimann, F., Green, N., Gribble, F. and Ashcroft, F. (2002), Sulfonylurea stimulation of insulin secretion, *Diabetes*, 51:S368-S376.

Punitha, I.S.R., Rajendran, K., Shirwaikar, A. and Shirwaikar, A. (2005), Alcoholic stem extract of *Coscinium fenestratum* regulates carbohydrate metabolism and improves antioxidant status in streptozotocin- nicotinamide induced diabetic rats, *Advance Access Publication*, 2:375-381.

Qu, W.H., Li, J.G. and Wang, M.S. (1991), Chemical studies on the *Helicteres isora*, *Zhongguo Yaoke Daxue Xuebao.*, 22:203-206.

Radha, V., Vimalaswaran, K.S., Deepa, R. and Mohan, V. (2003), The genetics of diabetes mellitus, *Indian J. Med. Res.*, 117:225-238.

Radin, N. (1981), In: *Methods in enzymology*, Lowenstein, J.M. (ed.), Academic Press, New York, pp.72.

Raghunath, R., Rai, E. and Madhavi, S. (2006), Plasma oxidant- antioxidant status in different respiratory disorders, Indian J. Clin. Biochem., 21:161-164.

Raghuramulu, N., Nair, M.K. and Kalyanasundaram, S. (1983), A manual of laboratory techniques, 1st edition, NIN, KMR, Hyderabad, pp. 31,32.

Rajagopal, K. and Saikala, K. (2008), Antidiabetic activity of hydro-ethanolic extracts of *Nymphaea stellata* flowers in normal and alloxan-induced diabetic rats, Afr. J. Pharm. Pharmacol., 2:173-178.

Rajendran, A., Narayanan, V. and Gnanavel, I. (2007), Study on the analysis of trace elements in *Aloe vera* and its biological importance, JASR, 3:1476-1478.

Ramachandran, A., Snehalatha, C., Baskar, A.D., Mary, S., Kumar, C.K., Selvam, S., Catherine, S. and Vijay, V. (2004), Temporal changes in prevalence of diabetes and impaired glucose tolerance associated with lifestyle transition occurring in the rural population in India, Diabetologia, 47:860-865.

Ramkumar, K.M., Rajaguru, P., Latha, M. and Ananthan, R. (2007), Ethanol extract of *Gymnema montanum* leaves reduces glycoprotein components in experimental diabetes, Nutr. Res., 27:97-103.

Ramprasath, V.R., Kumar, B.S., Shanthi, P. and Sachdanandam, P. (2006), Therapeutic efficacy of Kalpaamruthaa on lipid peroxidation and antioxidant status in experimental memory carcinoma in rats, J. Health Sci., 52:748-757.

Rani, H.S., Madhavi, G., Rao, V.R., Sahay, B.K. and Jyothy, A. (2005), Risk factors for the coronary heart disease in type 2 diabetes mellitus, Indian J. Clin. Biochem., 20:75-80.

Rattanachaikunsopon, P. and Phumkhachorn, P. (2007), Bacteriostatic effect of flavonoids isolated from leaves of *Psidium guajava* on fish pathogen, Fitoterapia, 78:434-436.

Raut, N.A. and Gaikwad, N.J. (2006), Antidiabetic activity of hydro-ethanolic extract of *Cyperus rotundus* in alloxan-induced diabetes in rats, Fitoterapia, 77:585-588.

Ravi, K., Ramachandran, B. and Subramanian, S. (2004), Effect of *Eugenia jambolana* seed kernel on antioxidant defense system in streptozotocin-induced diabetes in rats, Life Sci., 75:2717-2731.

Ravi, K., Rajasekaran, S. and Subramanian, S. (2005), Antihyperlipidemic effect of *Eugenia jambolana* seed kernel on streptozotocin-induced diabetes in rats, Food Chem. Toxicol., 43:1433-1439.

Ravi, S., Sadashiva, C.T., Tamizmani, T., Balasubramanian, T. and Rubeshkumar, M. (2009), *in vitro* glucose uptake by isolated rat hemi-diaphragm study of *Aegle marmelos* Correa root, BDPS, 4:65-68.

Reddy, K.P., Subhani, K.H., Khan, P.A. and Kumar, K.B. (1985), Effect of light and benzyladenine on dark-treated growing rice (*Oryza sativa*) leaves II changes in peroxidase activity, Plant Cell Physiol., 26:987-994.

Reitman, S. and Frankel, S. (1957), A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases, *Am. J. Clin. Pathol.*, 28:56-63.

Ribnicky, D.M., Poulev, A., Watford, M., Cefalu, W.T. and Raskin, I. (2006), Antihyperglycaemic activity of TarralinTM, an ethanolic extract of *Artemisia dracunculus* L., *Phytomed.*, 13:550-557.

Riose, J.L., Simon, S., Jimeney, F.J., Zafrapolo, M.C. and Villas, A. (1986), Reagents for screening medicinal plants by Thin layer chromatography- A review, *Fitoterapia*, 57:153-162.

Roberts, K. (2001), Bitter melon: Ready for mainstream medicine? *Sun Integrative Health*, 6-7.

Roe, J.H. and Kuether, C.A. (1953), The determination of ascorbic acid in whole blood and urine through 2, 4 dinitro phenylhydrazine derivative of dehydro ascorbic acid, *J. Biol. Chem.*, 147:399-407.

Rosenberg, H.R. (1992), *Chemistry and physiology of the vitamins*, Interscience publishers Inc, New York, pp. 452-453.

Rotruck, J.T., Pope, A.L., Ganther, H.E. and Swanson, A.B. (1984), Selenium: Biochemical roles as a component of glutathione peroxidase, *Science*, 179:588-590.

Ruiz-Teran, F., Martínez, A.M. and Ocana, A.N. (2008), Antioxidant and free radical scavenging activities of plant extracts used in traditional medicine in Mexico, *Afr. J. Biotechnol.*, 7:1886-1893.

Sabu, M.C. and Kuttan, R. (2000), Antidiabetic activity of some medicinal plants-relation with their antioxidant property, Amala research Bulletin, 20:81-86.

Sabu, M.C. and Kuttan, R. (2004), Anti-diabetic activity of *Aegle marmelos* and its relationship with its antioxidant properties, Indian J. Physiol. Pharmacol., 48:81-88.

Sampathkumar, R., Sivakumar, T., Shanmugasundaram, R., Sivakumar, P., Nethaji, R., Gupta, M. and Mazumdar, K.V. (2006), Antimicrobial and antioxidant activities of *Careyg arborea* Roxb. stem bark, IJPT, 5:35-41.

Sanderson, J.H. and Philips, C.E. (1981), An atlas of laboratory, Animal haematology, Clarendon Press, Oxford, p. 473.

Satake, T., Kamiya, K., Sazki, Y., Fujimoto, Y., Kitanake, S., Kimura, Y., Uzariva, J., Endang, H. and Umar, M. (1999), Studies on the constituents of fruits of *Helicteres isora* L., Chem. Pharm. Bull., 47:1444-1447.

Satdive, R.K., Abilash, P. and Fulzele, D.P. (2003), Antimicrobial activity of *Gymnema sylvestre* leaf extract, Fitoterapia, 74:699-701.

Satyanarayana, T., Katayani, B.M., Hemalatha, E., Anjana, A.M. and Eswaraiah, M.C. (2006), Hypoglycemic and anti hyperglycemic effect of alcoholic extract of *Euphorbia leucophylla* and its fractions in normal and in alloxan induced diabetic rats, PHCOG MAG., 2:244-253.

Schanderl, S.H. (1970), Methods in food analysis, Edition: Academic Press, New York, p. 709.

Seo, K.I., Choi, M.S., Jung, U.J., Kim, H.J., Yeo, J., Jeon, S.M and Lee, M.K. (2008), Effect of curcumin supplementation on blood glucose, plasma insulin and glucose homeostasis related enzyme activities in diabetic db/db mice, *Mol. Nutr. Food Res.*, 52:995-1004.

Seth, S.D. and Sharma, B. (2004), Medicinal plants in India, *Indian J. Med. Res.*, 120:9-11.

Sharma, B., Balomajumder, C. and Roy, P. (2008), Hypoglycemic and hypolipidemic effects of flavonoid rich extract from *Eugenia jambolana* seeds on streptozotocin-induced diabetic rats, *Food Chem. Toxicol.*, 46:2376-2383.

Shetty, A., Kumar, G., Sambaiah, K. and Salimath, P. (2003), Effect of bitter gourd (*Momordica charantia*) on glycaemic status in streptozotocin induced diabetic rats, *Plant Foods Hum. Nutr.*, 60:109-112.

Shi, Y., So, K.F., Man, R.Y.K. and Vanhoutte, P.M. (2007), Oxygen-derived free radicals mediated endothelium-dependent contractions in femoral arteries of rats with streptozotocin-induced diabetes, *Brit. J. Pharmacol.*, 152:1033-1041.

Shinoda, J. (1928), Color reactions of flavone and flavone like substances, *J. Pharmaceut. Society, Japan*, 48:214.

Shirwaikar, A., Rajendran, K. and Kumar, D.C. (2004), Oral antidiabetic activity of *Annona squamosa* leaf alcohol extract in NIDDM rats, *Pharmaceut. Biol.*, 42:30-35.

Siddique, O., Sun, Y., Lin, J.C. and Chien, Y.W. (1987), Facilitated transdermal transport of insulin, J. Pharm. Sci., 76:341-345.

Silva, J.A.F., Lorencini, M., Peroni, L.A., De La Hoz, C.L.R., Carvalho, H.F. and Stach-Machado, D.R. (2007), The influence of type I diabetes mellitus on the expression and activity of gelatinases (matrix metalloproteinases-2 and -9) in induced periodontal disease, J. Periodontal Res., 1- 2.

Singh, S.S., Pandey, S.C., Srivastava, S., Gupta, V.S., Patro, B. and Ghosh, A.C. (2003), Chemistry and medicinal properties of *Tinospora cordifolia* (Guduchi), IJP, 35:83-91.

Sinha, K.A. (1972), Colorimetric assay of catalase, Annal. Biochem., 47:389-394

Somani, R.S. and Singhai, A.K. (2008), Hypoglycaemic and antidiabetic activities of seeds of *Myristica fragrans* in normoglycaemic and alloxan-induced diabetic rats, Asian J. Exp. Sci., 22:95-102.

Somova, L.I., Shoda, F.O., Ramnanan, P. and Nadar, A. (2003), Antihypertensive, antiatherosclerotic and antioxidant activity of triterpenoids isolated from *Olea europaea*, subspecies *africana* leaves, J. Ethnopharmacol., 84:299-305.

Sridhar, S.B., Sheetal, U.D., Pai, M.R.S.M. and Shastri, M.S. (2005), Preclinical evaluation of the antidiabetic effect of *Eugenia jambolana* seed powder in streptozotocin diabetic rats, Brazl. J. Med. Biol. Res., 38:463-468.

Srinivasan, K., Muruganadhan, S. and Lal, J. (2001), Evaluation of anti-inflammatory activity of *Pongamia pinnata* leaves in rats, J. Ethnopharmacol., 78:151-157.

Steve, O.O., Joy, O. and Veronica, E.N. (2008), Evaluation of hypoglycaemic and hypolipidaemic effects of aqueous ethanolic extracts of *Treculia africana* Decne and *Bryophyllum pinnatum* Lam. and their mixture on streptozotocin (STZ)-induced diabetic rats, Afr. J. Biotechnol., 7:2535-2539.

Tapsell, L.C., Hemphill, I., Cobiac, L., Patch, C.S., Sullivan, D.R., Fenech, M., Roodenry, S.S., Keogh, J.B., Clifton, P.M., Williams, P.G., Fazio, V.A. and Inge, K.E. (2006), Health benefits of herbs and spices: the past, the present, the future, Med. J. Aust., 185:S4-24.

Tepe, B., Eminagaaghu, O., Akpulat, H.A. and Aydin, E. (2007), Antioxidant potentials and rosmarinic acid levels of the methanolic extracts of *Salvia verticillata* (L) subspecies *verticullate* and *Salvia verticullata* (L) subspecies *anasiaca* (Freyn and Bornm) Bornm, Food Chem., 100:985-989.

Tezuka, Y., Terazono, M., Kusumoto, I.T., Kawashima, Y., Hatanaka, Y., Kadota, S., Hattori, M., Namba, T., Kikuchi, T., Tanaka, K. and Supriyatna, S. (1999), Helisterculins A and B, two new (7.5', 8.5')-neolignans and helisorin, the first(6.4', 7.5', 8.2')-neolignan, from the Indonesian medicinal plant *Helicteres isora*, HVCA, 82:408-417.

The Medica, (2008), The effect of 8 dietary supplements on diabetes, 2-4.

Thomson, M., Al-Amin, Z.M., Al-Quttan, K.K., Shaban, L.H. and Ali, M. (2007), Anti-diabetic and hypolipidaemic properties of garlic (*Allium sativum*) in streptozotocin-induced diabetic rats, Intl. J. Diabetes and Metabolism, 15:108-115.

Tiwari, A.K. and Rao, M. (2002), Diabetes mellitus and multiple therapeutic approaches of phytochemicals: present status and future prospects, Curr. Sci., 83:30-38.

Toth, J., Mrlianova, M., Tekelova, D. and Korenova, M. (2003), Rosmarinic acid– an important phenolic active compound of lemon balm (*Melissa officinalis* L.), Acta Facult. Pharm. Univ. Comenianae, 50:139-146.

Tripathy, K.D. (1994), Adverse drug effects. In: essentials of Medical Pharmacology, 3rd Edition, Jaypee Brothers-Medical Publishers Ltd. New Delhi, pp. 65-69.

Ugochukwu., N.H. and Babady, N.E. (2003), Antihyperglycemic effect of aqueous and ethanolic extracts of *Gongronema latifolium* leaves on glucose and glycogen metabolism in livers of normal and streptozotocin-induced diabetic rats, Life Sci., 73:1925-1938.

Umpierrez, G.E., Murphy, M.B. and Kitabchi, A.E. (2002), Diabetic ketoacidosis and hyperglycemic hyperosmolar syndrome, Diabetes Spectrum, 15:28-36.

Umpierrez, G.E. and Kitabchi, A.E. (2003), Diabetic Ketoacidosis: risk factors and management strategies, Treatments in Endocrinology, 2:95-108.

Van Handel, E. and Zilversmit, D.B. (1957), Micromethod for the direct determination of serum triglycerides, J. Lab. Clin. Med., 50:152-157.

van Hecke, M.V., Decker, J.M. and Nijpels, G. (2006), Are retinal microvascular abnormalities associated with large endothelial dysfunction and intima-media thickness: the Hoorn study, Clin. Sci., 110:597-604.

Varughese, G.I., Patel, J.V., Lip, G.Y. and Varma, C. (2006), Novel concepts of statin therapy for cardiovascular risk reduction in hypertension, Curr. Pharm. Des., 12:1593-609.

Varughese, G.I. and Scarpello, J.H.B. (2006), Treating dyslipidaemia in the setting of diabetes mellitus and cardiovascular disease: a less commonly perceived therapeutic perspective in clinical practice, Journal compilation Blackwell Publishing Ltd., Intl. J. Clin. Pract., 60:884-88.

Venkatesh, S., Reddy, D.G. and Reddy, B.M. (2003), Antihyperglycemic activity of *Helicteres isora* roots in alloxan-induced diabetic rats, Pharm. Biol., 41:347-350.

Venkatesh, S.G., Reddy, D.Y.S.R., Sathyavathy, R.D. and Reddy, B.M. (2004), Effect of *Helicteres isora* root extracts on glucose tolerance in glucose- induced hyperglycemic rats, Fitoterapia, 75:364-367.

Venkatesh, S., Sailakshmi, K., Reddy, B. and Ramesh, M. (2007), Antimicrobial activity of *Helicteres isora* root, Indian J. Pharma. Sci., 69:687-689.

Venukumar, M.R. and Latha, M.S. (2002), Antioxidant activity of *Curculigo orchioides* in carbon tetrachloride induced hepatopathy in rats, IJCB, 17:80-87.

Vessby, J., Basu, S., Mohsen, R., Berne, C. and Vessby, B. (2002), Oxidative stress and antioxidant status in type 1 diabetes mellitus, Blackwell Science Ltd., JIM, 251:69-76.

Wahab, A.H.Y., O'harte, F.P.M., Mooney, H.M., Barnett, R.C. and Flatt, R.P. (2002), Vitamin C supplementation decreases insulin glycation and improves glucose homeostasis in obese hyperglycemic (ob/ob) mice, Metabolism, 51:514-517.

Wahren, J., Ekberg, K. and Jornvall. H. (2007), C-peptide is a bioactive peptide, Diabetologia, 50:503-509.

Wang, T.C., Ti, M.C., Lo, S.C. and Yang, C.C. (2007), Free radical-scavenging activity of aqueous extract of *Pteris multifida* Poiret, Fitoterapia, 78:248-249.

Warnick, G.R. and Albers, J.J. (1978), A comprehensive evaluation of the heparin-manganese precipitation procedure for estimating high density lipoprotein cholesterol, J. Lipid Res., 19:65-76.

Warrier, P.K., Nambrani, V.P.K. and Ramankutty, C. (2004), Indian medicinal plants: A compendium of 500 species, Orient Longmen Publishers, p. 113.

Watanabe, D., Nakara, H., Akagi, K., Ishii, T., Mizuguchi, H., Nagashima, Y. and Okaniwa, A. (2004), Oral glucose tolerance test and determination of serum fructosamine level in Beagle dogs, J. Toxicol. Sci., 29:33-36.

WHO, (2006), Definition and diagnosis of diabetes mellitus and intermediate hyperglycemia, Fact sheet No: 312.

WHO, (2008), Diagnosis and classification of diabetes mellitus, Geneva, 59.

Wild, S., Roglic, G., Green, A., Sicree, R. and King H. (2004), Global prevalence of diabetes: estimates for the year 2000 and projections for 2030, Diabetes Care, 27:1047-1053.

Winterbourn, C.C., Hawkins, R.E., Brain, M. and Carrell, R.W. (1975), The estimation of red cell superoxide dismutase activity, J. Lab. Clin. Med., 85:337-341.

Wohlrab, J., Wohlrab, D. and Meiss, F. (2007), Skin diseases in diabetes mellitus, JDDG, 5:37-53.

Wong, M.C., Chung, J.W.Y. and Wong, T.K.S. (2007), Effects of treatments for symptoms of painful diabetic neuropathy: systematic review, BMJ, 335:87.

Xia, T. and Wang, Q. (2006), Anti hyperglycemic effect of *Cucurbita ficifolia* fruit extract in streptozotocin-induced diabetic rats, Fitoterapia, 77:530-533.

Yale, J.F., Valiquatt, R.T., Ghazzi, N.M., Grillo, O.K.J., Whitcomb, W.R. and Foyt, L.H. (2001), The effect of a thiazolidinedione drug, troglitazone, on glycemia in patients with type 2 diabetes mellitus poorly controlled with sulfonylurea and metformin, *Ann. Internal Med.*, 134:737-745

Yoganarasimhan, S.N. (2000), *Medicinal plants of India - Tamilnadu*, Vol. II, International book publisher, Print Cyber Media, Bangalore, pp. 104-208.

Zachary, T. and Bloomgarden, M.D. (2002), The epidemiology of complications, *Diabetes care*, 25:924-932.

Zafrilla, P., Mulero, J., Xandri, J.M., Santo, E., Caravaca, G. and Morillas, J.M. (2006), Oxidative stress in Alzheimer's disease in different stages of the disease, *Curr. Med. Chem.*, 13:1075-1083

Zakaria, H., Simpson, K., Brown, P.R. and Krotulovic, A. (1979), Use of reverse phase HPLC analysis for the determination of provitamin A carotenes in tomatoes, *J. Chromatography.*, 176:109-117.

Zhang, B.B. and Moller, D.E. (2000), New approaches in the treatment of type 2 diabetes, *Curr. Opin. Chem. Biol.*, 4:461-467.

Zhang, R., Zhou, J., Jia, Z., Zhang, Y. and Gu, G. (2003), Hypoglycemic effect of *Rehmannia glutinosa* oligosaccharide in hyperglycemic and alloxan-induced diabetic rats and its mechanism, *J. Ethnopharmacol.*, 90:39-43.

Zhu, B.H., Guan, Y.Y., He, H. and Lin, M.J. (2000), Effects of scutellarein on diabetic rat aorta, *Acta Pharmacol. Sin.*, 21:353-356.

Zilversmit, D.B. and Davis, A.K. (1950), Microdetermination of phospholipids by TCA precipitation, *J. Lab. Clin. Med.*, 35:155-159.

Zimmerman, B.R. (2000), Postprandial hyperglycemia: Implications for Practia, *Am. J. Cardiol.*, 88:32H-36H.

Website

www.cstl.nist.gov/div831/DNATechologies/imgs/DNA_Damage_Repair.jpg.