

BIBLIOGRAPHY

BIBLIOGRAPHY

- Ahmed, Z., Sheikh, M.A., Hameed, A. and Salah ud Din. (2012), Investigation of Antioxidant Enzymes and Biochemical Changes in the Wheat Seeds (Freed) Induced by Different Pre-Sowing Treatments, *World Applied Sciences Journal*, 18 (1): 31-36.
- Al-Hinai, A.H., Al-Sadi, A.M., Al-Bahry, S.N., Mothershaw, A.S., Al-Said, F.A., Al-Harhi, S.A. and Deadman, M.L. (2010), Isolation and characterization of *Pseudomonas aeruginosa* with antagonistic activity against *Pythium aphanidermatum*, *Journal of Plant Pathology*, 92(3), 653-660.
- Almagro, L., Gomez Ros, L.V., Belchi-Navarro, S., Bru, R., Barcelo, A.R. and Pedren˜o, M.A. (2009), Class III peroxidases in plant defence reactions : Review paper, *Journal of Experimental Botany*, 60(2) ; 377–390.
- Almoneafy, A.A., Xi, G.L., Tian, W. X., Xu, L. H., Zhang, G.Q. and Ibrahim, M. (2012), Characterization and evaluation of Bacillus isolates for their potential plant growth and biocontrol activities against tomato bacterial wilt, *African Journal of Biotechnology*, 11(28); 7193-7201.
- Amara, M.T.A., Arafa, R.A., El-Sheref, E.M., Sidkey, N.M. and Khalifa, R.Y. (2009), Biological control of sore shin disease of cotton by fluorescent pseudomonad's isolated from egyptian desert soil, *Egyptian Journal of Natural Toxins*, 6(2); 32-64.
- Anis, M., Zaki, M.J. and Dawar, S. (2012), Development of a Na-alginate-based bioformulation and its use in the management of charcoal rot of sunflower (*Helianthus annuus* l.), *Pak. J. Bot.*, 44(3): 1167-1170.
- Anita, B. and Samiyappan, R. (2012), Induction of systemic resistance in rice by *Pseudomonas fluorescens* against rice root knot nematode *Meloidogyne graminicola*, *J Biopest*, 5 (Supplementary): 53-59.
- Anitha, A. and Rabeeth, M. (2009), Control of *Fusarium* Wilt of Tomato by Bioformulation of *Streptomyces griseus* in Green House Condition, *African Journal of Basic & Applied Sciences*, 1 (1-2): 9-14.

- Ardebili, Z.O., Ardebili, N.O. and Hamdi., S.M.M. (2011), Physiological effects of *Pseudomonas fluorescens* CHA0 on tomato (*Lycopersicon esculentum* Mill.) plants and its possible impact on *Fusarium oxysporum* f. sp. *Lycopersici*, *Australian Journal of crop science*, 5(12):1631-1638.
- Ashnaei, S.P., Tehrani, A.S., Ahmadzadeh, M. and Behboudi, K. (2008), Production of *Pseudomonas fluorescens* P-5 and P-6 for Bean Damping-off Disease, *International Journal of Agriculture and Biology*, 10: 573–576.
- Ashrafuzzaman, M., Hossen, F.A., Ismail, R.M., Hoque, A.Md., Islam, Z.M., Shahidullah, S.M. and Meon, S. (2009), Efficiency of plant growth-promoting rhizo bacteria, (PGPR) for the enhancement of rice growth, *African Journal of Biotechnology*, 8(7), 1247-1252.
- Babu, M. A., Singh, D. and Gothandam, K.M. (2012), The effect of salinity on growth, hormones and mineral elements in leaf and fruit of tomato cultivar PKM1, *The Journal of Animal & Plant Sciences*, 22(1), 159-164.
- Bakrim, A., Lamhamdi, M., Sayah, F. and Chibi, F. (2007), Effects of plant hormones and 20-hydroxyecdysone on tomato (*Lycopersicum esculentum*) seed germination and seedlings growth, *African Journal of Biotechnology*, 6 (24), 2792-2802.
- Bhagat, I. and Chakraborty, B. (2010), Defense response triggered by *Sclerotium rolfsii* in tea plants, *Ecoprint*, 17: 69-76.
- Bilkay, I.S., Karakoç, S. and Aksöz, N. (2010), Indole-3-acetic acid and gibberellic acid production in *Aspergillus niger*, *Turk J Biol*, 34: 313-318.
- Bolto K. & Rohrlach T. 2010 – Using Gibberellic Acid to Promote Winter Pasture Growth Mallee Sustainable Farming 2009 Research Compendium, 118 – 121.
- Celetti, M. (2012), Plant Pathologist - Horticulture Crops Program Lead, Fire blight prediction for May 28 – June 3, 2012, Hort matters, OMAFRA Specialists in Horticulture and Specialty Crops, 12(14): 4.
- Chakravarty, G. and Kalita, M.C. (2011), Comparative evaluation of organic formulations of *Pseudomonas fluorescens* based biopesticides and their application in the management of bacterial wilt of brinjal (*Solanum melongena* L.), *African Journal of Biotechnology*, 10(37): 7174-7182.

- Chang, A., Myung-Ho, L., Shin-Woo, L., Robb, E.J. and Nazar, R.N. (2008), Tomato Phenylalanine Ammonia-Lyase Gene Family, Highly Redundant but Strongly Underutilized, *The journal of biological chemistry*, 283(48): 33591–33601.
- Chen, C., Belanger, R.R., Benhamou, N. and Paulitz, T.C. (2000), Defense enzymes induced in cucumber roots by treatment with plant growth promoting rhizobacteria (PGPR) and *Pythium aphanidermatum*, *Physiological and Molecular Plant Pathology* 56: 13–23.
- Chérif, M., Arfaoui, A. and Rhaiem, A. (2007), Phenolic Compounds and their Role in Bio-control and Resistance of Chickpea to Fungal Pathogenic Attacks : Review Article, *Tunisian Journal of Plant Protection*, 2: 7-21.
- Christopher, D.J., Raj, T.S., Rani, S.U. and Udhayakumar, R. (2010), Role of defense enzymes activity in tomato as induced by *Trichoderma virens* against *Fusarium* wilt caused by *Fusarium oxysporum* f sp.lycopersici, *Journal of Biopesticides*, 3(1): 158 – 162.
- Dai, J. and Mumper, J.R. (2010), Plant Phenolics: Extraction, Analysis and Their Antioxidant and Anticancer Properties: Review, *Molecules*, 15: 7313-7352.
- Dar, W.A., Bhat, J.A., Rashid, R., Rehman, S. and Bhat, Z.A. (2012), Bioefficacy of *Pseudomonas fluorescens* in management of damping-off disease in papaya (*Carica papaya* L.), *Journal of Agricultural Technology*, 8(2): 693-697.
- Dickerson, D.P., Pascholati, S.F., Hagerman, A.E., Butler, L.G. and Nicholson, R.L. (1984), Phenylalanine ammonia- lyase and hydroxyl cinnamate CoA ligase in maize mesocotyls inoculated with *Helminthosporium matdis* or *Helminthosporium carbonum*, *Physiological plant pathology*, 25: 111-123.
- Doornbos, F.R., Cornelis van Loon, L. and Bakker, A.H.M.P. (2012) Impact of root exudates and plant defense signaling on bacterial communities in the rhizosphere. A review, *Agron. Sustain. Dev.* 32:227–24.
- El-beltagi, S.H., Farahat, A.A., Alsayed, A.A. and Mahfoud, A.N. (2012), Response of Antioxidant Substances and Enzymes Activities as a Defense Mechanism Against Root-Knot Nematode Infection, *Not Bot Horti Agrobo*, 40(1): 132-142.

- Gao, G., Yin, D., Chen, S., Xia, F., Yang, J., Li, Q. and Wang, W. (2012) Effect of Biocontrol Agent *Pseudomonas fluorescens* 2P24 on Soil Fungal Community in Cucumber Rhizosphere Using T-RFLP and DGGE, *PLoS one*, 7(2): 1-9.
- Gladden, L.A., Yu-Min, W., Ching-Hsiang, H and Tsou, I. (2012), Using deficit irrigation approach for evaluating the effects of water restriction on field grown tomato (*Lycopersicon esculentum*), *African Journal of Agricultural Research*, 7(14), 2083-2095.
- Gordon, A.S. and Weber, P.R. (1951), Colorimetric estimation of indole acetic acid, *Plant physiology*, 192-195.
- Gordon, S.A. and Paleg, L.G. (1957), Quantitative measurement of indole acetic acid, *Physiological Plant Pathology*, 10: 347-348.
- Gravel, V., Martinez, C., Antoun, H. and Tweddell, R.J. (2005), Antagonist microorganisms with the ability to control *Pythium* damping-off of tomato seeds in rockwool, *BioControl*, 50:771–786.
- Hammerschmidt, R., Nuckles, E.M. and Kuc, J. (1982), Association of enhanced peroxidase activity with induced systemic resistance of cucumber to *Colletotrichum lagenarium*. *Physiological Plant Pathology*, 20: 73-82.
- Hyun, M.W., Yun, Y.H., Kim, J.Y. and Kim, S.H. (2011) Fungal and Plant Phenylalanine Ammonia-lyase: Review article, *Mycobiology*, 39(4) : 257-265.
- Jack and Nelson (2010), Final report to the Organic Farming Research Foundation, <http://www.hort.cornell.edu/soilhealth/>
- Jenana, R.K.B., Haouala, R., Triki, M.A., Godon, J.J., Hibar, K., Khedher, M.B. and Henchi, B. (2009), Composts, compost extracts and bacterial suppressive action on *Pythium aphanidermatum* in tomato, *Pak. J. Bot.*, 41(1): 315-327.
- Jorjani, M., Heydari, A., Zamanizadeh, H.R., Rezaee, S., Naraghi, L. and Zamzami, P. (2012), Controlling sugar beet mortality disease by application of new bioformulations, *Journal of Plant Protection Research*, 52(3): 303-307.

- Kader, M.M.A., El-Mougy, N.S., Aly, M.D.E. and Lashin, S.M. (2012), Long Activity of Stored Formulated Bio-Agents Against Some Soil-Borne Plant Pathogenic Fungi Causing Root Rot of Some Vegetables, *Journal of Applied Sciences Research*, 8(4): 1882-1892.
- Kavina, J., Gopi, R. and Panneerselvam, R. (2011) Traditional and non- traditional Plant Growth Regulators alter the growth and Photosynthetic Pigments in *Mentha piperita* Linn., *International Journal of environmental sciences*, 1 (7): 124-134.
- Khan, T.N, Jeelani, G., Tari, S., Mahmood, T. and Hussain, S.J. (2011) Effect of different concentrations of rooting hormones on growth of tomato cuttings (*Solanum esculentusl.*), *J. Agric. Res.*, 2011, 49(2): 241-247.
- King, , E.O., Ward, M.K. and Raney, D.E. (1954), Two simple media for the demonstration of pyocyanin and fluorescein, *Journal of Laboratory and Clinical Medicine*, 44:301-307.
- Kumar, A., Kumar, A., Devi, S., Patil, S., Payal, C. and Negi, S. (2012) Isolation, screening and characterization of bacteria from Rhizospheric soils for different plant growth promotion (PGP) activities: an *in vitro* study, *Recent Research in Science and Technology*, 4(1): 01-05
- Laemmli, U.K. (1970), Cleavage of structural proteins during the assembly of the head of bacteriophage T4, *Nature*, 227: 680-685.
- Liang, J., Tao, R., Hao, Z., Wang, L. and Zhang, X. (2011), Induction of resistance in cucumber against seedling damping-off by plant growth-promoting rhizobacteria (PGPR) *Bacillus megaterium* strain L8, *African Journal of Biotechnology*, 10(36): 6920-6927.
- López-Ráez, A.J., Flors, V., García, M.J. and Pozo, J.M. (2010), AM symbiosis alters phenolic acid content in tomato roots, *Plant Signaling & Behavior* 5(9): 1138-1140.
- Mahadevan, A. and Sridhar, R (1982), Hydrolytic enzymes in: *Methods in Physiological plant pathology*, 2nd Edition, Sivakami Publishers, Madras, 24-44.

- Maisuria, K. M. and Patel, S. T. (2009), Seed germinability , root and shoot length and vigour index of soybean as influenced by rhizosphere fungi, *Karnataka J. Agric. Sci.*, 22 (5), 1120-1122.
- Manonmani, K., Bhaskaran, R., Jagannathan, R. and Jayasekhar, M. (2009), Biochemical changes in acid lime during pathogenicity of canker pathogen, *Indian J. Agric. Res.*, 43(3) : 157-165.
- Matthews, V. D. (1931) Studies on the Genus *Pythium*, Chapel Hill, NC, University of North Carolina Press.
- Mayer, A.M., Harel, E. and Shaul, R.B. (1965). Assay of catechol oxidase a critical comparison of methods, *Phytochemistry*, 5: 783-789.
- Mazid, M., Khan, T.A. and Mohammad, F. (2011), Role of secondary metabolites in defense mechanisms of plants: Review Article, *Biology and Medicine*, 3(2): 232-249.
- Morsy, E.M., Abdel-Kawi, K.A. and Khalil, M.N.A. (2009), Efficiency of *Trichoderma viride* and *Bacillus subtilis* as Biocontrol Agents against *Fusarium solani* on Tomato Plants, *Egypt. J. Phytopathol*, 37(1): 47-57.
- Muthukumar, A., Eswaran, A. and Sanjeevkumas, K. (2011) Exploitation of trichoderma species on the growth of *Pythium aphanidermatum* chilli, *Brazilian Journal of Microbiology*, 42: 1598-1607.
- Newman, S.M., Tantasawat, P. and Steffens, J.C. (2011), Tomato Polyphenol Oxidase B Is Spatially and Temporally Regulated during Development and in Response to Ethylene, *Molecules*, 16, 493-517.
- Noori, M.S.S. and Saud, H.M. (2012), Potential Plant Growth-Promoting Activity of *Pseudomonas* sp Isolated from Paddy Soil in Malaysia as Biocontrol Agent, *J Plant Pathol Microb*, 3:2, 1-4.
- Noureen, F., Jilani, M.S., Waseem, K. and Kiran, M. (2010), Performance of tomato hybrids under hydroponic culture, *Pak. J. Agri. Sci.*, 47(1): 19-25.
- Ojha, S. and Chatterjee, N.C. (2012) Induction of resistance in tomato plants against *fusarium oxysporum f. sp. lycopersici* mediated through salicylic acid and *Trichoderma harzianum*, *Journal of plant protection research*, 52(2): 220-225.

- Omer, A.M. (2010), Bioformulations of *Bacillus* Spores for using as Biofertilizer, *Life Science Journal*, 7(4); 124-131.
- Ortiz, H.O., Mendoza, A.B., Villarreal, R.M., Rodríguez, H.R. and KarimDeAlba Romenus, (2007), Enzymatic Activity in Tomato Fruits as a Response to Chemical Elicitors, *J.Mex.Chem.Soc*, 51(3): 141-144.
- Passam, C.H., Karapanos, I.C., Bebeli, J.P. and Savvas, D. (2007), A Review of Recent Research on Tomato Nutrition, Breeding and Post-Harvest Technology with Reference to Fruit Quality, *The European Journal of Plant Science and Biotechnology*, 1(1), 1-21.
- Queiroz, C., Lopes, M.L.M., Fialho, E. and Mesquita, V.L.V. (2008), Polyphenol Oxidase: Characteristics and Mechanisms of Browning Control, *Food Reviews International*, 24: 361–375.
- Raja, P., Uma, S., Gopal, H. and Govindarajan, K. (2006), Impacts of bio inoculants consortium on rice root exudates, biological nitrogen fixation and plant growth, *Journal of biological sciences*, 6(5): 815-823.
- Raju, S., Jayalakshmi, S.K. and Sreeramulu, K. (2008), Comparative study on the induction of defense related enzymes in two different cultivars of chickpea (*Cicer arietinum*L) genotypes by salicylic acid, spermine and *Fusarium oxysporum* f. sp. *Ciceri*, *Australian Journal of Crop Science*, 2(3): 121-140.
- Riker, A.J. and Riker, R.S. (1936), Introduction to research on plant Disease, John Swift, New York. 117pp.
- Saharan, B.S. and Nehra, V. (2011), Plant Growth Promoting Rhizobacteria: A Critical Review, *Life Sciences and Medicine Research*, LSMR-21, 1-30.
- Salaheddin, K., Valluvaparidasan, V., Ladhakshmi, D. and Velazhahan, R. (2010), Management of Bacterial Blight of Cotton Using a Mixture of *Pseudomonas fluorescens* and *Bacillus subtilis*, *Plant Protect. Sci.*, 46(2): 41–50.
- Sarwar, N., Hayat zahid, M., Ashfaq, S. and Jamil, F. (2011), Induced systemic resistance in chickpea against Ascochyta blight by safe chemicals, *Pak. J. Bot.*, 43(2): 1381-1387.

- Shahab, S., Ahmed, N. and Khan, S.N. (2009), Indole acetic acid production and enhanced plant, growth promotion by indigenous PSBs, *African Journal of Agricultural Research*, 4 (11): 1312-1316.
- Suleiman, M. N. (2011), The *in vitro* chemical control of *Pythium aphanidermatum*, an agent of tomato root rots in the north central, Nigeria, *Scientia Africana*, 10(2): 48 – 54.
- Thipyapong, P., Michael J.S. and Attajarusit, J. (2007), Functional Analysis of Polyphenol Oxidases by Antisense/Sense Technology, *Molecules*, 12: 1569-1595.
- Ulrichs, C., Fischer, G., Büttner, C. and Mewis, I. (2008), comparison of lycopene, b-carotene and phenolic contents of tomato using conventional and ecological horticultural practices, and arbuscular mycorrhizal fungi (AMF), *Agronomía Colombiana*, 26(1): 40-46.
- Umashankari, J. and Sekar, C. (2011), Comparative evaluation of different bioformulations of PGPR cells on the enhancement of induced systemic resistance (ISR) in Rice *P. oryzae* pathosystem under upland condition, *Curr. Bot*, 2(3):12-17.
- Vidhyasekaran, R, and Muthamilan, M. (1995), Development of formulation of *Pseudomonas fluorescens* for control of chickpea wilt, *Plant Disease*,79: 782-786.
- Wintemans, J.F.G.M. and Demots, A. (1965), Spectrophotometer characteristics of chlorophyll a and b and their phenophytins in ethanol, *Biochem. Biophys. Acta*, 109: 448- 453.
- Zieslin, N. and Ben-Zaken, R. (1993), Peroxidase activity and presence of phenolic substances in peduncles of rose flowers. *Plant Physiology and Biochemistry*, 31: 333-339.

WEBSITES

http://www.nature.com/mt/journal/v10/n2/fig_tab/mt20041219f1.html