



BIBLIOGRAPHY

- Abbas, A., Murtaza, S., Shahid, K., Munir, M., Ayub, R. and Akber, S. (2012). Comparative study of adsorptive removal of congo red and brilliant green dyes from water using peanut shell, Middle- East Journal of Scientific Research, 11(2012), 828-832.
- Abdul-Baki, A.A. and Anderson, J.D. (1973). Vigour determination in soybean seed by multiple criteria, Crop Sci. 3(1973), 630-633.
- Abdel- Moneium, M., Abousabana, N.M., Khadre, S.E.M. and Abdel Kader, H.H. (2008). Physiological and Histopathological effects in cat fish exposed to dyestuff and chemical waste water, International Journal of Zoological Research, 4, 189-202
- Abraham, K., Sridevi, R., Suresh, B. and Damodharam, T. (2013). Effect of heavy metals (Cd, Pb and Cu) on seed germination of *Arachis hypogea*.L. Asian J of plant Science and Research, 3(1), 10-12.
- Adeogun, A. O. and Chukwuka, A. V. (2011). Effect of textile factory effluent on otolith and somatic growth parameters in *Clarias gariepinus*. Zoologist, 9, 70-77.
- Ahlawat, K. and Kumar, A. (2009), Analysis of industrial effluents and its comparison with other effluents from residential and commercial areas in Solan H.P, Journal of Theoretical and Applied Sciences, 1(2), 42-46.
- Ahmad, I., Hamid, T., Fatima, M., Chand, H. S., Jain, S. K., Athar, M., & Raisuddin, S. (2000). Induction of hepatic antioxidants in freshwater catfish (*Channa punctatus* Bloch) is a biomarker of paper mill effluent exposure. Biochimica et Biophysica Acta (BBA)-General Subjects, 1523(1), 37-48.

- Ahmed, M.A. (2012). Physicochemical Assessment of Textile Effluents in Chittagong Region of Bangladesh and Their Possible Effects on Environment, *International Journal of Research in Chemistry and Environment*, 2(3), 220-229.
- Ahmed, T.F., Sushil, M and Krishna, M. (2012), Impact of dye industrial effluent on physicochemical characteristics of Kshipra River, Ujjain City, India, *International Research Journal of Environment Sciences*, 1(2), 41-45.
- Al Dahaan, S., Al-Ansari, N. and Knutsson, S. (2016). Influence of Groundwater Hypothetical Salts on Electrical Conductivity Total Dissolved Solids. *Engineering*, 8(11), 823-830.
- Allen, J.L., Gofus, J.E. and Meinertz, J.R. (1994). Determination of malachite green residues in the eggs, fry and adult muscle tissue of rainbow trout (*Oncorhynchus mykiss*). *JAOAC Int.* 77: 553- 557.
- Alimba, C.G., Ogunkanmi, A.L. and Ogunmmola, F.J. (2013). Cytotoxic and genotoxic assessment of textile effluent using *Allium* assay. *Current Topics in Toxicology*. 9: 65-74.
- Alvarado, L., Torres, I. R., & Chen, A. (2013). Integration of ion exchange and electrodeionization as a new approach for the continuous treatment of hexavalent chromium wastewater. *Separation and Purification Technology*, 105, 55-62.
- Alwan, S.F. (2009). Effect of aluminum exposure on some blood parameters in the *Tilapia zillii* fish, *International Journal of Pharmacy & Life Sciences*, 4(7), 2766-2769.
- Amin, N.K. (2009). Removal of direct blue-106 dye from aqueous solution using new activated carbons developed from pomegranate peel: adsorption equilibrium and kinetics, *J. Hazard. Mater.*, 15(165), 52-62.
- Angelin, A. J., Jones , R.D. S. and Das, S.S.M. (2015), Investigation of acute toxicity and the effect of chrome plating industry effluent on behavior of the guppy (*Poecilia reticulata*), *European Journal of Biotechnology and Bioscience*, 3 (9), 01-05.

- APHA, AWWA and WEF. Standard Methods for the Examination of Water and Wastewater, 21st edn, Washington (2005).
- Arora, S. (2014). Textile Dyes: It's Impact on Environment and its Treatment, Journal of J Bioremediation & Biodegradation, Arora, J Bioremed Biodeg, 5,3 DOI: 10.4172/2155-6199.1000e146.
- Arul, J.M., Revathi, M.N. and Saravanan, J. (2011). Decolorization and Physicochemical analysis of textile azodye by Bacillus, International Journal of Applied Bioengineering, 5: 35-39.
- Asgher, M. and Bhatti, H.N. (2012). Evaluation of thermodynamics and effect of chemical treatments on sorption potential of citrus waste biomass for removal of anionic dyes from aqueous solutions. Ecological Engineering, 38, 79-85.
- Asgher, M. and Bhatti, H.N. (2012). Removal of reactive blue 19 and reactive blue 49 textile dyes by citrus waste biomass from aqueous solution: equilibrium and kinetic study, The Canadian Journal of Chemical Engineering, 90, 412-419.
- Asgher, M., Bhatti, H.N., Ashraf, M and Legge, R.L. (2008). Recent developments in biodegradation of industrial pollutants by white rot fungi and their enzyme system, Biodegradation, 19, 771– 783.
- Ashish S. Sartape., Aniruddha M. Mandhare., Vikas V. Jadhav., Prakash D. Raut ., Mansing A. Anuse and Sanjay S. Kolekar. (2014). Removal of malachite green dye from aqueous solution with adsorption technique using Limonia acidissima (wood apple) shell as low cost adsorbent Arabian Journal of Chemistry. pp: 1-10.
- Ashraf, H and Hussain, Q. (2010). Studies on bitter gourd peroxidase catalyzed removal of P- bromophenol from wastewater. Desalination, 262 (1-3), 267-272.
- Asouhidou, D.D., Triantafyllidis, K.S., Lazaridis, N.K., Matis, K.A., Kim, S.S. and Pinnavia, T.J. (2009). Sorption of reactive dyes from aqueous solutions by ordered hexagonal and disordered mesoporous carbons, Microporous and Mesoporous Materials. 117, 257-267.

- Aswin, M. (2015). Biosorption of Textile Effluent Using *Sargassum longifolium*. International Journal of Advance Research in Science and Engineering, 4(01), ISSN-2319-8354.
- Bakhiet, S.H.H.A. (2015). Determination of Heavy Metals in Fish Tissues and Water from White Nile Khartoum City –Sudan, Journal of Environment Protection and Sustainable Development, 1(3), 178-181.
- Bakker, P., Van Doorne, H., Gooskens, V. and Wieringa, N.F. (1992). Activity of gentian violet and brilliant green against some microorganisms associated with skin infections. Int J Dermatol , 31, 210-213.
- Bal, A.S. (1999), Waste water management for textile industry- an overview, Indian Journal of Environmental Health, 41 (4): 264-290.
- Balarak, D., Bazrafshan, E. and Mostafapour, F.K. (2016). Equilibrium, Kinetic studies on the adsorption of Acid Green 3 (Ag₃) dye onto *Azolla filiculoides* as adsorbent, American Chemical Science Journal, 11, 1-10.
- Bhanu, D.A. and Deepak, M. (2013). The influence of Textile Mill Effluent on Haematological Changes of *Labeo rohita*. International Journal of Innovative Research in Science, Engineering and Technology. 2(12): 8001 – 8003.
- Barot, J. and Bahadur, A. (2013). Behavioural and histopathological effects of azodye on kidney and gills of *Labeo rohita* fingerlings. Journal of environmental biology, 34(2), 147.
- Bassyouni, D. G., Hamad, H. A., El-Ashtoukhy, E. Z., Amin, N. K. and El-Latif, M. A. (2017). Comparative performance of anodic oxidation and electrocoagulation as clean processes for electrocatalytic degradation of diazo dye Acid Brown 14 in aqueous medium. Journal of hazardous materials, 335, 178-187.
- Behera, B. and Rathna, R.V. (2002), Economic and Political Weekly, Jan.19.
- Benguella, B. and Benaissa, H. (2002), Cadmium removal from aqueous solutions by chitin: kinetic and equilibrium studies, Water Re- search, 36, 2463–2474.

- Bhatkar, N. V. (2011). Chromium, nickel and zinc induced histopathological alterations in the liver of Indian common Carp *Labeo rohita* (Ham.). *Journal of Applied Sciences and Environmental Management*, 15(2), 331 – 336.
- Bhattacharyya, K.G, Sarma, A. (2003) Adsorption characteristics of the dye, brilliant green, on neem leaf powder. *Dyes Pigments*, 57:211–222.
- Bobmanuel, N. O. K. , Gabriel U. U. and Ekweozor I. K. E. (2006), Direct toxic assessment of treated fertilizer effluents to *Oreochromis niloticus*, *Clarias gariepinus* and catfish hybrid (*Heterobranchus bidorsalis* and *Clarias gariepinus*), *African Journal of Biotechnology*, 5 (8), 635-642.
- Bold, H.C. and Wynne, M.J. (1978). Introduction to the algae: Structure and reproduction, Englewood Cliffs, New Jersey, Prentice Hall, Inc, 706.
- Boudechiche, N., Mokaddem, H., Sadaoui, Z. and Trari, M. (2015). Biosorption of cationic dye from aqueous solutions onto lignocellulosic biomass (*Luffa cylindrica*): characterization, equilibrium, kinetic and thermodynamic studies, *Int J Ind Chem*, DOI 10.1007/s40090-015-0066-4.
- Brahmabhatt, N.H. and Jasrai, R.T. (2016) Bidegradation of reactive dyes by two microalgal species, 4(5), 1697-1702.
- Brahmabhatt, N.H. and Jasrai, R.T. (2016). The Role of Algae in Bioremediation of Textile Effluent, *International Journal of Engineering Research and General Science*. 4, 21- 28.
- Brusle, J. and Anadon, O.I.G. (1996). The structure and function of fish liver: In: *Fish Morphology*, (Eds) J.S.D. Munshi and H.M. Dutta. Science Publishers Inc.
- Chou, C.H., Chaiang, C. and Kao, C.I. (1978). Impacts of water pollution on crop growth in Taiwan, *Bot. Bull. Sinica*, 19, 107-124.
- Cha, S. M., Nagaraju, G., Sekhar, S. C., Bharat, L. K. and Yu, J. S. (2018). Fallen leaves derived honeycomb-like porous carbon as a metal-free and low-cost counter

electrode for dye-sensitized solar cells with excellent tri-iodide reduction. *Journal of colloid and interface science*, 513, 843-851.

- Chapman, D.J. and Gellenbeck, K.W. (1983). An historical prospective of algal biotechnology. In: *Algae and cyanobacterial biotechnology*, Longman group, U.K, 1-27.
- Cock, J.G. (1964). *Handbook of textile fibers*, Merrow Publication Co. Ltd., England.
- Correia, V.M. (1998). Sulphanated surfactants and related compounds: Fact of their desulfonation by aerobic and anaerobic bacteria, *Tenside surfactants detergents*, 35, 52- 56.
- Culp, S.J. and Beland, F.A. (1996). Malachite green a toxicological review. *J. Am. Coll. Toxicol.* 15: 219-238.
- Dahri, M. K., Lim, L. B. L., Kooh, M. R. R., and Chan, C. M. (2017). Adsorption of brilliant green from aqueous solution by unmodified and chemically modified Tarap (*Artocarpus odoratissimus*) peel. *International Journal of Environmental Science and Technology*, 14(12), 2683-2694.
- Daneshvar, E., Kousha, M., Jokar, M., Koutahzadeh, N. and Guibal, E. (2012). Acidic dye biosorption onto marine brown macroalgae: Isotherms, kinetic and thermodynamic studies. *Chemical Engineering Journal*, pp: 204–206, 225–234.
- Daneshvar, E., Vazirzadeh, A., Niazi, A., Sillanpää, M., and Bhatnagar, A. (2017). A comparative study of methylene blue biosorption using different modified brown, red and green macroalgae–Effect of pretreatment. *Chemical Engineering Journal*, 307, 435-446.
- Daneshvar, N., Ayazloo, M., Khataee, A.R. and Pourhassan, M. (2007). Biological Decolourisation of dye solution containing Malachite Green by microalgae *Cosmarium* sp, *Bioresour Technol*, 98, 1176-1182.

- Davidson, W. W. and Henry, W. W. (1969). *Clinical Diagnosis by laboratory Methods*, 14th ed. London, Toronto: W. B. Saunders.
- de Castro Silva, F., da Silva, M. M. F., Lima, L. C. B., Osajima, J. A. and da Silva Filho, E. C. (2018). Modifying cellulose with metaphosphoric acid and its efficiency in removing brilliant green dye. *International journal of biological macromolecules*, 114, 470-478.
- de Souza, S.G., Peruzzo, L.C. and de Souza, A.U. (2008). Numerical study of the adsorption of dyes from textile effluents, *Applied Mathematical Modelling*, 32, 1711-1718.
- Deniz, F. and Saygideger, S.D. (2010). Equilibrium, kinetic and thermodynamic studies of Acid Orange 52 dye biosorption by *Paulownia tomentosa* Steud. leaf powder as a low-cost natural biosorbent, *Bioresource Technology* 101 (2010) 5137–5143.
- Deokar, R. and Sabale, A. (2015). Biosorption of methylene blue from aqueous solution onto green seaweeds, *International Journal of Recent Scientific Research*, 5, 347-351.
- Dhanve, R., Jadhav, J. and Govindwar, S. (2014), A study of textile effluent ecotoxicity and its biodegradation by an *Exiguobacterium* sp. RD3, *Int. J. Curr. Biotechnol*, 2(4):45-50.
- Durairaj, S. and Durairaj, S. (2012). Colour removal from textile industry waste water using low cost adsorbents, *International Journal of Chemical, Environmental and Pharmaceutical Research*, 3, 52-57.
- Duran, C., Ozdes, D., Gundogdu, A. and Senturk, H. B. (2011). Kinetics and isotherm analysis of basic dyes adsorption onto almond shell (*Prunus dulcis*) as a low cost adsorbent. *Journal of Chemical & Engineering Data*, 56(5), 2136-2147.
- Ebrahiem, E. E., Al-Maghrabi, M. N., & Mobarki, A. R. (2017). Removal of organic pollutants from industrial wastewater by applying photo-Fenton oxidation technology. *Arabian Journal of Chemistry*, 10, S1674-S1679.

- El-Kassas, H. Y. and Mohamed, L. A. (2014). Bioremediation of the textile waste effluent by *Chlorella vulgaris*. The Egyptian Journal of Aquatic Research, 40(3), 301-308.
- Ellman, G.L. (1959). Tissue sulphhydryl groups. Arch Biochem Biophys, 82: 70-77.
- Esan, O.S., Abiola, O.N., Owoyomi, O., Aboluwoye, C.O. and Osundiya, M.O. (2014). Adsorption of Brilliant Green onto Luffa Cylindrical Sponge: Equilibrium, Kinetics, and Thermodynamic Studies, ISRN Physical Chemistry, 2014, 1-12.
- F. Deniz and S.D. Saygideger, Equilibrium, kinetic and thermodynamic studies of Acid Orange 52 dye biosorption by *Paulownia tomentosa* Steud. leaf powder as a low-cost natural biosorbent, Bioresource technology. 101(2010) 5137-5143.
- Fan, J., Chen, D., Li, N., Xu, Q., Li, H., He, J. and Lu, J. (2018). Adsorption and biodegradation of dye in wastewater with Fe₃O₄@ MIL-100 (Fe) core-shell bio-nanocomposites. Chemosphere, 191, 315-323.
- Fatima N, Omm-e-Hany, Alamgir A (2015) Decolourization of Textile Effluent through Indigenous Microorganism. Bulletin of environment, pharmacology and life sciences: 4(5), 109-115.
- Fiskesjo, G. (1985). The allium test as standard in environmental monitoring. Hereditas, 102, 99-112.
- Francis, R. and Portia, G.A. (2013), Impact of sago effluent on the levels of acid phosphatase activity in the liver tissue of the fresh water fish *Clarias batrachus*, Int. J. Bioassays, 02 (10), 1388-1391.
- Freundlich, H.M.F. (1906). Over the adsorption in solution, J Phys Chem. 57, 385-470.
- Gajare, S.M. and Menghani, S. (2012). Biosorption of malachite green by naturally grown algal biomass from Girna River, Jalgaon District, Maharashtra. J. Algal Biomass Utiln. 3 (4): 60-65. ISSN: 2229- 6905.

- Ganesan, R. and Thanasekaran, K. (2011), Decolourisation of textile dyeing wastewater by modified solar photo fenton oxidation, *International Journal of Environmental Sciences*, 1(6), 1168-1176.
- Garg, S.K. and Tripathi, M. (2013). Process parameters for decolourization and biodegradation of orange II (Acid Orange 7) in dye-simulated minimal salt medium and subsequent textile effluent treatment by *Bacillus cereus* (MTCC 9777) RMLAU1, *Environ Monit Assess*, 185, 8909–8923.
- Garg, S.K., Tripathi, M., Singh, S.K. and Tiwari. (2012). Biodecolorization of textile dye effluent by *Pseudomonas putida* SKG-1 (MTCC 10510) under the conditions optimized for monoazo dye orange II color removal in simulated minimal salt medium. *International Biodeterioration & Biodegradation* 74: 24-35.
- Garg, V.K., Amita, M., Kumar, R. and Gupta, R. (2004a). Basic dye (methylene blue) removal from simulated wastewater by adsorption using Indian Rosewood sawdust: a timber industry waste, *Dyes and pigments*, 63, 243-250.
- Garg, V.K., Kumar, R. and Gupta, R. (2004b). Removal of malachite green dye from aqueous solution by adsorption using agro-industry waste: a case study of *Prosopis cineraria*, *Dyes and Pigments*, 62, 1-10.
- Golob, V., Vinder, A., & Simonic, M. (2005). Efficiency of the coagulation/flocculation method for the treatment of dyebath effluents. *Dyes and pigments*, 67(2), 93-97.
- Gomes, K. M. S., Oliveira, M. V. G. A. D., Carvalho, F. R. D. S., Menezes, C. C. and Peron, A. P. (2013). Cytotoxicity of food dyes sunset yellow (E-110), bordeaux red (E-123), and tatzazine yellow (E-102) on *Allium cepa* L. root meristematic cells. *Food Science and Technology (Campinas)*, 33(1), 218-223.
- Haq, I., Kumari, V., Kumar, S., Raj, A., Lohani, M. and Bhargava, R. N. (2016). Evaluation of the Phytotoxic and Genotoxic Potential of Pulp and Paper Mill Effluent Using *Vigna radiata* and *Allium cepa*. *Advances in Biology*, 2016, 1-10.

- Hebeish, A., Ramadan, M.A., Abdel-Hadim, A. and AboAkeil, E. (2011). An effective adsorbent based on sawdust for removal of direct dye from aqueous solutions. *Clean Technol Environ Policy*, 13, 713- 718.
- Himri, I., Bellahcen, S., Souna, F., Belmekki, F., Aziz, M., Bnouham, M., & Saalaoui, E. (2011). A 90-day oral toxicity study of tartrazine, a synthetic food dye, in wistar rats. *International Journal of Pharmacy and Pharmaceutical Sciences*, 3, 159-169.
- Humason, G.L. (1979). *Animal tissue technique*. 4th Edn., WH Freeman and company, San Francisco, USA., pp: 3-331
- Husain, I. and Hussain, J. (2012). Ground water pollution by discharge of dyeing and printing industrial waste water in Bandi river, Rajasthan, India, *International Journal of Environment and Bioenergy*, 2(2): 100-119.
- Ilyas,S. and Rehman, A. (2013). Decolourization and detoxification of synozol red HF- 6BN azo dye, by *Aspergillus niger* and *Niogrospora* sp. *Iranian Journal of Environmental Health Science & Engineering*. 10, 12.
- Iqbal, F., Quereshi, I.Z. and Ali, M. (2004). Histopathological changes in the kidney of common carp, *Cyprinus carpio*, following nitrate exposure, *J. Fish. Biol*, 36(4), 481-488.
- Jadhav, J.P., Kalyani, D.C., Telke, A.A., Phugare, S.S. and Govindwar,S.P. (2010). Evaluation of the efficacy of a bacterial consortium for the removal of color reduction of heavy metals and toxicity from textile dye effluent. *Bioresource technology*, 101, 165-173.
- Jagruti, B. (2015). Evaluation of azo dye toxicity using some hematological and histopathological alterations in fish *Catla catla*, *International Journal of Biological, Biomolecular, Agricultural food and Biotechnological Engineering*, 9(5), 422-425.
- Jayakumar, R., Manivasagan, R. and Karthikeyan, C. (2014), Sorption of hexavalent chromium from aqueous solution using marine green algae *Halimeda*

gracilis: Optimization, equilibrium, kinetic, thermodynamic and desorption studies, Journal of Environmental Chemical Engineering 2(3):1261–1274

- Jayaraj, R., Mohan, M.C., Prasath, P.M.D. and Khan, T.H. (2011), Malachite Green Dye Removal Using the Seaweed *Enteromorpha*, E-Journal of Chemistry, 8(2), 649-656.
- Jerold, M., Vidya, E.V., Ramya, S., Naveenji, A. and Sivasubramanian, V. (2017) Nanoscale zerovalent iron-Sargassum swartzii biocomposites for the removal of malachite green from an aqueous solution, Separation Science and Technology, 52:6, 965-974, DOI: 10.1080/01496395.2016.1276188.
- Jody, T. and Dons, J. (2003). Meeting the challenges of swine manure management, Biocycle, 44(10), 24.
- Jolly, Y.N., Islam, A. and Mustafab, A.I. (2012), Impact of Dyeing Industry Effluent on Soil and Crop, Universal Journal of Environmental Research and Technology, 2 (6): 560-568.
- Jos, A., Repetto, G., Riosc, J.C., Hazend, M.J., Molerod, M.L., Del Pesoa, A., Salgueroa, M., Fernandez-Freired, P., Perez-Martind, J.M. and Cameanb, A. (2003). Ecotoxicological evaluation of carbamazepine using six different model systems with eighteen endpoints, Toxicol. In Vitro, 17: 525-532.
- Joshi, V.J. and Santani, D.D. (2012), Physicochemical characterization and heavy metal concentration in effluent of textile industry, Universal Journal of Environmental Research and Technology, 2(2), 93-96.
- Kahraman, S. and Yalcin, P. (2005). Antibacterial Effect of Untreated and Treated (Decolourized) Dyes by Agricultural wastes. American Journal of Biochemistry and Biotechnology, 1(1): 50-53.
- Kanawade, S.M. and Gaikwad, R.W. (2011), Removal of dyes from dye effluent by using sugarcane bagasse ash as an adsorbent, International Journal of Chemical Engineering and Applications, 2(3), 202-206.

- Karadag, D., Akgul, E., Tok, S., Erturk, F., Kaya, M. A., and Turan, M. (2007). Basic and reactive dye removal using natural and modification zeolites. *Journal of Chemical and Engineering Data*, 52, 2436–2441.
- Kaur, A., Vats, S., Rekhi, S., Bhardwaj, A., Goel, J., Tanwar, R. S. and Gaur, K. K. (2010). Physico-chemical analysis of the industrial effluents and their impact on the soil microflora. *Procedia Environmental Sciences*, 2, 595-599.
- Kavitha, C., Malarvizhi, A., Kumaran, S.S. and Ramesh, M. (2010), Toxicological Effects of Arsenate Exposure on Hematological, Biochemical and Liver Transaminases Activity in an Indian Major Carp, *Catla catla*, *Food and chemical toxicology*, 48(10):2848-54.
- Kavitha, D. and Namasivayam, C. (2007), Experimental and kinetic studies on methylene blue adsorption by coir pith carbon, *Bioresour. Technol*, 98, 14-21.
- Khalaf, M.A. (2008). Biosorption of reactive dye from textile wastewater by non-viable biomass of *Aspergillus niger* and *Spirogyra* species, *Bioresource Technology*, 99, 6631-6634.
- Kismir, Y. and Aroguz, A.Z. (2011). Adsorption characteristics of the hazardous dye brilliant green on Saklikent mud. *Chem. Eng. J.* 172, 199–206.
- Kousha, M., Daneshvar, E., Dopeikar, H., Taghavi, D. and Bhatnagar, A. (2012). Box–Behnken design optimization of Acid Black 1 dye biosorption by different brown macroalgae, *Chemical Engineering Journal*, 179, 158-168.
- Kumar, K.V. and Kumaran, A. (2005). Removal of methylene blue by mango seed kernel powder, *Biochem. Eng. J.* 27, 83-93.
- Kumar, P. S., Varjani, S. J. and Suganya, S. (2018). Treatment of dye wastewater using an ultrasonic aided nanoparticle stacked activated carbon: Kinetic and isotherm modelling. *Bioresource technology*, 250, 716-722.

- Kumaraguru, K. and Sureshkumar, P. (2014), Kinetics, Modeling, Thermodynamic Analysis and Optimization of Biosorption Parameters for Biomass of *Sargassum wighiti* using Response Surface Methodology, International Journal of ChemTech Research, 6(5), 3040-3052.
- Lade, H., Govindwar, S. and Paul, D. (2015). Low-cost biodegradation and detoxification of textile azo dye CI Reactive Blue 172 by *Providencia rettgeri* strain HSL1. Journal of Chemistry, 2015, pp.1- 10.
- Lagergran, S. (1898). About the theory of So- Called Adsorption of Soluble Substances, Kung Sven. Vetén. Hand, 24, 1.
- Langmuir, I. (1916). The constitution and fundamental properties of solids and liquids, J Am Chem Soc. 38, 2221–2295.
- Lodeiro, P., Herrero, R. and De Vicente, M.E.S. (2006). Batch desorption studies and multiple sorption–regeneration cycles in a fixed-bed column for Cd (II) elimination by protonated *Sargassum muticum*, Journal of hazardous materials, 137, 1649-1655.
- Lowry, O.H., N.J. Rosenbrough, A.L. Farr and R. Randall: Protein measurement with the folin phenol reagent. J. Biochem., 193, 265-275 (1951).
- Maharajan., Kitto, M.R., Paruruckmani, P.S. and Ganapiriya, V. (2016). Histopathology biomarker responses in Asian sea bass, *Lates calcarifer* (Bloch) exposed to copper, The Journal of Basic & Applied Zoology, 77, 21-30.
- Malaviya, P., Hali, R. and Sharma, N. (2012), Impact of dyeing industry effluent on germination and growth of pea (*Pisum sativum*), J. Environ. Biol. 33, 1075-1078.
- Mali, P.L., Maharajan, M.M., Patil, D.P. and Kulkarni, M.V (2000). Biodecolourization of members of Triphenyl methane and azo group of dyes. Journal of Scientific and Industrial Research, 59 (3), 221-224.

- Mane, V.S., Mall, I.D. and Srivastava, V.C. (2007). Kinetic and equilibrium isotherm studies for the adsorptive removal of brilliant green dye from aqueous solution by rice husk ash, *Journal of Environmental Management*, 84(4), 390–400.
- Manikandan, N., Kuzhali, S.S. and Kumuthakalavalli, R. (2011), Quality assessment of dye effluent from small scale textile units of Chinnalapatti, Dindigul district, Tamilnadu, India, *Elixir pollution* 37, 4018-4019.
- Mao, J., Won, S.W., Min, J. and Yun, Y.S. (2008). Removal of Basic Blue 3 from aqueous solution by *Corynebacterium glutamicum* biomass: Biosorption and precipitation mechanisms, *Korean Journal of Chemical Engineering*, 25.5, 1060-1064.
- Mata, Y., Blazquez, M., Ballester, A., Gonzalez, F. and Munoz, J. (2010). Studies on sorption, desorption, regeneration and reuse of sugar-beet pectin gels for heavy metal removal, *Journal of hazardous materials*, 178, 243-248.
- Maximous, N. N., Nakhla, G. F. and Wan, W. K. (2010). Removal of heavy metals from wastewater by adsorption and membrane processes: a comparative study. *World Acad. Sci. Eng. Technol*, 64, 594-599.
- McKay, G. and Ho, Y.S. (1999). Pseudo Second Order Model for Sorption Process. *Process Biochem.* 34, 451.
- Mittal, A., Kaur, D. and Mittal, I. (2008). Applicability of waste materials- bottom ash and deoiled soya- as adsorbents for the removal and recovery of a hazardous dye, Brilliant green, *Journal of colloid and Interface Science.* 326 (2008) 8-17.
- Mittal, A., Mittal, J., Malviya, A. and Gupta, V. K. (2009). Adsorptive removal of hazardous anionic dye “Congo red” from wastewater using waste materials and recovery by desorption. *Journal of Colloid and Interface Science*, 340(1), 16-26.
- Mohabansi, N.P., Tekade, P.V. and Bawankar, S.V. (2011). Physico-chemical and microbiological analysis of textile industry effluent of Wardha region, *Water Research and Development*, 1(1), 40-44.

- Mohapatra, U.K. and Singh, B.C. (1998). Inorganic Anions in Drinking Water. Collected from Different Sources in the Old Capital City of Cuttack. *Ind J Env. Pollu*, 18(7), 532-535.
- Muley, D. V., Karanjkar, D. M., & Maske, S. V. (2007). Impact of industrial effluents on the biochemical composition of fresh water fish *Labeo rohita*. *Journal of environmental biology*, 28(2), 245-249.
- Murugesan, K. and Kalaichelvan, PT. (2003), Synthetic dye decolourization by white rot fungi, *Indian journal of experimental biology*, 41 (9), 1076-1087.
- Nagarani, N., Vahitha, H. and Kumaraguru, A.K. (2012). Comparative studies on the removal of textile effluents by various eco-friendly adsorbent strategies, *Journal of Toxicological & Environmental Chemistry*, 94(7), 1259-1271.
- Nandi, B.K., Goswami, A. and Purkait, M.K. (2009). Adsorption characteristics of brilliant green dye on kaolin, *Journal of Hazardous Materials*, 161, 387-395.
- Narat, J. K. (1931). Brilliant Green: A Clinical Study of Its Value As a Local Antiseptic. *Annals of Surgery*. 94 (6), 1007–1012. doi:10.1097/00000658-193112000-00003.
- Nelson, D.A. and Morris, M.W. (1989). Basic methodology. Chap. 27. Hematology and coagulation, part IV, in: *Clinical diagnosis and management by laboratory methods*. Nelson, D.A., Henry, J.B. seventieth ed. Henry, J.B, W.B. Saunder Company., Philadelphia, U.S.A. 578-625.
- Naveen, N., Saravanan, P., Baskar, G. and Renganathan, S. (2011). Equilibrium and Kinetic modeling on the removal of Reactive Red 120 using positively charged *Hydrilla verticillata*, *Journal of the Taiwan Institute of chemical Engineers*, 42, 463- 469.
- Nemcsok J, Benedeczky I, Boross L, Asztalos B, Orban L (1981). Subcellular localization of transaminase enzymes in fishes and their significance in the detection of water pollution. *Acta Biol Szeged*, 27: 9-15.

- Nemerow, N.L. (1978). *Industrial Water Pollution: Origins, Characteristics and Treatment*. AddisonWesley, Reading, Massachusetts, pp. 738.
- Nikalje, S. B., Muley, D. V., & Angadi, S. M. (2012). Histopathological changes in gills of a freshwater major carp *Labeo rohita* after acute and chronic exposure to textile mill effluent (tme). *International Journal of Environmental Sciences*, 3(1), 108.
- Noel, S. D., and Rajan, M. R. (2015). Phytotoxic Effect of Dyeing Industry Effluent on Seed Germination and Early Growth of Lady's Finger. *Journal of Pollution Effects & Control*, 3(2), 1-4.
- Nwani, C.D., Mkpadobi, B.N., Onyishi, G., Echi, P.C., Chukwuka, C.O., Oluah, S.N. and Ivoke, N. (2014). Changes in behavior and hematological parameters of freshwater African catfish *Clarias gariepinus* (Burchell 1822) following sublethal exposure to chloramphenicol, *Drug Chem Toxicol*, 37(1):107-13.
- Ogunlaja, O. and Aemere, O. (2009). Evaluating the efficiency of a textile wastewater treatment plant located in Oshodi, Lagos, *African Journal of Pure & Applied Chemistry*, 3, 189-196.
- Ohioma, A.I., Luke, N.O. and Amraibure, O. (2009). Studies on the pollution potential of wastewater from textile processing factories in Kaduna, Nigeria. *Journal of Toxicology & Environmental Health Sciences* 1, 34-37.
- Oke, I.A.; Okuofu, C.A. and Otun, J.A. (2006). A statistical classification of textile wastewaters in Nourthern Nigeria. *Journal of Applied Sciences Research*, 2 (4), 209-216.
- Oliveira, G. A., Carissimi, E., Monje-Ramírez, I., Velasquez-Orta, S. B., Rodrigues, R. T., & Ledesma, M. T. O. (2018). Comparison between coagulation-flocculation and ozone-flotation for *Scenedesmus* microalgal biomolecule recovery and nutrient removal from wastewater in a high-rate algal pond. *Bioresource technology*, 259, 334-342.

- Ozer, A., Akkaya, G. and Turabik, M. (2005). Biosorption of Acid Red 274 (AR 274) on *Enteromorpha prolifera* in a batch system. *Journal of hazardous materials*, 126(1-3), 119-127.
- Pan, X. and Zhang, D. (2009). Removal of malachite green from water by *Firmiana simplex* wood fiber. *Electronic Journal of Biotechnology*, 12(4), 9-10.
- Panaskar, D. B., and Pawar, R. S. (2011). Effect of textile mill effluent on growth of *Vigna unguiculata* and *Pisum sativum* seedlings. *Indian Journal of Science and Technology*, 4(3), 266-272.
- Pansamrit, K. and Ruangsomboon, S. (2010). Effect of pH on Biosorption of Basic Dye Malachite Green by Algae, The 8th International Symposium on Biocontrol and Biotechnology, 132-136.
- Papenfuss, G.F. (1955). Classification of the algae. In: *A century of progress in the natural sciences*, California Academy Science, San Francisco, 115-224.
- Parihar, A. and Malaviya, P. (2013). Textile wastewater treatment using sawdust as adsorbent. *Intl J Environ Sci*, 2, 110-113.
- Parshetti, G.K., Parshetti, S.G., Telke, A.A., Kalyani, D.C., Doonq, R.A. and Govindwar, S.P (2011). Biodegradation of crystal violet by *Agrobacterium radiobacter*. *J. Environ Sci*, 23(18), 1384-93.
- Patel, S. (2011). Potential of fruit and vegetable wastes as novel biosorbents, *Rev Environmental Science and Biotechnology*, 11:365–380.
- Paul, S.A., Chavan, S.K. and Khambe, S.D. (2012). Studies on characterization of textile industrial waste water in Solapur city, *Int. J. Chem. Sci.*: 10(2), 635-642.
- Pearse, A. G. E. (1968). *Histochemistry: Theoretical and Applied*. 3rd Edn., J and A Churchill Ltd., London, pp: 13-102
- Phugare, S., Patil, P., Govindwar, S. and Jadhav, J. (2010). Exploitation of yeast biomass generated as a waste product of distillery industry for remediation of textile industry effluent, *International Bioremediation and Biodegradation*, 64, 716-726.

- Prasad, A. and Rao, K.V.B. (2011), Physicochemical analysis of textile effluent and decolourization of textile azo dye by *Bacillus endophyticus* strain vitabr13, *The IIOAB Journal*, 22(), 55-62.
- Prola, L.D.T ., Acayanka, E ., Lima, E.C ., Umpierres, C.S ., Vaggetti, J.C.P ., Santos, W.O ., Laminsi, S ., and Djifon, P.T. (2013), Comparison of *Jatropha curcas* shells in natural form and treated by non-thermal plasma as biosorbents for removal of Reactive Red 120 textile dye from aqueous solution. *Ind Crop Prod*, 46:328–340.
- Rajan, M.R., Prabhakaran, M. and Noel, S.D. (2016). Field Level Study on The Utilization of Dyeing Industry Effluent Residue on Growth of Brinjal (*Solanum melongena*), *Indian Journal of Research*, 5(3), 50-52.
- Ramavandi, B., Ebrahimi, A., Hashemi, S. E., Akbarzadeh, S. and Asgari, G. (2015). The Potential of *Sargassum oligocystum* Harvested From Persian Gulf for the Adsorption of Copper Ions from Aqueous Solutions. *Avicenna Journal of Environmental Health Engineering*, 2(1), e3155.
- Ramesh, M., Sankaran, M., Veera Gowtham, V. and Poopal, P.K. (2014), Hematological, biochemical and enzymological responses in an Indian major carp *Labeo rohita* induced by sublethal concentration of waterborne selenite exposure, *Chemico-biological interactions* , 207, 67-73.
- Rani, B., Kumar, V., Singh, J., Bisht, S., Teotia, P., Sharma, S. and Kela, R. (2014). Bioremediation of dyes by fungi isolated from contaminated dye effluent sites for bio-usability. *Brazilian Journal of Microbiology*, 45, (3), 1055-1063.
- Rao, C.V., Giri, A.S., Goud, V.V. and Golder, A.K. (2016). Studies on pH Dependent colour variation and decomposition mechanism of Brilliant Green Dye in Fenton Reaction, *International Journal of Industrial Chemistry*, 7, 71- 80.
- Rashidi, H. R., Sulaiman, N. M. N., Hashim, N. A., Hassan, C. R. C., & Ramli, M. R. (2015). Synthetic reactive dye wastewater treatment by using nano-membrane filtration. *Desalination and Water Treatment*, 55(1), 86-95.

- Reddy, D.H.K. and Lee, S.M. (2013), Application of magnetic chitosan composites for the removal of toxic metal and dyes from aqueous solutions, *Adv Colloid Interface*, 201–202:68–93.
- Rehman, M.S.U., Munir, M., Ashfaq, M., Rashid, N., Nazar, M.F., Danish, M. and Han, J. (2013). Adsorption of Brilliant Green dye from aqueous solution onto red clay, *Chemical Engineering Journal.*, 228, 54–62.
- Rehman, R., Mahmud, T. and Irum, M. (2015). Brilliant green dye elimination from water using *Psidium guajava* leaves and *Solanum tuberosum* peels as adsorbents in environmentally benign way, *Journal of Chemistry*, 2015, 8.
- Remyla, S.R., Ramesh, M., Sajwan, K.s. and Kumar, K.S. (2008). Influence of zinc on cadmium induced haematological and biochemical responses in a freshwater teleost fish *Catla catla*, *Fish Physiology and Biochemistry* 34(2), 169-74.
- Roberts, R. J. (1978). *The Pathophysiology and systemic pathology of Teleosts and Laboratory methods*. 1st Edn., Bailliere Tindall, London, UK., pp: 235-246.
- Robinson, T., McMullan, G., Marchant, R., & Nigam, P. (2001). Remediation of dyes in textile effluent: A critical review on current treatment technologies with a proposed alternative. *Bioresource Technology*, 77(3), 247-255. DOI: 10.1016/S0960 8524(00)00080-8.
- Rohit, K.C. and Ponmurugan, P. (2013), Physicochemical analysis of textile, automobile and pharmaceutical industrial effluents, *International Journal of Latest Research in Science and Technology*, 2(2), 115-117.
- Roopadevi , H. and Somashekar R.K. (2012), Assessment of the toxicity of waste water from a textile industry to *Cyprinus carpio*, *J Environ Biol*, 33(2):167-71.
- Rosales, E., Anasie, D., Pazos, M., Lazar, I. and Sanromán, M. A. (2018). Kaolinite adsorption-regeneration system for dyestuff treatment by Fenton based processes. *Science of The Total Environment*, 622, 556-562.

- Rosales, E., Anasie, D., Pazos, M., Lazar, I., and Sanromán, M. A. (2018). Kaolinite adsorption-regeneration system for dyestuff treatment by Fenton based processes. *Science of The Total Environment*, 622, 556-562.
- Rubin, E., Rodriguez, P., Herrero, R and Cremades, J., Barbara, I and Manuel E Sastre de Vicente (2005). Removal of Methylene Blue from aqueous solutions using as biosorbent *Sargassum muticum*: an invasive macroalga in Europe. *Journal of Chemical Technology and Biotechnology*, 80:291–298. DOI: 10.1002/jctb.1192.
- Sahli (1962). Determination of haemoglobin by acid haematin method. In: *Practical haematology* (Ed: Dacie J.V. and Lewis S.M.) 5th Edn. Churuchill, London.
- Sahoo P.K., Mukherjee S.C., Jain A.K. and Mukherjee A. (2003). Histopathological and electron microscopic studies of gills and opisthonephros of rohu, *Labeo rohita*, to acute and subchronic aflatoxin B1 toxicity. *Asian Fisheries Science*, 16, 257-268.
- Saravanan, M., Kumar, P. and Ramesh, M. (2011). Haematological and biochemical responses of freshwater teleost fish *Cyprinus carpio* (Actinopterygii: Cypriniformes) during acute and chronic sublethal exposure to lindane, *Pestic. Biochem. Physiol*, 100, 206-211.
- Saravanan, M., Suganya, R., Ramesh, M. and Nagamony, P. (2015). Iron oxide nanoparticles induced alterations in haematological, biochemical and ionoregulatory responses of an Indian major carp *Labeo rohita*, *J Nanopart Res*, 17:274.
- Sartape, A.S., Mandhare, A.M., Jadhav, V., Raut, P.D., Anuse, M.A. and Kolekar, S.S. (2013). Removal of Malachite green dye from aqueous solution with adsorption technique using *Limonia acidisimma* (wood apple) shell as low cost adsorbent. *Arabian Journal of Chemistry*. <http://dx.doi.org/10.1016/j.arabjc.2013.12.019>.

- Sasikala, T. and Poongodi, N. (2013), Impact of Dye Effluent on Seed Germination of Blackgram (*Vigna mungo*.L.hepper), Indian Journal of Applied Research, 3(8), 47. ISSN - 2249- 555X.
- Sawyer, C.N. and P.L. McCarty. (1978). Chemistry for Environmental Engineering. McGraw-Hill Book Company, New York.
- Sayed, A.H. and Moneeb, R.H. (2015). Hematological and biochemical characters of monosex tilapia (*Oreochromis niloticus*, Linnaeus, 1758) cultivated using methyltestosterone, The Journal of Basic & Applied Zoology, 72, 36-42.
- Selvaraj, D., Leena, R. and Kamal. D.C. 2015. Toxicological and histopathological impacts of textile dyeing industry effluent on a selected teleost fish *Poecilia reticulata*. Asian Journal of Pharmacology and Toxicology, 03(10): 26-30
- Sen, G., Behra, M.K. and Patel, P.N. (1992). Effect of zinc on haemato- biological parameters of *Channa punctatus*. J. Ecotoxicol. Environ. Monit, 2: 89-92.
- Shaffi, S.A. (2001), Influence of sub lethal HCH on differential distribution of dehydrogenases, glycogen, pyruvate and lactate in different brain region in *Labeo rohita*. Int. J. Fish, 48 (2), 151-157.
- Sharma, S and Adholeya, A. (2011). Detoxification and Accumulation of chromium from tannery effluent and spent chrome effluent by *Paecilomyces lilacinus* fungi. International Biodeterioration & Biodegradation 65: 309-317.
- Sharmila, S.K., Rebecca, J., Chandran, P.N. and Kripanand, P.N. (2015). Extraction of biofuel from seaweed and analyse its engine performance, International Journal of Pharmacy and Technology, 7(2), 8870-8875.
- Shedbalkar, U. and Jadhav, J.P. (2011). Detoxification of malachite green and textile industrial effluent by *Penicillium ochrochloron*, Biotechnology and Bioprocess Engineering, 16(1), 196-204.

- Shertate, R.S. and Thorat, P.R. (2016). Bioremedial potential of marine bacterium. *World journal of pharmaceutical research*, 5(5), 1130-1150.
- Singh, S.M., Varshneya, I. and Nagarkoti. (1998). Assessment of physocp-chemical parameters of effluents of three factories of Bareilly district and their possible effects on grazing animals and cereals, *J. Environ. Bio*, 19(3), 271.
- Sivakumar, P., Kanagappan, M. and Sam Manohar Das, S. (2015). Chronic Histopathological Changes in Selected Tissues of *Danio Rerio* Exposed to Effluents from Leather Tanning Industry, 4(4), 1009-1009.
- Siyanbola, T O., Ajanaku, K O., James, O O., Olugbuyiro, J A O. and JAdekoya, J O. (2011). Physico- Chemical Characteristics of Industrial Effluents In Lagos State, Nigeria, *G. J. P & A Sc and Tech*. 01: 49-54.
- Sobecka, E. (2001). Changes in the iron level in the organs and tissues of wels cat fish, *silurus Glanis L.* Caused by nickel. *Acta Ichthyol . Piscat*, 31 (2), 127-143.
- Southamani, C., Shanti, G., & Deivasigamani, M. (2015). Hematological response in three Indian major carps in relation to supplementary feeding. *International journal of fisheries and aquatic studies*, 3(2), 287-294.
- Souza, R. P., Freitas, T. K., Domingues, F. S., Pezoti, O., Ambrosio, E., Ferrari-Lima, A. M., & Garcia, J. C. (2016). Photocatalytic activity of TiO₂, ZnO and Nb₂O₅ applied to degradation of textile wastewater. *Journal of Photochemistry and Photobiology A: Chemistry*, 329, 9-17.
- Srivastava, S., Sinha,R. and Roy, D (2004). Toxicological effects of malachite green. *Aquat Toxicol*. 66: 319-329.
- STATSOFT, INC, Statistica for Windows [Computer program manual]. Tulsa, OK: StatSoft, Inc., 2300 East 14th Street, Tulsa, OK 74104 (1999).
- Stoskopf, M.K. (1993). Clinial pathology. In: Stoskopf M.K. (Ed.). Philadelphia, Pa: WB Saunder. Fish Med. pp. 113-131

- Sultana, M., Islam, M., Sahaa, R. and Al-Mansur, M. (2009). Impact of the effluents of textile dyeing industries on the surface water quality inside D.N.D (Dhaka- Narayanganj-Demra) Embankment, Narayanganj. *Bangladesh Journal of Scientific and Industrial Research*, 44, 65-80.
- Taghipour, T., Karimipour, G., Ghaedi, M., and Asfaram, A. (2018). Mild synthesis of a Zn (II) metal organic polymer and its hybrid with activated carbon: Application as antibacterial agent and in water treatment by using sonochemistry: Optimization, kinetic and isotherm study. *Ultrasonics sonochemistry*, 41, 389-396.
- Tahir, H., Sultan, M. and Jahanzeb, Q. (2008). Removal of basic dye methylene blue by using bioabsorbents *Ulva lactuca* and *Sargassum*, *African journal of Biotechnology*, 7(15), 2649-2655.
- Tan, P., Wong, C., Ong, S. and Hii, S. (2009). Equilibrium and Kinetic studies for Basic yellow 11 removal by *Sargassum binderi*, *Journal of Applied Sciences*, 9, 3005-3012.
- Tangaromsuk, J., Pokethitiyook, P., Kruatrachue, M. and Upatham, E.S. (2002), Cadmium biosorption by *Sphingomonas paucimobilis* biomass. *Biores Technol*, 85, 103–5.
- Tekoglu, O. and Ozdemir, C. (2010). Wastewater of textile industry and its treatment processes, *Agricultural and Economic Research Review*, 23, 245-252.
- Tortman, S.R, Tortman, E.R. (1964), *The bleaching, dyeing chemical technology of Textile fibers*. Charless Griffen and Comp. Ltd., London.
- Tripathi, S., Sahu, D. B., Kumar R. and Kumar A. (2003). Effect of acute exposure of sodium arsenite (Na₃ AsO₃) on some haematological parameters of *Clarias batrachus* (common Indian cat fish) in vivo, *Indian J Environ Health*, 45(3), 183-8.

- Tripathy, S. K. and Patel, S. (2014). Abnormal mitosis in root meristem cells of *Allium cepa* L. induced by a fabric dye reactive turquoise blue (Procion MX). African Journal of Biotechnology, 13(38).
- Turner, R.G. and Marshall, C. (1972). Accumulation of zinc by subcellular fractions of roots of *Agrostis tannis Sibth* in relation of zinc tolerance, New Phytologist, 71, 671- 676.
- Varma, L. and Sharma, J. (2011), Analysis of physical and chemical parameters of textile waste water, Journal of International Academy of Physical Sciences, 15(2), 269-276.
- Varsha, G., Sudesh and Seeema, S. (2013). Physico-chemical analysis of textile effluents of dye and printing clusters of bagru region, Jaipur, India : Journal of Environmental Research And Development, 8(1), July-September 2013.
- Verma, S.R. and Dalela, R.C. (1975). Studies on the pollution of the Kalinadi by industrial wastes near Mansurpur Part-II: Biological Index of Pollution and Biological Characteristics of the River, Acta Hydrochim Hydrobiol, 3(3), 259-274.
- Verschuere K. (2009), Handbook of Environmental Data on Organic Chemicals, Four Volume Set, 5th edition, 4486 pages.
- Vieira, R. and Volesky, B. (2000). Biosorption: a solution to pollution?, International Microbiology, 3, 17-24.
- Vigneshpriya, D. and Shanthi, E. (2016), Biosorption of textile effluent using sugarcane bagasse, International Journal of Current Research, 8, 31324-31331.
- Vigneshpriya, D., Krishnaveni, N., and Renganathan, S. (2017). Marine brown macroalga *Sargassum wightii* as a novel biosorbent for removal of brilliant green dye from aqueous solution: kinetics, equilibrium isotherm modeling and phytotoxicity of treated and untreated dye. Desalination and water treatment, 78, 300-312.

- Vigneshpriya, D., Krishnaveni, N., Bhargavi, G., Sri Sakthi Priyadarshini, R. and Renganathan, S. (2017), Effect of textile effluent on growth and germination of cow pea *Vigna unguiculata* L. Indian Journal of Environmental Protection, 37(2):163-168.
- Vijayaraghavan K, Yun YS (2008). Biosorption of C.I. reactive black 5 from aqueous solution using acid-treated biomass of brown seaweed *Laminaria* sp. Dyes Pigments 76: 726-732.
- Vijayaraghavan, K., Mao, J. and Yun, Y. S. (2008). Biosorption of methylene blue from aqueous solution using free and polysulfone-immobilized *Corynebacterium glutamicum*: batch and column studies. Bioresource technology, 99(8), 2864-2871.
- Wamba, A. G., Lima, E. C., Ndi, S. K., Thue, P. S., Kayem, J. G., Rodembusch, F. S. and de Alencar, W. S. (2017). Synthesis of grafted natural pozzolan with 3-aminopropyltriethoxysilane: preparation, characterization, and application for removal of Brilliant Green 1 and Reactive Black 5 from aqueous solutions. Environmental Science and Pollution Research, 24(27), 21807-21820.
- Wang, Y.J., Jia, D.A, Sun R.J., Zhu, H.W. and Zhou, D.M. (2008) Adsorption and cosorption of tetracycline and copper(II) on montmorillonite as affected by solution pH. Environ Sci Technol 42(9):3254–3259.
- Wanyonyi, W.C., Onyari, J.M. and Shiundu, P.M. (2014). Adsorption of Congo Red Dye from Aqueous Solutions Using Roots of *Eichhornia crassipes*: Kinetic and Equilibrium Studies. Energy Procedia, 50, 862 – 869.
- Wepener, V. and Du, H. P. (1992). The effect of hexavalent chromium at different pH values on the haematology of *Tilapia sparrmanii* (Cichlidae). Comparative biochemistry and physiology. C, Comparative pharmacology and toxicology, 101(2), 375-381.
- Wetzel, R.G. (1972). Primary production. In: Mitchel RA. Water pollution microbiology, Wiley Inter science, New York. 230.

- Wins, J.A. and Manavalan, M. (2010). Effect of textile mill effluent on growth and germination of black gram - *Vigna mungo* (L.) Hepper, International Journal of Pharma and Bio Sciences, 1(1).
- Wu, Chung- Hsin. (2007). Adsorption of reactive dye onto carbon nanotubes: equilibrium, kinetics and thermodynamics, Journal of Hazardous Materials, 144 (1), 93-100.