

Bibliography

- Appanah, T. P., Oogarah-Pratap, B. and Ruggoo, A. (2009). Awareness and consumption of iron among Mauritian female factory workers., *Nutrition and Food Science*,39(2): pp. 168–175.
- Venkaiah, K., Brahmam, G.N. V. and Vijayaraghavan, K. (2011). Application of factor analysis to identify dietary patterns and use of factor scores to study their relationship with nutritional status of adult rural populations. *J Health Popul Nutr.*; 29:327–38.
- Kossioni, A and Bellou, O. (2011) Eating habits in older people in Greece: The role of age, dental status and chewing difficulties, *Archives of Gerontology and Geriatrics*, Vol. 52, issue 2;pp. 197–201.
- National Center for Education Statistics. (2008). Percentage of high school dropouts among persons 16 through 24 years old (status dropout rate), by income level, and percentage distribution of status dropouts, by labor force status and educational attainment: 1970 through 2007. Retrieved from http://nces.ed.gov/programs/digest/d08/tables/dt08_110.asp
- Intawongse, M and Dean, J.R. (2008) Use of the physiologically-based extraction test to assess the oral bioaccessibility of metals in vegetable plants grown in contaminated soil. *Environ Pollut*; 152:60–72
- Singh, A.P., Luthria, D., Wilson, T., Vorsa, N., Singh, V., Banuelos, G.S., Pasakdee, S., (2009). Polyphenols content and antioxidant capacity of eggplant pulp. *Food Chem.*, 114: 955 961.
- Sandstrom, B. and Cederblad, A. (1987). Effect of ascorbic acid on the absorption of zinc and calcium in man. *International Journal of Vitamin Nutrition Research* 57,87-90.
- Harriss-White, B. and Hoffenberg, R (1994) eds. *Food: Multidisciplinary Perspectives*, Blackwell, Oxford

- Afridi, H.I., Kazi, T.G., Kazi, N., Kandhro, G.A., Baig, J.A., Shah, A.Q., Kolachi, N.F., Wadhwa, S.K., Khan, S. and Shah, F. (2010). Potassium, calcium, magnesium, and sodium levels in biological samples of Pakistani myocardial infarction patients at different stages as related to controls. *Clin Lab.*, 56:427–439
- Agarwal, S., Millett, C.J., Dhillon, P. K., Subramanian, S. V and Ebrahim., S.(2014) Type of vegetarian diet, obesity and diabetes in adult Indian population., *Nutrition Journal* ;2014, 13:89
- Al Hasan S, M., Hassan, M., Saha, S., Islam, M., Billah, M. and Islam, S. (2016) Dietary phytate intake inhibits the bioavailability of iron and calcium in the diets of pregnant women in rural Bangladesh: a cross-sectional study. *BMC Nutrition.*, 21; 2(1):1.
- Association of Official Analytical Chemists (2005). (AOAC). Official methods of analysis of the association of official analytical chemists, 18th Ed.
- Arnold, F., Parasuraman, S., Arokiasamy, P. and Kotha, M. (2009). Nutrition in India, National Family Health Survey. (In NFHS 3) International Institute for Population Sciences. Calverton, Maryland, USA: ICF Macro;
- Badii, A., Nekouei, N., Fazilati, M., Shahedi, M. and Badiei, S. (2012). Effect of consuming zinc-fortified bread on serum zinc and iron status of zinc-deficient women: A double blind randomized clinical trial. *International Journal of Preventive Medicine* 3: S 124–130.
- Baech, S.B., Hansen, M., Bukhave, K., Jensen, M., Sorensen, S.S., Kristensen, L., Purslow, P.P., Skibsted, L.H., and Sandström B. (2003). Non haeme-iron absorption from a phytate-rich meal is increased by the addition of small amounts of pork meat. *Am J Clin Nutr.* 77(1):173-9.
- Bains, K and Shruti (2007). Analysis of various vegetable preparations for calcium, iron and zinc intake of Punjabi urban and rural families., *Natural Product Radiance*: 6(2): pp.106-110
- Baker, C. J. L. (1952). The determination of oxalates in fresh plant material. *Analyst*, 77, 340-344.

- Balarajan, Y., Ramakrishnan, U., Ozaltin, E., Shankar, A. H. and Subramanian, S. V. (2011). Anaemia in low-income and middle-income countries. *Lancet* 378: 2123–35.
- Baynes, R. D. and Bothwell, T.H. (1990). Iron deficiency. *Ann. Rev. Nutr.* 10:133-148.
- Benito, P and Miller, D. (1998) Iron absorption and bioavailability: an updated review. *Nutr Res*; 18: 581-603.
- Biradar, S. S., Biradar, S.P., Alalagi, A.C., Wantamutte, A.S. and Malur, P.R. (2012) Prevalence of Anaemia among Adolescent Girls: A One Year Cross-Sectional Study. *Journal of Clinical And Diagnostic Research*; 6:372-377.
- Black, R.E., Allen, L. H., Bhutta, Z. A., Caulfield, .L.E., Onis, M.D., Ezzati, M., Mathers, C. and Rivera, J. (2008). Maternal and child under nutrition: global and regional exposures and health consequences. *The Lancet* 34:404.
- Blenford, D. (1995). Bioavailability is key to nutrient effectiveness. *Food Ingrid Process Int.*17:28-30
- Brown, K. H., Peerson, J. M., Baker, S. K. and Hess, S.Y. (2009). Preventive zinc supplementation among infants, preschoolers, and older prepubertal children. *Food and Nutrition Bulletin*: 30 (1 Supplement): S12–S40.
- Census of India (2011). Census data shows numbers rising more in urban areas. The Indian Express. New Delhi: 2011. <http://archive.indianexpress.com/news/census-data-shows-numbers-rising-more-in-urban-areas/818130/2> [Accessed 28 May 2015]
- Centers for Disease Control and Prevention (2013). About BMI for children and teens. Atlanta, GA: Centers for Disease Control and Prevention.
- Chandyo, R.K., Strand, T.A., Mathisen, M., Ulak, M., Adhikari, R.K. and Bolann, B.J. (2009) Zinc deficiency is common among healthy women of reproductive age in Bhaktapur, Nepal. *J Nutr.*, 139:594–597.
- Chavan, J. K. and Kadam, S.S. (1989). Nutritional improvement of cereals by fermentation. *Critical Reviews in Food Science and Nutrition* 28:349–400.
- Chellappa, A. R. and Karunanidhi. S. (2012). Supplementation with Iron and Zinc Selectively Improves Cognitive and Behavioral Functions in Female Adolescents. *International Journal of Chemical Engineering and*

- Applications*, 3(4), 274–281. doi:10.7763/IJCEA.2012.V3.199
- Cousins, R.J. (1985). Absorption, transport, and hepatic metabolism of copper and zinc: Special reference to metallothionein and ceruloplasmin. *Physiol Rev.* 65:238–309. [PubMed]
- Cozzolino, S.C., (2007) Estudos Avançados *Print version* ISSN 0103-4014 *On-line version* ISSN 1806-9592, Estud. av. vol.21 no.60 Sao ., <http://dx.doi.org/10.1590/S0103-40142007000200009>
- Daniel, C. R., Prabhakaran, D., Kapur, K., Graubard, B. I., Devasenapathy, N., Ramakrishnan, L., George, P.S., Shetty, H., Ferrucci, L.M., Susan ,Y., Chatterjee, N., Reddy, K.S., Rastogi, T., Gupta, P.C., Mathew, A and Sinha, R. (2011). A cross-sectional investigation of regional patterns of diet and cardio-metabolic risk in India. *Nutrition Journal*, 10(1), 12. <http://doi.org/10.1186/1475-2891-10-12>
- Deshpande, N.S., Karva, D., Agarkhedkar, S. and Deshpande, S. (2013). Prevalence of anemia in adolescent girls and its co-relation with demographic factors. *Int J Med Public Health*;3:235-9.
- Dhakar, R.C., Maurya, S.D., Pooniya, B.K., Bairwa, N., Gupta, M. and Sanwamal, S. (2011). Moringa : The herbal gold to combat malnutrition. *Chron Young Sci* [serial online], 2:119-25. <http://www.cyonline.org/text.asp?2011/2/3/119/90887>
- Diaz, M., Rosado, J.L., Allen, L.H., Abrams, S. and Garcia, O.P: (2003). The efficacy of a local ascorbic acid-rich food in improving iron absorption from Mexican diets: a field study using stable isotopes. *Am J Clin Nutr*, 78:436-440.
- District Level Household and Facility survey (2012) District Fact sheet., District Level Household and Facility survey -IV (DLHFS -IV)., International Institute of Population Sciences., Mumbai- 400088.
- Eaton, S.B., Eaton, S.B. and Konner, M.J. (1999). Paleolithic nutrition revisited: a twelve-year retrospective on its nature and implications" *European Journal of Clinical Nutrition* 51 (4): 207–16.
- Eleventh five year plan (2012). Report of the 11th 5 year plan., Planning Commission, Government of India., New Delhi-110002.
- Elizabeth, K. (2001). A novel growth assessment chart for adolescents. *Indian paediatrics* , 1061-1064.
- Eng, P. M., Kawachi, I., Fitzmaurice, G. and, Rimm, E. B. (2005) Effects of marital transitions on changes in dietary and other health behaviours in US male health professionals, *Journal of Epidemiology and Community Health*: 59 (1) ;pp. 56–62
- Engelmann, M.D., Davidsson, L., Sandstrom, B., Walczyk, T., Hurrell, R.F. and Michaelsen, K.F. (1988). The influence of meat on non haeme iron absorption in infants. *Pediatr Res* 43:768–73.

- Engle-Stone, R., Yeung, A., Welch, R. and Glahn, R. (2005). Meat and ascorbic acid can promote Fe availability from Fe-phytate but not from Fe-tannic acid complexes. *J Agric Food Chem.* 28; 53 (26):10276-84.
- Fabbri, A. D. T. and Crosby, G. A. (2015). A review of the impact of preparation and cooking on the nutritional quality of vegetables and legumes. *International Journal of Gastronomy and Food Science*, 3, 1–10. <http://doi.org/10.1016/j.ijgfs.2015.11.001>
- Fairweather-Tait, S. J. (1993). Bioavailability of nutrients. In: Macrae, R. Robinson, R.K. and Sadler, M.J. (editors.) *Encyclopedia of Food Science, Food Technology And Nutrition*. London: Academic Press. p. 384-8.
- FAO / WHO. (2004) Expert Consultation on Human Vitamin and Mineral Requirements, Vitamin and mineral requirements in human nutrition: Report of joint FAO/WHO expert consultation; 2nd ed. Bangkok, Thailand; p. 341.
- FAO/WHO. 2001. Joint report: Human vitamin and mineral requirements. Rome, Italy: Food and Nutrition Division, FAO/WHO.
- Fayeta, F., Flood, V., Peter, P. and Samman, S. (2011). Relative and biomarker-based validity of a food frequency questionnaire that measures the intakes of vitamin B12, folate, iron, and zinc in young women. *Nutrition Research*, 31: 14 -20.
- Fesharakinia, A.Z.A., Sharifzadeh, G.R. (2009). Prevalence of zinc deficiency in elementary school children of South Khorasan Province (East Iran). *Iran J Pediatr.*, 19: 249–254.
- Fidler, M.C., Davidsson, L., Zeder, C. and Hurrell, R. F. (2004). Erythorbic acid is a potent enhancer of nonhaeme-iron absorption. *Am J Clin Nutr.*;79(1):99-102.
- Food and Agriculture Organization (2010). Food Balance Sheets. (FAO). Available: <http://faostat.fao.org/site/345/default.aspx> [accessed June 2011].
- Food Security Bill (2011). The National Food Security Bill- Report., Government of India. New Delhi-110002
- Fordyce, E. J., Forbes, R. M., Robbins, K. R. and Erdman, J. W. (1987). Phytate x calcium / zinc molar ratios: are they predictive of zinc bioavailability; *J. Food Sci.*, 52: 440-444.
- Garcia-Casal, M.N. (2006): Carotenoids increase iron absorption from cereal-based food in the human. *Nutr Res*, 26:340-344.
- Geissler, C. and Singh, M. (2011). Iron, Meat and Health, *Nutrients*; 3(3): 283–316.
- Gibson, R.S., Bailey, K.B., Gibbs, M. and Ferguson, E.L. (2010); A review of phytate, iron, zinc, and calcium concentrations in plant-based complementary foods used in low-income countries and implications for bioavailability. *Food Nutr Bull.*, 31:S134–S146

- Gopalan, C., Sastri, R.B. V. and Balasubramaniam, S.C. (2010) Nutritive Value of Indian Foods, Indian Council of Medical Research; National Institute of Nutrition Hyderabad; Pp11-18.
- Green, R., Milner, J., Joy, E. and Dangour, A. (2015). Dietary patterns in India: Systematic Literature Review; *Br. J. Nutr.* ; 116(1): 142–148.
- Gupta, A. and Kumar, P. (2015). Regional Patterns of Food intake and Nutrients in Rural India: Evidence from NSS 66th round., International Research Journal of Social Sciences, ISSN 2319–3565 Vol. 4(7), 1-11
- Gupta, S, Lakshmi, A.J. and Prakash, J. (2006). *In vitro* bioavailability of calcium and iron from selected green leafy vegetables. *Journal of the Science of Food and Agriculture*, 86(13), 2147–2152
- Gupta, S.K., Agarwal, S.S., Kaushal, R., Jain, A., Gupta, V.K. and Khare, N. (2014). Prevalence of anemia among rural population living in and around of rural health and training center, Ratua Village of Madhya Pradesh. *Muller J Med Sci Res*, 5(1): 15-18
- Hambidge, M. (2000). Human zinc deficiency. *J Nutr* 130(5S Suppl):1344S-1349S.
- Heaney, R. P. (2001). Factors influencing the measurement of bioavailability, taking calcium as a model. *Journal of Nutrition* 131(suppl):1344S-1348S.
- Hemalatha, S., Platel, K. and Srinivasan, K. (2006). Influence of heat processing on the bioaccessibility of zinc from cereals and pulses consumed in India. *J Trace Elements Med Biol*; 21(1):1-7.
- Hemalatha, S., Platel, K. and Srinivasan, K. (2007) Zinc and iron contents and their bioaccessibility in cereals and pulses consumed in India. *Food Chem* 102(4):1328– 1336.
- Hemalatha, S., Platel, K. and Srinivasan, K. (2005). Influence of food acidulants on bioaccessibility of zinc and iron from selected food grains. *Mol. Nutr. Food Res.*, 49: 950-956
- Holst, B. and Williamson, G. (2008) Nutrients and phytochemicals: from bioavailability to bioefficacy beyond antioxidants. *Current Opinion in Biotechnology* 19:73-82. doi: 10.1016/j.copbio.2008.03.003
- Hotz, C. and Brown, K. H. (2004). Assessment of the risk of zinc deficiency in populations and options for its control; International Zinc Nutrition Consultative Group (IZiNCG): 25: 1-204
- Hunt, J.R. and Vanderpool, R.A. (2001). Apparent copper absorption from a vegetarian diet. *Am J Clin Nutr.* ; 2001;74:803–7. 10
- Hurrell, R. and Egli, I. (2010). Iron bioavailability and dietary reference values. *American Journal of Clinical Nutrition*; 91(5):1461S-1467S. doi: 10.3945/ajcn.2010.28674F
- International Nutritional Anaemia Consultative Group (1984) Iron Deficiency In Women. Washington, DC: Nutrition Foundation.

- IZiNCG (International Zinc Nutrition Consultative Group). 2004. Assessment of the risk of zinc deficiency in populations and options for its control (Hotz C and Brown KH, eds.). *Food and Nutrition Bulletin* 25:S91–S202.
- Jackson, M.J. (1997). The assessment of bioavailability of micronutrients: introduction,; *Eur. J. Clin. Nutr.*, 51: S1-S29.
- Jelliffe, D.B. (1966). Assessment of nutritional status of community, WHO monograph series 53.
- Jideani, V. A, and Jideani, I. A. (2006). Laboratory Manual of Food Bacteriology. Amana Printing and Advertising Ltd , Kaduna, Nigeria. Pp. 96-108
- Joshi, M. and Gumashta, R. (2013). Weekly iron folate supplementation in adolescent girls--an effective nutritional measure for the management of iron deficiency anaemia. *Global journal of health science*, 5(3), 188-94. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/23618489>
- Joshi, P. and Mehta, D. (2010)., Effect of dehydration on the nutritive value of drumstick leaves., *Journal of Metabolomics and Systems Biology.*, 1: pp.5—9.
- Kalasuremath, S., Kurpad, A., V. and Thankachan, P. (2013). Effect of iron status on iron absorption in different habitual meals in young south Indian women, *Indian Journal of Medical Research.*, 137 p: 324—330
- Kanani, S. and Poojara, R.M (2000) Supplementation with IFA enhances growth in adolescent Indian girls. *J Nutr.* ; 130: 452S-455S.
- Kaur, M., Bassi, R. and Sharma, S. (2011). Impact of nutrition education in reducing iron deficiency anemia in adolescent girls., *Indian Journal of Fundamental and Applied Life Sciences*; 1 (4): pp.222-228 .
- Kaur, K. (2014). Anaemia ‘a silent killer’ among women in India: Present scenario., *European journal of Zoological Research*, 3(1): 32-36.
- Kawade, R. (2012). Zinc status and its association with the health of adolescents: a review of studies in India. *Global Health Action* 5:7353. (doi:10.3402/gha.v5i0.7353)
- King, J.C. (2011). Zinc: an essential but elusive nutrient. *American Journal of Clinical Nutrition.*, 94(2):679S– 684S.
- Kodkany, B.S., Bellad, R.M., Mahantshetti, N.S., Westcott, J.E., Krebs, N.F., Kemp, J.F and Hambidge, K.M. (2013). Bio fortification of pearl millet with iron and zinc in a randomized controlled trial increases absorption of these minerals above physiologic requirements in young children. *Journal of Nutrition.* (doi:10.3945/jn.113. 176677)
- Kolb, L.E. and Beard, J.L. (2007). Iron treatment normalizes cognitive functioning in young women. *Am J Clin Nutr.*;85: 778–787.
- Kollipara, U.K. and Brittin, H.C. (1996). Increased iron content of some Indian foods due to cookware. *J Am Diet Assoc.* 96(5):508-10.
- Kotecha, P. V., Nirumpam, S. and Karkar, P. D. (2009). Adolescent girl’s

- anaemia control programme, Gujarat, India., *Indian J Med Res*; 130: 584-589
- Kowsalya, S. and Shimaray, A.S. (2008). Impact of supplementation of lotus stem (*Nelumbium Nelumbo*) on the iron nutriture of adolescent girls (13-18 years) in Manipur. *The Indian Journal of Nutrition and Dietetics*,; 45: 47-53.
- Krebs, N.F. (2000). Overview of zinc absorption and excretion in the human gastrointestinal tract., *J Nutr*;130(5S Suppl):1374S-7S.
- Kristin, L., Welle, N., Tone, M., Steinar, H., and Anne, G. *J Prim Health Care*. (2011); 29(1):39–44. doi: 10.3109/02813432.2010.545209
- Kuligowski, J. and Halperin, K.M. (1992). Stainless steel cookware as a significant source of nickel, chromium, and iron. *Arch Environ Contam Toxicol*. 23(2):211-5.
- Kulkarni, M. V., Durge, P. M. and Kasturwar, N. B. (2012). Prevalence of anaemia among adolescent girls in an urban slum., *NJCM*: 3(1), 108-11
- Kumar, V. and Rajagopalan, M. (2008). Trial using multiple micronutrient food supplement and its effect on cognition, *Indian Journal of Pediatrics*., 75(7): 671-678
- Kumar, V. M. and Rajagopalan, S. (2006) Impact of a multiple- micronutrient food supplement on the nutritional status of schoolchildren. *Food Nutr Bull* ; 27: 203:10.
- Larmond, E. (1977) Laboratory methods for sensory evaluation of foods. Publication No. 1637, Department of Agriculture, Ottawa, pp 17–22
- Lestienne, I., Icard-Verniere, C., Mouquet, C., Picq. C. and Treche, S. (2005). Effect of soaking whole cereal and legume seeds on iron, zinc and phytate contents. *Food Chem*; 89: 421-425
- Lombardi-Boccia, G., De Santis, N., Di Lullo, G. and Carnovale, E. (1995). Impact of processing on Fe dialysability from bean (*Phaseolus vulgaris* L.). *Food Chem* 53, 191–195
- Luo, Y. and Xie, W. (2013) Effect of soaking and sprouting on iron and zinc availability in green and white faba bean (*Vicia faba* L.); *J Food Sci Technol* (December 2014) 51(12):3970–3976
- Luten, J., Crews, H., Flynn, A., Vandael, P., Kastenmeyer, P., Hurrell, R., Deelstra, H., Shen, L.H., Tait, S.F., Hickson, K., Farre, R., Schlemmer, W. and Frohlich, W., (1996), Inter-laboratory trial on the determination of the *in vitro* dialyzability from food. *J.Sci. Food Agri.*, 72: 415-424.
- Patted, M., Pushpa, B., and Hemalatha, S. (2013). Influence of acidulants on bioaccessibility of zinc from green leafy vegetable products. *Karnataka*

- Journal of Agricultural Sciences*; 26(2):128–132.
- Mala, V., Kumar, D., Dwivedi, S and Dabral S.B., (2007). Psychological behavioural pattern of unmarried adolescent girls in urban area of Allahabad, Uttar Pradesh. *Indian J Community Med*; 32:7-9.
- Malleshi, N, G. and Desikachar, H.S.R.(1985). Milling, popping and malting characteristics of some minor millets. *J Food Sci Technol*; 22: 400–403.
- Mathangi, S.K., Hemalatha, G., Amutha, S. and Arulmozhichelvan, K. (2013). Food consumption patterns and the nutritional profile of the urban south Indian Dwellers., *Asian Journal Of Dairy and Food Research*, P: 235-240
- Micronutrient Initiative(2009). *Vitamin and Mineral Deficiency: a global progress report* .
- Misra, A., Khurana, L., Isharwal, S., & Bhardwaj, S. (2009). South Asian Diets and insulin resistance. *British Journal of Nutrition* , 101 (4), 465-473.
- Mohamed, A., Perera, P.J., Hafez, Y.S. (1986). New chromophore for phytic acid determination., *Cereal Chem*; 63(6): 475–478.
- Moradi, N. and Moradi, K. (2013). “Physiological and pharmaceutical effects of fenugreek (*Trigonella foenum - graecum* L.) as a multipurpose and valuable medicinal plant”, *Global journal of medicinal plant research*, 1(2):199-206
- Mubarak, A.E. (2005) Nutritional composition and antinutritional factors of mung bean seeds (*Phaseolus aureus*) as affected by some home traditional processes. *Food Chem*; 89:489–495
- Nair, K. M and Iyengar (2009)., Iron content, bioavailability & factors affecting iron status of Indians., *Indian J Med Res.*, 130,pp: 634-645.
- Nair, K. M., and Brahmam, G. N. V., Radhika, M.S., Choudhury, D.R., Punjal, R., Nagalla, B., Zhensheng, C., Keli , H.M. and Abrams, S. A. (2013)., Inclusion of Guava Enhances Non-Haeme Iron Bioavailability but Not Fractional Zinc Absorption from a Rice-Based Meals in adolescents., *The Journal of Nutrition, Nutrient Requirements and Optimal Nutrition.*,pp:852-858
- Nair, K.M. and Choudhury, D.R. (2013)., Zinc nutrition in health and diseases., *Journal of SAT Agricultural Research* 11.
- Nambiar, V. S., Guin, P., Parnami, S. and Daniel, M. (2010). Impact of antioxidants from drumstick leaves on the lipid profile of hyperlipidemics. *Health (San Francisco)*, 4(1), 165–172.
- Nambiar, V.S. and Parnami, S. (2008) Standardization and organoleptic evaluations of freshly blanched drumstick leaves (*Moringa oliefera*) incorporated recipes for young girls. *Trees for Life Journal*; 3 (3):1-7.
- National Council for Applied Economic Research. (2014) , An Analysis of Changing Food Consumption Pattern in India- a paper ., National Council

- of Applied Economic Research., Parisila Bhawan, 11, I.P Estate, New Delhi
- National Family Health Survey –IV (2016), International Institute for Population Sciences, Mumbai - 400 088 Maharashtra, India.
- National Nutrition Monitoring Bureau (2006). Diet and nutritional status of population and prevalence of hypertension among adults in rural areas. NNMB Technical Report No. 24. Hyderabad, India: National Institute of Nutrition, Indian Council of Medical Research. (Available from: <http://www.nnmbindia.org/NNMBReport06Nov20.pdf>)
- National Nutrition Monitoring Bureau (2012). Diet and Nutritional Status of Rural Population, Prevalence Of Hypertension & Diabetes Among Adults And Infant & Young Child Feeding Practices , (NNMB) -Report Of Third Repeat Survey National Institute Of Nutrition, Indian Council Of Medical Research, Hyderabad - 500 007. India
- National Sample Survey Organization (2012), NSS Report No.558: Household Consumption of Various Goods and Services in India, 2011-12.
- Navas-Carretero, S., Pérez-Granados, A.M., Sarriá, B., Carbajal, A., Pedrosa, M.M., Roe, M.A., Fairweather-Tait, S.J. and Vaquero, M.P. (2008). Oily fish increases iron bioavailability of a phytate rich meal in young iron deficient women. *J Am Coll Nutr.* 27(1):96-101.
- Neelam, D.S., Karva, D., Agarkhedkar, S. and Deshpande, S. (2013). Prevalence of anemia in adolescent girls and its co-relation with demographic factors.; *International Journal of Medicine and Public Health*; 3(4) : 236-239
- Odunayo, C.A. and Singh, V. (2007) Effect of cooking on the profile of phenolics, tannins, phytate, amino acid, fatty acid, mineral nutrients of whole-grain and decorticated vegetable cowpea *Vigna unguiculata* (L.) Walp. *J Food Quali* 30:1101–1120
- Ogunniy, L.T., Ajao, A.O. and Oladejo, J.A. (2012). Food consumption pattern in Ogbomoso Metropolis of Oyo State Nigeria. *JASR*, 12 (1): 74-83.
- Parimalavalli, P. and Sangeetha, M. (2011). Anthropometric Measurements and Nutrient Adolescent Girls. *Anthropologist*, 13(2): 111-115.
- Patted, M., Bharati, P and Hemalatha, S. (2010). Influence of acidulants on bioaccessibility of zinc from green leafy vegetable products ., *Karnataka J. Agric. Sci.*,26 (1) :pp128-132
- Paul, I., Turner, R.E. and Ross, D. (2004). Nutrition. Jones and Barlett Publishers.
- Pavithra, B.S., Basavaraja, H, Kiresur, V.R., Mahajanshetty and Mageri,S.N. (2009), An economic analysis of food consumption pattern in Karnataka, *Karnataka J. Agric. Sci.*; 22 (4):840-845
- Pereira, E.J., Lucia M. J. Carvalho, U., Gisela, M.. Dellamora-Ortiz, Flavio, S. N. Cardoso, J. L. V., Carvalho, D.S. V., Sidinea, C. F., and Maurisrael ,M. R. (2015). Effects of cooking methods on the *Vigna* iron and zinc contents in

cowpea to combat nutritional deficiencies in Brazil

- Popkin, B.M., Adair, L.S. and Ng, S.W. (2012) Global nutrition transition and the pandemic of obesity in developing countries.; *Nutrition Reviews*; 70: 3–21.
- Prasad, R., Gupta, A., Parihar, R. and Gangwar, K. (2014). In vitro method for predicting the bioavailability of iron from Bathua Indian cookies, 9411.
- Priya, S., S. and Kowsalya, S. (2015). Formulation and evaluation and convenience food mixes from malted millets., *International of Scientific Research*; 4(11):11-13.
- Radhika, G., R. Sathya, A., Ganesan, R., Saroja, P., Vijayalakshmi, V., Sudha, and Mohan. , (NNMB) .(2011), Dietary profile of urban adult population in South India in the context of chronic disease epidemiology (CURES-68). *Public Health Nutrition* 14 (4): 591–598.
- Ragaee, S., Abdel-Aal, E.M. and Noaman, M. (2006). Antioxidant activity and nutrient composition of selected cereals for food use. *Food Chemistry*, 98(1): 32-38
- Raghuram, V., Manjula, A. and Jayaram, S. (2012). Prevalence of anaemia amongst women in the reproductive age group in a rural area in South India. *Int J Biol Med Res*, 3(2): 1482-1484.
- Raghuramulu, N., Nair, K. M. and Kalyansunderaram, S. (2003). A manual of laboratory techniques. National Institute of Nutrition, Jamai-Osmania, Hyderabad, India.
- Rao, B. S. N. and Prabhavathi, T. (1982), Tannin content of foods commonly consumed in India and its influence on ionisable iron. *J. Sci. Food Agric.*, 33: 89–96. doi: 10.1002/jsfa.2740330116
- Reaburn, J.A., Krondl, M. and Lau, D, (1979). Social determinants in food selection. *J Am Dietet Assoc.*, 74: 637-641
- Recommended Dietray Allownaces (2010). Indian council of Medical Research, National Institute of Nutrition, Hyderabad.
- Sahai, R. and Gupta, A. (2012). In vitro Bioavailability of iron and nutritional composition of the products prepared from the leaves of the weed, Indian Sorrel (*Oxalis Corniculata*). M.Sc., Thesis of S.H.I.A.T.S., Allahabad, U.P. India.
- Sandberg, A.S. (2002). Bioavailability of Minerals in Legumes. *British Journal of Nutrition*, 88: 281-285.
- Sathya, R. Amirthaveni, M. and Vijaylakshmi, P. (2002). Enhancing the bioavailability of iron from mung bean (*Vigna radiata* L.) through simple modifications in cooking. *Ind. J. Nutr. Diet.*, 39: 45-54.
- Shayeghi, M., Latunde-Dada, G.O., Oakhill, J.S., Laftah, A.H., Takeuchi, K., Halliday, N., Khan, Y., Warley, A., McCann, F.E., Hider, R.C., Frazer, D.M., Anderson, G.J., Vulpe, C.D., Simpson, R.J. and McKie, A.T. (2005). Identification of an intestinal haeme transporter. *Cell*; 122(5):789-801.

- Singh, G. and Kawatra, A. (2006). Development and nutritional evaluation of recipes prepared using fresh and dried amaranthus (*Amaranthus tricolor*) leaves. *J. Food Sci. Tech.*, 43 (5): 509-511.
- Singh, G. and Sehgal, S., (2008) Nutritional evaluation of laddoo prepared from popped pearl millet. *Nutr Food Sci.*, 38(4):310–315
- Sinha, N.K., Chattopadhyay, J.C., Das, P.K., Maiti, S. and Maiti, K. (2013). Prevalence of anemia and its possible attributing factors in psychologically healthy women of reproductive ages in Midnapore (Jangalmahal-area), India. *Indian J Community Health*; 25(3): 226-232
- State of Food security in the World (2013): The multiple dimensions of food security (2013), <http://www.fao.org/docrep/018/i3434e/i3434e.pdf>
- Stevens, G.A., Finucane, M.M., De-Regil, L.M., Paciorek, C.J. and Flaxman, S.R. (2013) Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995-2011: a systematic analysis of population representative data.; *The Lancet Global Health* 1: 16-25
- Sucharitha, K.V. (2013). Zinc status in adolescent girls of low income groups. Presented at 2nd International Conference and Exhibition on Nutritional Science and Therapy, Philadelphia, USA. (Available at: <http://www.omnicsgroup.com/conferences/nutritional-sciencetherapy-2013>) (Accessed on: 20.09.2013)
- Suliburska ,J. and Krejpcio, Z. (2014). Evaluation of the content and bioaccessibility of iron, zinc, calcium and magnesium from groats, rice, leguminous grains and nuts. *J Food Sci Technol* (March 2014) 51(3):589–594 DOI 10.1007/s13197-011-0535-5
- Tandon, B.N.,(2002). Report of the Indo-US Workshop on Nutrition and Health of Women, Infants and Children. National Institute of Nutrition, Hyderabad. February 10-12, 2000. *Nutrition Reviews*, 60(5): (II)S1.
- Teucher, B., Olivares, M. and Cori, H. (2004). Enhancers of iron absorption: ascorbic acid and other organic acids. *Int J Vitam Nutr Res*; 74(6): 403-19.
- Thankachan, P., Muthayya, S., Walczyk, T., Kurpad, A.V. and Hurrell,R.F.(2007): An analysis of the etiology of anemia and iron deficiency in young women of low socioeconomic status in Bangalore, India. *Food Nutr Bull.*, 28(3):328–336.
- Thimmayamma, B.V, Rau P, and Rao, K.V. (2003). Socioeconomic status, diet and nutrient adequacies of different population groups in urban and rural Hyderabad. *Indian J Nutr Diet* 1982;19:173-83.
- Thompson, B. and Amoroso L.(2011). Combating micronutrient deficiencies: food-based approaches. Rome, Italy .,The Food and Agriculture Organization of the United Nations and CAB International; 2011.

- Torheim, L. E., Ferguson, E. L., Penrose, K., and Arimond, M. (2010). Women in resource-poor settings are at risk of inadequate intakes of multiple micronutrients. *The Journal of Nutrition*, 140(11), 2051S–8S. <http://doi.org/10.3945/jn.110.123463>
- Toteja, G.S., Singh, P., Dillon, B.S., Saxena, B.N., Ahmed, F.U., Singh, R.P., Prakash, B., Vijayaraghavan, K., Singh Y, Rauf, A., Sharma VC, Gandhi, S., Behl, L., Mukherjee, K., Swami, S. S., Meru, V., Chandra, P. and Mohan U (2006). Prevalence of anemia among pregnant women and adolescent girls in 16 districts of India. *Food Nutrition Bulletin* 27(4): 311–15.
- Underwood, B.A. (1998). Prevention of vitamin A deficiency. In Prevention of micronutrient deficiencies. Tools for policymakers and public health workers. Eds., Howsen C.P., E.T. Kennedy and Horwitz A. Washington, DC: *Natl Acad Sci*, pp: 207 .
- United States Department of Agriculture (USDA) (2011) Food and Nutrition Information Center. <http://www.nal.usda.gov/fnic> [cited 2011]
- Vijayakumari, K., Pugalenti, M., and Vadivel, V. (2007) Effect of soaking and hydrothermal processing methods on the levels of antinutrients and in vitro protein digestibility of *Bauhinia purpurea* L. seeds. *Food Chem* 103:968–975 Zielinski H, Frias M, Mariusz K
- Wessells, K.R. and Brown, K.H. (2012). Estimating the global prevalence of zinc deficiency: results based on zinc availability in national food supplies and the prevalence of stunting. *PLoS One* 7: e50568.
- World Health Organization (1995). Physical Status: The Use and Interpretation of Anthropometry. Report of a WHO Expert Committee. Technical Report Series, No. 854. Geneva: World Health Organization.
- World Health Organization (2007). Assessing the iron status of populations: including literature reviews: report of a Joint World Health Organization / Centers for Disease Control and Prevention Technical Consultation on the Assessment of Iron Status at the Population Level, Geneva, Switzerland, 6–8 April 2004. – 2nd ed.
- World Health Organization (2008 a) Worldwide Prevalence of Anaemia 1993–2005 WHO Global Database on Anaemia. Geneva: WHO.
- World Health Organization (2009) Global prevalence of vitamin A deficiency in populations at risk 1995–2005. WHO Global Database on Vitamin A Deficiency. Geneva: World Health Organization.
- World Health Organization. (2008 b), The WHO Child Growth Standards. WHO, Geneva. Available from: <http://www.who.int/childgrowth/standards/en/>, accessed on December 24, 2008.)
- Yadav, S. and Sehgal, S. (2002). In vitro and in vivo availability of iron from bathua (*Chenopodium album*) and spinach (*Spinacia oleracea* L.) leaves. *J. Food Sci. Tech.*, 39 (1): 42-45.
- Yokoi, K., Sandstead, H. H., Egger, N. G., Alcock, N. W., Sadagopa Ramanujam,

- V. M., Dayal, H. H. and Penland, J. G. (2007). Association between zinc pool sizes and iron stores in premenopausal women without anaemia. *Br. J. Nutr.*, 98(6), 1214–1223. doi:10.1017/S0007114507803394
- Yoon, S. H., Kim, D.S., Yu, S.T., Shin, S.R. and Choi, D.Y. (2015). The usefulness of soluble transferrin receptor in the diagnosis and treatment of iron deficiency anemia in children, *Korean Journal of Pediatrics.*, 58(1):15-19
- Zimmerman, M.B., Winichagoon, P., Gowachirapant, S., Hess, S.Y., Harrington, M., Chavasit, V., Lynch, S.R. and Hurrell, R.F. (2005): Comparison of the efficacy of wheat based snacks fortified with ferrous sulfate, electrolytic iron, or hydrogen reduced elemental iron: randomized, double-blind, controlled trial in Thai women. *Am J Clin Nutr.*, 82:1276-1282

Websites

- http://fns.dpi.wi.gov/files/fns/pdf/ra_stand.pdf
- http://nnmbindia.org/1_NNMB_Third_Repeat_Rural_Survey_Technical_Report_26.pdf
- http://www.cdc.gov/healthyweight/assessing/bmi/childrens_bmi/about_childrens_bmi.html, accessed on 7 February 2013.
- <http://www.cibtech.org/jls.htm>
- <http://www.diet.com/g/regional-diet-american>
- <http://www.food-india.com>
- <http://www.metabolics.com/blog/a-practitioners-guide-to-zinc-supplements/>
- <http://www.nfsmi.org/documentlibraryfiles/PDF/20080215091404.pdf>
- <http://www.slideshare.net/obanbrahma/trace-minerals-60303676>
- <http://www.slideshare.net/princy77malik/role-of-iron-in-human-health>
- <http://www.tamilnadu.com>
- http://www.who.int/nutrition/publications/infantfeeding/essentail_nutrition_actions/en/