

V. SUMMARY AND CONCLUSION

Adolescence is a developmental period which may be conservatively understood as the years between the onset of puberty as well as the establishment of social autonomy. It occurs between ages 13 and 19 and is usually subjected to numerous exposures. Several studies in recent years had manifestly demonstrated that this section of the population group is going through more levels of psychological distress which included Depression, Anxiety and Stress (DAS) and that can lead to enormous problems in later years.

The world scenario witnesses that one in every eight people live with a mental disorder, the anxiety and depressive disorders as the most common ones (GHDx, 2022). WHO, in its 2021 report stated that more than 13% of adolescents aged 10–19 live with a diagnosed mental disorder with 40% of the disorders being anxiety and depression. UNICEF, 2021 has declared that every year almost 46,000 children between the ages of 10 and 19 end their own lives.

World Health Organization (WHO) report on India as cited in India Today (2022) conveyed that 56 and 38 million Indians suffer from depression and anxiety disorders respectively, out of which 14% were adolescents. Sagar et al. (2020) forewarn that the mental disorders as being one of the leading causes of non-fatal disease burden in India. Tamil Nadu, being one of the states in the high socio-demographic index group, accounted for a high prevalence of depressive and anxiety disorders. The Crude Disability Adjusted Life Years Rate of depressive disorders was highest in Tamil Nadu in the adolescent group (The Hindu, 2020). Indian Council of Medical Research survey conducted during the year 2022 revealed that 12-13% of students in Tamil Nadu suffers from psychological, emotional and Behavioural conditions.

The coinciding symptoms and behaviours of these three negative mental states can lead to all sorts of educational issues, which in turn impact the academic achievement of students. It has also been proved that academic performance from school education to the higher education level is influenced by the symptoms of stress, anxiety, and depression, as these mental disturbances lead to difficulties in concentration, lack of motivation and interest, poor attendance, and poor physical health (Ibrahim et al., 2013).

Depression, anxiety and stress in the school network are considered as significant markers for low degree of emotional well-being. The inability to recognize and manage these upsetting attributes would lamentably prompt or expand mental dreariness with bothersome effects all through their lives (Teh et al., 2015). Various studies had discovered the progress from youthfulness to adulthood loaded up with anxiety, stressfulness, and depression as well. Hence, these three negative mental states that affect the mental health conditions of adolescents need to be addressed.

Shastri (2009) stated that prevention and early intervention are recognized as key elements for minimizing the impact of any potentially serious health condition. He also cited the recommendation of WHO on early identification and intervention while examining the extent of the gap between the prevalence and treatment of psychiatric disorders. The investment made to identify the mental health problems early and intervene at the right time was cost-effective, as the intervention would be preventing further breakdown and avoid an adult treatment and rehabilitation programme which is much more expensive. Based on such evidences, it was well understood that there exists a pressing need to develop and improve adolescent mental healthcare models that could be implemented as preventive and early intervention strategies.

After a detailed exploration of the existing mental health care models, the investigator found it imperative to redesign and merge the two major interventions namely Cognitive Behavioural Therapy (CBT) and Stress Inoculation Training (SIT). CBT has proved its effectiveness in alleviating anxiety and depression through the psychotherapy technique, wherein the individual could recognise and change the incorrect and/or negative thoughts that have been influencing their behaviour. CBT is always one of the identified suitable treatment methods for individuals with anxiety and depression. Miller et al. (2012) also stated that several methodical reviews of CBT-based interventions in the school setting have been done, with the main focus on prevention of anxiety disorders and depression.

Whereas SIT, though had found an domineering place in clinical setting to reduce stress, it has a unique characteristic of customising for the normal population by preparing an individual to quickly defend against stress triggers by exposing the individual to a milder form of stress (just like vaccination). Stress Inoculation Training (SIT), a set of Cognitive Behavioural Techniques (CBT) was developed as a treatment by Donald

Meichenbaum with the objective of facilitating a person to cope up with the magnitude of facing stressful. SIT works on a preventative basis in terms of inoculating the person to present and the future stressor. The uniqueness of SIT is that it is ready to be tailored to the patron's needs. The SIT was pilot tested among 588 middle school students as per the local needs by Maraichelvi (2016) in the term of School Stress Inoculation Training (SSIT) and was found to be effective in reducing the stress levels of the experimental group of students. On the whole, the term 'inoculation' implies, SSIT was designed to impart skills to enhance resistance to stress.

Put together, adolescence is a transitional period and they are overloaded with multiple things. Hence, the prevalence of mental disorders is common among them. The relationship between the development of depression, anxiety and stress among adolescents becomes a promising area of inquiry. Hence, identification of the best preventive measure to combat depression, anxiety and stress among school-going adolescents at the initial stage has become essential. Accordingly, a preventive intervention model (merging CBT and SIT) was formulated and titled as Comprehensive School-based Intervention Training to alleviate Depression, Anxiety and Stress (CSIT-DAS). The outcome of the intervention was to build a repertoire of coping skills that could improve their adaptability to physical, mental, and psycho-social changes as it is directly linked with their overall well-being.

Objectives

Primary Objective: To formulate, and analyse the effectiveness of a suitable preventive intervention to alleviate Depression, Anxiety and Stress (DAS) in school-going adolescents

Secondary Objectives

- Explore the level of Depression, Anxiety and Stress (DAS) and its interrelationship among the selected school-going adolescents
- Assess the influence of socio-demographic factors on Depression, Anxiety and Stress (DAS) of the selected school-going adolescents
- Identify the potential determinants triggering Depression, Anxiety and Stress (DAS) among the selected school-going adolescents
- Identify and customize a suitable preventive intervention to alleviate Depression, Anxiety and Stress (DAS) among school-going adolescents and

- Analyze the effectiveness and the sustainability of the formulated preventive intervention.

Based on the objectives framed above certain null hypothesis were formulated as given below:

- H₀ 1- There is no relationship between Depression, Anxiety and Stress (DAS)
- H₀ 2 - There is no difference in the levels of Depression, Anxiety and Stress (DAS) based on gender
- H₀ 3- There is no difference in the levels of Depression, Anxiety and Stress (DAS) based on class of study
- H₀ 4 - There is no difference in the levels of Depression, Anxiety and Stress (DAS) based on Area of Residence
- H₀ 5 - There is no difference in the levels of Depression, Anxiety and Stress (DAS) based on Family type
- H₀ 6 - There is no difference in the levels of Depression, Anxiety and Stress (DAS) based on the education status of the father
- H₀ 7 - There is no difference in the levels of Depression, Anxiety and Stress (DAS) based on the education status of the mother
- H₀ 8 - There is no relationship between Depression and its potential determinants
- H₀ 9 - There is no relationship between Anxiety and its potential determinants
- H₀ 10 - There is no relationship between Stress and its potential determinants
- H₀ 11 - The formulated preventive intervention shows no difference in the levels of Depression, Anxiety and Stress (DAS) among the experimental group of respondents based on gender and
- H₀ 12 - The formulated preventive intervention does not decrease the level of Depression, Anxiety and Stress (DAS) among the experimental group of participants.

METHODOLOGY

The methodology of the study were discussed based on the two phases of study:

- ✓ Phase I: Identify the levels of Depression, Anxiety and Stress (DAS), their interrelationship, and their determinants
- ✓ Phase II: Effectiveness of the formulated preventive intervention

Phase I: Identify the levels of Depression, Anxiety and Stress (DAS), their interrelationship, and their determinants

A descriptive cross-sectional study predominantly adopting questionnaires was embraced. Coimbatore district in the state of Tamil Nadu was selected as the area for the study. The urban and the extended corporation wards (referred to as semi-urban) of each of the five zones were selected and only higher secondary schools with certain inclusion criteria from each of the places were taken into consideration. Accordingly, a total of 30 schools from all five zones of Coimbatore District were shortlisted. However, after seeking permission from the administrators of all the short-listed schools, only a total of fifteen schools expressed their willingness to participate and permitted the researcher to collect data.

The study focused on target respondents of adolescents enrolled in Class 8 and Class 9, as it was during this stage the adolescents were getting ready for the Board exams (10th, 11th and 12th standard based on the Indian Education System) in a year or two. They were exposed to much psychological adversity in the process of getting prepared to face board exams added to the burden of transitional hardships. All students from classes 8 and 9 from the 15 schools constituted the population sample. Accordingly, the population sample accounted for a total of 1038 students. The tools used were;

- *General Profile Questionnaire*: This tool was used to collect the general and personal profile of the selected school-going adolescents.
- *Standardized Depression, Anxiety and Stress Scale (DASS-42)*: A standardized tool developed by Lovibond and Tilteld as DASS-42 was used to determine the levels of DAS among the school-going adolescents. The scale had 42 items segregated into three mental states – Depression, Anxiety and Stress having 14 items each. Higher the score, higher was the DAS level in them.
- *Self-formulated Scale on determinants of Depression, Anxiety and Stress*– A scale to identify the determinants of DAS among the school-going adolescents was formulated as three subscales (one for each mental state). The number of determinants and the items within each determinant varies for each sub-scale based on the exploratory factor analysis. The details were as follows:
 - a) Determinants of Depression scale composed of five determinants namely Attitudinal issues, Present/future life concerns, Academic pressure, Family & relationship issues, and Psychological concerns with 6, 5, 3, 3, and 3 items respectively.

- b) Determinants of anxiety scale composed of four determinants namely Generalized panic, Social anxiety, Social phobia, and Academic anxiety with 6, 4, 3, and 3 items respectively.
- c) Determinants of Stress scale composed of four determinants namely Family stress, Drastic change in life, Attitudinal issue and Academic stress with 6, 5, 4, and 2 items respectively.

The reliability and validity of the scales were tested. The items were formulated in such a way that higher the score in the determinant scale lower was the effect of determinants in causing the negative mental states.

As the data of the first phase was significantly deviating from normal distribution, nonparametric tests were performed. Both qualitative (Chi-Square) and quantitative analysis (Mann Whitney test for two independent variables and Kruskal - Wallis Test for more than two) were performed to see the significant association between socio-demographic variables and levels of DAS. Spearman's Rank Correlations was carried out to see the significant correlation between DAS and as well as for the determinants of DAS.

Phase II: Effectiveness of the formulated preventive intervention

As the study embodies the implementation of the preventive intervention (CSIT-DAS) and analyse its effectiveness in terms of alleviated DAS, out of the 15 schools identified with a sample size of 1038, only 2 schools provided consent and were ready to spare a few structured hours within a period of 4 months needed for intervention. Out of these two schools, the school with a greater mean score on DAS constituted the experimental group and the other school was considered the control group. Accordingly, all students in the 8th and 9th classes of these two schools constituted the sample for the second phase of the study (Experimental group - 60 students, Control group – 60 students).

The tool for this phase was the preventive intervention called as Comprehensive School-based Intervention Training to alleviate Depression, Anxiety and Stress abbreviated as (CSIT-DAS). Understanding the prevalence of mental health problem in terms of DAS among the school-going adolescents, the researcher focused on a primary intervention model as it aims to prevent the adolescent who were at the risk to become exposed to DAS. An intervention customizing the CBT and SIT framework in accordance with the results of the first phase of the study, was formulated by a team of experts in the field of Psychology, and implemented by a handful of experienced trainers. The

Institutional Ethical clearance approved the study with an approval number - IHEC /17-18HD/29.

The CSIT-DAS, exclusively designed for the student population, was formulated to be implemented through four-stage training phases namely: IQ Test & Trust Building; Orientation and pretest evaluation; Education and Training (Psycho Education and Behavioural & Cognitive training); and Posttest evaluation with stipulated follow ups. The intervention was conducted in 15 sessions (2 hours each) spread across four months.

The data set of the phase II test results was not significantly deviating from normal distribution, hence parametric tests were used. A Two Way MANOVA of repeated measures [2 (experimental and control groups) x 2 (Pre and Post evaluation scores of the three dependent variables, depression, anxiety and stress)] was computed to find the effectiveness of CSIT-DAS. To examine the sustainability of the CSIT-DAS, two-way MANOVA of repeated measures (Pre intervention, Post intervention, Follow up 1 and Follow up 2) was carried out.

RESULTS AND DISCUSSION - KEY FINDINGS

Phase I: Identify the levels of Depression, Anxiety and Stress (DAS), their interrelationship, and their determinants

A. General profile of the respondents

- ✓ Female respondents (50.7%) were more than the male respondents (49.3%)
- ✓ Most of the respondents were from 9th standard (51.4%) followed by class 8 (48.6%)
- ✓ Around 60% of them hailed from nuclear family and 30% from by joint family
- ✓ The ratio of respondents from urban, semi-urban and rural stands at 2:1:1
- ✓ Educational status of parents – The mothers' educational status though seems to be on par with fathers' educational status, the %age of mothers (8.8%) surpassed fathers (4.6%) in terms of not having formal education.

B. Categorization of the respondents based on the levels of Depression, Anxiety and Stress

- ✓ Majority of them were moderately depressed (35%), followed by severely depressed (18.2%) and extremely severe (5.4%).
- ✓ The threshold score for the participants being labelled as anxious was also categorized under five categories, wherein, 35.5% (368 out of 1038) were anxious at a moderate level. Also, it was observed that 25.8% and 15.9% had severe and extremely severe anxiety respectively.

- ✓ The percentage of (32.9%) of the respondents who were moderately stressed was more in number (342) when compared to the other levels. The table further states that a good percentage of respondents were severely stressed (14.5%) and extremely severely stressed (3.4%).
- ✓ Anxiety tops the list with 91% of the selected respondents being affected followed by depression (84.2%) and stress (75%).

C. Interrelationship between Depression, Anxiety and Stress of the respondents

- ✓ The three mental issues – Depression, Anxiety, and Stress have a significant positive moderate association with each other.
- ✓ Depression and anxiety had a moderate positive relationship ($R = .495, p < .01$) with each other
- ✓ Stress with a weak positive association with depression ($R = .378, p < .05$) and anxiety ($R = .301, p < .05$)
- ✓ Therefore H_0 1 Stating that ‘there is no relationship between Depression, Anxiety and Stress (DAS)’ stands rejected

D. Association between sociodemographic variables and levels of Depression, Anxiety and Stress of the respondents

- ✓ The stress means score was high among male students (19.03) than female counterparts (18.64). The anxiety and depression scores revealed that females were more anxious and depressed (14.17 and 15.66 respectively) in comparison to males (13.74 and 15.34 respectively). However, the Mann-Whitney U test reveals that these results were not statistically significant for Depression with $|z| = 0.95, p > 0.05$; Anxiety with $|z| = 1.27, p > 0.05$ and Stress with $|z| = 0.66, p > 0.05$. Hence gender did not have an impact on DAS, thereby H_0 2 - ‘there is no difference in the levels of Depression, Anxiety and Stress (DAS) based on gender’ was accepted
- ✓ The Mann-Whitney U test between the 8th and 9th class of study reveals a statistically significant value of ($|z| = 2.85; p < .01$) in relation to stress thereby accepting the hypothesis H_0 3 for depression and anxiety and rejecting for stress. The comparison of the respondents’ mean scores on stress was found to be high among class 9 against class 8 (19.49 against 18.18).
- ✓ The significance of the association between the area of residence (Rural, Semi-Urban, Urban) of the selected respondents in relation to the DAS mean score, reported a

statistically significant value of $(\chi^2 (2) = 9.651; p <.01)$ for anxiety. Hence the hypothesis H₀ 4 is accepted for depression and stress but rejected for anxiety.

- ✓ Type of family does not significantly influence the occurrence of DAS ($\chi^2 (2) = 3.084; p >.05$); ($\chi^2 (2) = 4.374; p >.05$); ($\chi^2 (2) = 3.453; p >.05$) among the selected section of respondents and hence hypothesis H₀ 5 is accepted.
- ✓ The mean scores of the respondents in the stress domain were found to be amplified among the students of graduated fathers (19.85), closely followed by the children of fathers who had their primary school level of education (18.87). A statistically significant difference between the levels of fathers' education and the stress mean score at a 5% level ($\chi^2 (2) = 11.748; p <.05$) was noted only for the stress domain, thereby the hypothesis H₀ 6 is accepted for depression and anxiety and rejected for stress.
- ✓ A statistically significant association was found between the level of mothers' education and the anxiety scores ($\chi^2 (2) = 14.992; p <.01$). Hence the hypothesis H₀ 7 is accepted for depression and stress and rejected for anxiety. The mean scores showed that the children of mothers with no formal education were more anxious.

E. Relationship between Depression, Anxiety and Stress and its potential indicators among the respondents

- ✓ *Determinants of Depression:* Though all the five determinants are significantly correlated with the depression score, psychological concerns ($r_s = -.741, p = .0001$) attitudinal issues, ($r_s = -.728, p = .0001$) and family and relationship issues ($r_s = -.701, p = .0001$) had a strong negative correlation to the depression of the selected respondents. Whereas the two determinants namely present/future life concerns ($r_s = -.342, p = .0001$) and academic pressure ($r_s = -.305, p = .0001$) had a weak negative correlation. Hence the hypothesis H₀ 8 that stated 'there is no relationship between Depression and its potential determinants' was rejected.
- ✓ *Determinants of Anxiety:* The relationship between generalized panic ($r_s = -.612, p = .0001$) and specific phobia ($r_s = -.745, p = .0001$) had a strong negative correlation to the anxiety of the selected respondents. Whereas, social anxiety ($r_s = -.581, p = .0001$) and Academic anxiety ($r_s = -.426, p = .0001$) had a moderate negative correlation. Hence the hypothesis H₀ 9 that stated 'there is no relationship between Anxiety and its potential determinants' was rejected.
- ✓ *Determinants of Stress:* Stress in the students had a very strong negative correlation with two potential determinants namely family stress ($r_s = -.718, p = .0001$) and

attitudinal stress ($r_s = -.707, p = .0001$). Drastic life changes ($r_s = -.317, p = .001$) and academic stress ($r_s = -.236, p = .004$) though had a significant correlation, the magnitude of the relationship was weak. Hence the hypothesis H_0 10 that stated ‘there is no relationship between Stress and its potential determinants’ was rejected.

Phase II: Effectiveness of the formulated preventive intervention

- **Efficacy of Comprehensive School-based Intervention Training to alleviate Depression, Anxiety and Stress - Comparison between pre-test and post-test Depression, Anxiety and Stress levels between Experimental and Wait-list Control Group**
- ✓ The experimental group of respondents’ mean score showed a tremendous decline in the post-evaluation measures on all the three dependent variables Depression (8.72 from 25.43), Anxiety (7.55 from 24.03) and Stress (13.18 from 25.28). On the other hand, the control group of respondents’ mean score on all the three dependent variables was augmented {Depression (22.60 from 18.92), Anxiety (18.55 from 14.30) and Stress (24.00 from 20.22)}
- ✓ Multivariate tests indicates a statistically significant effect confirming that the experimental and control group do indeed differ in terms of intervention. The effect of time measures endorse that the pre and post-level of DAS vary in terms of time. The partial eta square of both the 2 groups of intervention 32.7% and 2 groups of time measures 62% showed a large effect size.
- ✓ The Multivariate tests also showed a statistically significant interaction between intervention and time over DAS and explained 82.5% variance between the interacting groups
- ✓ To determine which of the dependent variable accounts for the significance in multivariate tests, univariate analysis was computed. The interaction between the group and the time indicated a significant effect on all the three dependent variables – Depression ($\eta^2 = .770, p = .0001$), Anxiety ($\eta^2 = .785, p = .0001$) and Stress ($\eta^2 = 2.17, p = .0001$). It also showed a large effect size between the group and time measure.
- ✓ The MANOVA plots showing the change in the direction and magnitude of the mean score of the three dependent variables (Depression, Anxiety and Stress) with reference to the two-time measure (Pre and Post) vividly portrays that the mean score of the experimental group has a significant decline in all the three dependent variables. This

decline is indicative of lower depression, anxiety and stress among the experimental group respondents. Whereas, for the control group, the mean score showed an inclination towards higher score in all the three dependent variables. The increase in the score is indicative of higher depression, anxiety and stress among the control group respondents.

- ✓ Hence hypothesis H₀ 11 that stated ‘the formulated preventive intervention shows no difference in the levels of Depression, Anxiety and Stress (DAS) among the experimental group of respondents based on gender’ was accepted.
- **Sustainability of Comprehensive School-based Intervention Training to alleviate Depression, Anxiety and Stress (CSIT- DAS) among the experimental group**
- ✓ The decline in the mean score of all the three dependent variable – depression (7.80 from 25.43), anxiety (6.40 from 24.03) and stress (9.75 from 25.28) - within the four-time repeated measures authenticates the sustainability of the intervention titled CSIT-DAS. However the follow up 2 score in both anxiety and stress (6.40 and 9.75 respectively) had a mild increase in mean score when compared with follow up 1 (6.35 and 9.27 respectively)
- ✓ The multivariate tests of the interaction between the four times measures of the experimental group of participants was significant with partial eta square showing a strong effect size ($\eta^2 = .840, p = .0001$) on the three dependent variables put together.
- ✓ To find out which of the dependent variable contributed to the combined significant effect, univariate analysis of variance was computed. The (partial eta square) η^2 of .639, .681, and .502 respectively, demonstrates large effect that the time measures (4 measures) had on the three dependent variables – Depression, Anxiety and Stress. Hence, the two ways repeated measures MANOVA statistics has rejected the hypothesis H₀ 12 – The formulated preventive intervention does not decrease the level of Depression, Anxiety and Stress (DAS) among the experimental group of participants.

Depression, Anxiety and Stress directly or indirectly have an impact on one’s lifestyle in different aspects. These issues mandate research in the domain of mental health. The key finding of the present study is that out of 1038 total respondents, majority of them had perceived to be depressed, anxious, and stressed, at varying levels between mild and extremely severe where more children were found to be anxious, followed by depression

and stress. All the five determinants of depression are significantly correlated. All the four determinants of anxiety and stress are also significantly correlated. The CSIT-DAS formulated for this sect of population based on the theoretical models of CBT and SIT and the results of the first phase of study, was found to be effective in two terms. Firstly, the experimental group showed a tremendous decline in the post-intervention mean score on DAS compared to the pre-evaluation mean score and was statistically significant. Secondly, the sustainability of CSIT-DAS was also witnessed with the stipulated follow up cycles. The finding clearly states that the CSIT-DAS should be a continuous training with periodic sessions of rejuvenation.

RECOMMENDATIONS AND IMPLICATIONS

- Establishing a student counselling centre within the school campus handled by qualified staff is highly recommended. Altogether, every school need to have a full time counsellor with a deeper understanding of the principles of child and adolescent development.
- The teachers should be familiarised with basic coping techniques such as knowing the "Do's and Don'ts" on students who experience negative mental health symptoms.
- Schools should be encouraged to create programmes such as individual and group interventions by applying one-on-one interviews with the guidance counsellors of the school and occasional programmes that promote and explain good mental health on both parents and the student. Parents should also be encouraged by school authority to participate in students' performance activities.
- Launching programmes focused on mental health awareness and culture change, with the goal of enlightening an environment in which students feel encouraged and empowered to care for their overall well-being is strongly recommended.
- Since mental health problems are so common among this sect of population, routine administration of assessment test should be encouraged in monitoring the mental problems among early adolescents
- The tool DASS-42 is just a screening tool so children who are depressed, anxious and stressed at extremely severe level should be referred to clinical assessment. Moreover, the alarming prevalence rate of DAS in this sect of the population has to be immediately addressed, as DAS poses a threat to the holistic living of adolescents. Although many schools have a self-formulated school health programme, adequate importance is not given in all the dimensions.

- The alarming risk of DAS among school students is also caused by the family pressure to perform better; hence the researcher feels that the focus of intervention should not only be on children but also on the parents.
- CSIT-DAS has boosted the students' awareness and developed a sense of mastery over the usage of coping skills, which undoubtedly validates the efficacy of the CSIT-DAS intervention provided to them. Therefore, CSIT-DAS especially tailored for school children is strongly recommended to be used by every school to alleviate the risks of mental issues prevalent among the student population. .