



INFLUENCE OF VITAMIN-E AND SELENIUM SUPPLEMENTATION ON THE GROWTH PERFORMANCE AND ANTIBODY RESPONSES OF LAYER CHICKS

G. KANCHANA ^{*1} AND G. P. JEYANTHI ²

¹Department of Biochemistry, Muthayammal College of Arts & Science, Rasipuram, Tamilnadu, India.

²Department of Biochemistry, Biotechnology and Bioinformatics, Avinashilingam University for Women, Coimbatore, Tamilnadu, India

*Corresponding Author kanchmoorthi@yahoo.co.in

ABSTRACT

An experiment was conducted to study the effect of vitamin E and selenium yeast supplementation on the growth performance and some immune parameters with two hundred and ten commercial straight run day-old layer chicks. The chicks were randomly allotted into seven treatment groups with three replicates of ten chicks each. The chicks were fed basal diet (T₁), basal diet with 100 mg/kg vitamin E (T₂), basal diet with 200 mg/kg vitamin E (T₃), basal diet with 0.2 mg/kg selenium (T₄), basal diet with 0.4 mg/kg selenium (T₅), basal diet with 100 mg/kg vitamin E plus 0.2 mg/kg selenium (T₆), and basal diet with 200 mg/kg vitamin E plus 0.4 mg/kg selenium (T₇) for eight weeks period. No significant difference was found in the body weight of layer chicks for the first four weeks among the treatment groups but the body weight of groups (T₆ and T₇) receiving both vitamin E and selenium was significantly increased ($P < 0.05$) at the end of 8 weeks period. The feed intake of layer chicks did not vary significantly between treatment groups. Chicks receiving supplements of 100 mg vitamin E/kg and 0.2 mg selenium/kg produced significantly higher HA titre against SRBC ($P < 0.01$), HI titre against NDV ($P < 0.01$) and QAGPT titre against IBDV ($P < 0.01$). The results suggested that vitamin E and selenium might provide higher level of protection against natural immune challenges.

KEY WORDS

Trace elements, antibody, immunity, antioxidant, selenium

INTRODUCTION

Poultry is one of the fastest growing segments of the agricultural sector in developing countries today. In context of poultry industry, the birds encounter numerous stressors during their lives. These stressors cause hormone changes, reduces feed intake, alter nutrient metabolism and suppresses immune function. Therefore, it is highly essential to find ways and means for enhancement of immune response by nutritional manipulation. The

immune system benefits greatly from proper nutrition of the bird and it also prepare the bird for periods of stress, reducing the adverse effects of stress and enhancing recovery from stressful periods. Therefore, it is an increasing challenge to formulate diets that allow highly productive, intensively-reared modern livestock and poultry to reach genetic potential. In recent years, there has been a growing understanding that marginal vitamin and trace element status is a factor limiting health and



INFLUENCE OF VITAMIN-E AND SELENIUM SUPPLEMENTATION ON THE GROWTH PERFORMANCE AND ANTIBODY RESPONSES OF LAYER CHICKS

productivity. It was reported that, nutritionists have to take into account several factors including stress management and immunity enhancement while formulating feed¹.

In birds, free radical generation and lipid peroxidation are responsible for the development of various diseases as well as for a decrease in bird's productivity and product quality^{2,3}. In this respect, antioxidants play an important role in maintaining bird health, productive and reproductive performance⁴. Of the known antioxidants within animal cells, vitamin E, and selenium are considered the most important. Vitamin E functions as a chain breaking antioxidant which prevents free radical induced oxidative damage by trapping reactive oxyradicals in biological membranes⁵. Selenium, an essential component of the enzyme glutathione peroxidase (GSH-Px) destroys peroxides before they can damage body tissues. GSH-Px concentration and activity are directly related to the selenium status of the animal⁶. Selenium and vitamin E are inter-related; hence complete protection of living cells requires both vitamin E and selenium in the diet. Traditionally, selenium has been added to poultry diet in the form of inorganic sources either as sodium selenite or sodium selenate⁷. Selenium yeast has been reported to be an excellent source of organic selenium⁸. Results of current studies provided evidence that organic form of selenium is generally safer and better absorbed⁹.

Even though most nutrients required by the immune system are present in the diet in sufficient concentrations, there is evidence that increased dietary supplementation of certain nutrients especially antioxidants, above that needed for maximum growth and feed efficiency, is of benefit to the immune response. Thus, the aim of the present investigation was to assess the effect of dietary supplementation of different combinations of vitamin E and selenium in excess of the recommended level either independently or

simultaneously on growth and immune response of layer chicks.

MATERIALS AND METHODS

Two hundred and ten commercial straight run day-old layer chicks belonging to a single hatch were purchased from Venkateswara Hatcheries, Namakkal. The birds were wing banded, weighed and randomly allotted into seven treatment groups with three replicates of ten chicks each. The chicks were fed basal diet (T₁), basal diet with 100 mg/kg vitamin E (T₂), basal diet with 200 mg/kg vitamin E (T₃), basal diet with 0.2 mg/kg selenium (T₄), basal diet with 0.4 mg/kg selenium (T₅), basal diet with 100 mg/kg vitamin E plus 0.2 mg/kg selenium (T₆), and basal diet with 200 mg/kg vitamin E plus 0.4 mg/kg selenium (T₇).

The chicks of all treatment groups were reared in cages under standard managerial condition throughout the study period of 8 weeks. The experimental diet was formulated according to the standard prescribed in Bureau of India Standards¹⁰; except the vitamin E and selenium level in basal diet. Vitamin E in the form of dl- α -tocopheryl acetate (Promix E) and selenium in the form of Selplex containing mainly as selenomethionine were incorporated into the basal diet either independently or simultaneously in the basal diet. The chicks were fed with weighed quantity of feed ad libitum and have free access to whole some water throughout the investigation period.

(i) Growth rate study

Individual body weight of the chicks in each replicate in all the treatment groups was recorded at the end of every 28 days period up to 8 weeks of age. Feed consumption of all the treatment groups was recorded once in 28 days period and the mean total feed consumption per bird was calculated.



INFLUENCE OF VITAMIN-E AND SELENIUM SUPPLEMENTATION ON THE GROWTH PERFORMANCE AND ANTIBODY RESPONSES OF LAYER CHICKS

(ii) Immunity against Sheep Red Blood Cells

Whole blood was collected from Mecheri breed of sheep in Alsever's solution (5.125 g of dextrose, 2 g of sodium citrate and 1.05 g of sodium chloride dissolved in 250 ml of distilled water). The sheep red blood cells (SRBC) were washed thrice in phosphate buffered saline (PBS, pH 7.4) and resuspended in PBS to prepare 25 % (v/v) SRBC suspension. On the twenty eighth day of age, six chicks per dietary treatment were randomly picked up and immunized with 0.5ml of 25% SRBC on each thigh muscle¹¹. Blood samples were collected fifteen days after immunization and haemagglutination (HA) titre was assessed. HA titre was expressed as the log₂ of the reciprocal of the highest dilution showing 100% agglutination¹².

(iii) Immunity against Newcastle Disease Virus

Blood samples were collected randomly from six birds in each treatment group fifteen days after immunization with Newcastle disease vaccine (Ventri Biologicals, Pune, India). The serum samples were separated and utilized for the haemagglutination inhibition (HI) test to find out the immunity developed against the Newcastle disease virus (NDV)¹³.

(iv) Immunity against Infectious Bursal Disease Virus

Blood samples were collected randomly from six birds in each treatment group fifteen days after immunization with IBD vaccine (Ventri Biologicals, Pune, India). The serum samples were separated and utilized for analyzing the antibody level. Quantitative agar gel precipitation test (QAGPT) was used to assess the antibody level against Infectious Bursal Disease Virus (IBDV) as per the method described by Wood et al., with slight modification¹⁴.

About 1.2g of agar was dissolved in 100 ml hypertonic saline (8% sodium chloride) and boiled. To the above boiled solution, 0.02% sodium azide was added as a preservative to prevent bacterial

contamination. The solution was cooled to 50-55° C and then cast approximately 5 ml on the clean glass slide and allowed for solidification. After solidification, with the aid of gel puncher two sets of wells were made on the agar plates, each set was made of a central well surrounded by six equidistant satellite peripheral wells of about 5 mm diameter with an inter-space of 0.3 cm. The peripheral wells were numbered from 1 to 12. The two central wells were loaded with infectious bursal disease virus antigen. The wells 6 and 7 were kept as known negative and positive control respectively. The first peripheral well was filled with test serum and in the remaining wells, serial two fold dilutions of test serum were added. The slides were then incubated at 37° C for 24 to 28 h in a humid chamber. The reciprocal of the highest dilution of serum showing precipitin line was taken as the QAGPT titre of the serum.

(v) Statistical analysis of data

The collected data on various parameters were subjected to analysis of variance for statistical significance as per the methods of Snedecor and Cochran¹⁵.

RESULTS


1. Growth performance

The influence of vitamin E and selenium supplementation independently and simultaneously on the body weight gain of layer chicks from 1 to 8 weeks of age is presented in Figure 1. The mean body weight of layer chicks during the first 4 week growth period did not differ significantly among the treatment groups. The body weight was significantly ($P < 0.05$) higher in T₆ and T₇ at the end of eighth week, as compared to control and other treatment groups. However, no significant difference was observed between the experimental groups T₂, T₃, T₄ and T₅.



INFLUENCE OF VITAMIN-E AND SELENIUM SUPPLEMENTATION ON THE GROWTH PERFORMANCE AND ANTIBODY RESPONSES OF LAYER CHICKS

Figure 1 Effect of vitamin E and selenium supplementation on body weight gain of layer chicks




2. Feed consumption

The mean daily feed consumption of layer chicks from 1 to 8 weeks of age as influenced by supplementation of vitamin E and selenium independently and simultaneously is presented in Figure 2. The feed intake of layer

chicks did not differ significantly among the treatment groups throughout the investigation period. However, the treatments groups (T₆ and T₇) that received combination of vitamin E and selenium had lower feed consumption.

Figure 2 Effect of vitamin E and selenium supplementation on feed consumption of layer chicks



3. Immunity

The HA, HI and QAGPT antibody titres of layer chicks are shown in Table 1. The

mean log₂ HA titre against SRBC was significantly ($P < 0.01$) higher in T₆ and T₇ with mean values of 4.88 and 4.50 respectively as compared to T₁



INFLUENCE OF VITAMIN-E AND SELENIUM SUPPLEMENTATION ON THE GROWTH PERFORMANCE AND ANTIBODY RESPONSES OF LAYER CHICKS

which recorded HA titre of 2.13. The HI titre against NDV was significantly ($P < 0.01$) higher in T₆ (5.13) and T₇ (4.63) followed by selenium supplemented groups T₄ (4.25) and T₅ (3.88). The lowest HI titre was recorded in T₁, T₃ and T₂ with

mean values of 2.88, 3.50 and 3.63 respectively. The QAGPT titre against IBDV was significantly ($P < 0.01$) higher in T₆ (3.50) followed by T₇ (3.25) as compared to the control and other treatment groups.

Table 1

Mean (\pm S.E.) log₂ HA titre against SRBC, HI titre against NDV and QAGPT titre against IBDV as influenced by supplementing vitamin E and selenium

Treatment groups	HA titer against SRBC	HI titre against NDV	QAGPT titre against IBDV
T ₁ - Control	2.13 \pm 0.13 ^A	2.88 \pm 0.23 ^A	2.25 \pm 0.16 ^A
T ₂ - Vitamin E 100 mg/kg	3.38 \pm 0.18 ^B	3.63 \pm 0.18 ^{AB}	2.75 \pm 0.25 ^{AB}
T ₃ - Vitamin E 200 mg/kg	3.13 \pm 0.23 ^B	3.50 \pm 0.19 ^{AB}	2.50 \pm 0.27 ^{AB}
T ₄ - Selenium 0.2 mg/kg	3.75 \pm 0.25 ^{BC}	4.25 \pm 0.25 ^{BC}	3.00 \pm 0.19 ^{AB}
T ₅ - Selenium 0.4 mg/kg	3.63 \pm 0.26 ^B	3.88 \pm 0.23 ^{BC}	2.88 \pm 0.23 ^{AB}
T ₆ - Vit-E 100 mg/kg + Se-0.2 mg/kg	4.88 \pm 0.23 ^D	5.13 \pm 0.13 ^D	3.50 \pm 0.19 ^C
T ₇ - Vit-E 200 mg/kg + Se-0.4 mg/kg	4.50 \pm 0.19 ^{CD}	4.63 \pm 0.18 ^{CD}	3.25 \pm 0.16 ^{BC}

Value given in each cell is the mean of six observations.

^{A-D} Means within a column with no common superscript differ significantly ($P < 0.01$).

DISCUSSION

The supplementation of vitamin E (100 & 200 mg/kg) and selenium (0.2 & 0.4 mg/kg) independently and simultaneously in the basal diet improved the body weight gain of layers chicks. However, higher body weight gains were recorded

in combination of vitamin E and selenium supplemented groups (T₆ & T₇) than vitamin E and selenium alone fed group. Salman et al. ¹⁶ showed that body weight of broilers was higher ($P > 0.05$) in vitamin E (250 mg/kg) and vitamin E plus organic selenium supplemented groups (250 mg+0.2 mg/kg) than that of control and vitamin E plus inorganic selenium supplemented groups (250



INFLUENCE OF VITAMIN-E AND SELENIUM SUPPLEMENTATION ON THE GROWTH PERFORMANCE AND ANTIBODY RESPONSES OF LAYER CHICKS

used, age of animals and the amount of vitamin E and selenium supplemented to the diet.

The present findings suggest that vitamin E may have an immunomodulator effect, increasing the resistance to diseases. A possible mechanism of this enhanced immune function may be a down regulation of the biosynthesis of prostaglandins, which inhibit several immunity parameters²⁸. Vitamin E and GSH-Px are two molecules that help to prevent the oxidative damage. Free radicals are scavenged by vitamin E as a first line of defense and selenium as a constituent of cytosolic enzyme glutathione peroxidase convert free radicals to inert substances rendering them harmless. Thus, vitamin E and selenium have biosynergistic effect and complete protection of living cells require both the nutrients in the diet.

CONCLUSION

The present investigation suggested that combination of vitamin E at a level of 200 mg and selenium at 0.4 mg/kg diet are good for improving growth rate and 100 mg vitamin E and 0.2 mg selenium per kg diet improves the immune responses of layer chickens. Thus, supplementation of vitamin E and selenium at levels above those recommended as nutritional requirements enhances disease resistance which has been attributed to significant stimulation of humoral and cellular immunity and phagocytosis.

REFERENCES

1. Linge P, The use of probiotic and yeast derivatives in India. *World Poult*, 21: 12-15, (2005).
2. McDowell LR, Reevaluation of the metabolic essentiality of the vitamins. *Asian- Australian J. Anim. Sci*, 13: 115-125, (2000).
3. Surai PF and Dvorska EJ, Dietary organic selenium and eggs: From improvements in egg quality to production of functional food. *Proceedings of IX European Symposium on the quality of eggs and egg products*, Sep 9-12, Kusadasi, Turkey (2001).
4. Surai PF, Selenium in poultry nutrition. I. Antioxidant properties, deficiency and toxicity. *World's Poult. Sci. J*, 58: 333-345, (2002).
5. Packer L, Protective role of vitamin E in biological system., *Am. J. Clin. Nutr*, 53: 1050-1055, (1991).
6. Brigelius-Flohe R, Tissue specific functions of individual glutathione peroxidases. *Biol. Med*, 27: 951-965, (1999).
7. Wolfram S, Absorption and metabolism of selenium: difference between inorganic and organic sources. In: *Biotechnology in the Feed industry. Proceedings of the 15th Annual Symposium*, Edited by T.P. Lyons, and K.A. Jacques, Nottingham University Press, Nottingham, UK, 1999, pp. 547-566.
8. Kelly MP and Power RF, Fractionation and identification of the major selenium compounds in selenized yeast. *J. of Dairy Sci*, 78 (Suppl 1): 237, (1995).
9. Edens FW, Practical applications for selenomethionine: broiler breeder reproduction. In: *Biotechnology in the Feed Industry. Proceedings of the 18th Alltech's Annual Symposium*, Edited by T.P. Lyons, and K.A. Jacques, Nottingham University Press, Nottingham, UK, 29-42, (2002)..
10. B.I.S. Nutrient requirement for poultry, Bureau of Indian Standards, I.S. 13574, (1992).
11. Kundu A, Singh DP, Mohapatra SC, Dash BB, Moidgal RP and Bisht GS, Antibody response to sheep erythrocytes in Indian



INFLUENCE OF VITAMIN-E AND SELENIUM SUPPLEMENTATION ON THE GROWTH PERFORMANCE AND ANTIBODY RESPONSES OF LAYER CHICKS

mg+0.2 mg/kg). It was also reported that body weight of sel-plex (0.2 ppm) fed broilers were increased compared to control and sodium selenite treatment groups (0.2 ppm) and the combination of sodium selenite(0.1 ppm) and sel-plex (0.1 ppm) was no more effective than sel-plex alone¹⁷.

On contrary, Yoon et al.¹⁸ reported that broiler chicks fed supplemental selenium from organic (selenium yeast A and B) or inorganic (sodium selenite) sources did not affect ($P > 0.05$) the final body weight and average daily gain. It was also reported that dietary supplementation of *Ocimum sanctum* leaf powder (0.25% and 0.5%) and organic selenium (0.3 ppm) and their combination did not significantly influence the growth rate of broilers¹⁹. The body weight gain of vitamin E and selenium supplemented groups may be due to the fact that vitamin E is an excellent biological chain breaking antioxidant that protects cells and tissues from lipoperoxidative damage induced by free radicals. Vitamin E interacts with selenium containing enzyme glutathione peroxidase to prevent the oxidative breakdown to tissue membranes and cell integrity and membrane damage²⁰.

Vitamin E and selenium supplementation either alone or in combination in layer chicks did not differ significantly on feed consumption from 1 to 8 weeks of age. These findings were in accordance with previous studies in which, Payne et al.²¹ found that feed intake was not affected ($P > 0.05$) by organic or inorganic selenium source or level of supplementation in any period of growth of broilers. The inclusion of vitamin E (250 mg/kg) in combination with either organic or inorganic selenium (0.2 mg/kg) in the diet of broilers had no effect on feed intake¹⁶. It was reported that feed intake was not affected by vitamin E supplementation under thermo-neutral conditions. However, feed intake increased with the vitamin E supplementation either singly or in combination

with vitamin C in heat-stressed Japanese quail²². On the other hand, it was reported that supplementation of vitamin E (150 and 300 IU/Kg) in the basal diet of broilers caused significant reduction in feed take²³. The variation in feed intake may be due to dosages of vitamin E and selenium supplementation, strain of the birds, season, energy content of the diet, ambient temperature, housing designs, hygienic conditions and rearing environment prevailed at the time of experiment.

The mean HA titre against SRBC, HI titre against NDV, and QAGPT titre against IBDV were significantly higher in chicks fed diets containing vitamin E (100 mg/kg) and selenium (0.2 mg/kg) compared to other treatments and control group. This report is well supported that combination of vitamin E and selenium supplementation significantly improved both humoral and cell mediated immunity in birds²⁴. Similar results have been observed in broilers fed different levels of vitamin E (0, 10, 25, 50, 100 and 200 IU/kg diet) and found better immune response against IBDV and SRBC at levels of 25 and 50 IU/kg feed respectively²⁵. Significantly higher antibody titres (HI and ELISA) against NDV were found in broilers fed diets receiving 0.06 mg/kg selenium and 150 IU/kg vitamin E²³. Increased antibody production against SRBC was observed in laying hens supplemented with combination of probiotics, yeast, vitamin E and vitamin C²⁶.

On the other hand, no significant effect on antibody titres against SRBC was observed in broilers fed vitamin E (10 and 300 IU/kg) and concluded that vitamin E supplementation increases heterophil/lymphocyte ratio, indicating that vitamin E improved the phagocytic capacity of the immune system, promoting the birds against the invasion of pathogenic microorganisms²⁷. The reason for the contradiction among the results might be the result of the difference in the species



INFLUENCE OF VITAMIN-E AND SELENIUM SUPPLEMENTATION ON THE GROWTH PERFORMANCE AND ANTIBODY RESPONSES OF LAYER CHICKS

- native vis-a-vis imported breeds of chickens. Br. Poult. Sci, 40: 40-43, (1999).
12. Siegel PB and Gross WB. Production and non-persistence of antibodies chicken to sheep erythrocytes. 1. Directional selection. Poult. Sci, 59: 1-5, (1980).
 13. Allan WH and Gough RE, A standard haemagglutination inhibition test for Newcastle disease. 1. A comparison of macro and micro methods. Veterinary Record, 95: 120-123, (1974).
 14. Wood GW, Muskett JC, Hebert CN and Thronton DH, Standardization of the quantitative agar gel precipitin test for antibodies to Infectious Bursal Disease. J. of Biological Standards, 7: 89-96, (1979).
 15. Snedecor GW and Cochran WG, Statistical methods, 8th ed., Iowa State University Press/Ames, Iowa-50010, (1989).
 16. Salman M, Selcuk Z and Muglali OH, Effect of vitamin E and selenium on performance, plasma and tissue GSH-Px activity in broilers. J. of Istanbul Vet. Sci, 3: 25-34 (2007).
 17. Upton JR, Edens FW and Ferket PR, Selenium yeast effect on broiler performance. Int. J. of Poult.Sci, 7: 798-805 (2008).
 18. Yoon I, Werner TM and Butler JM, Effect of source and concentration of selenium on growth performance and selenium retention in broiler chickens. Poult. Sci, 86: 727-730, (2007).
 19. Vara Prasad Reddy LSS, Thangavel A, Leela V and Narayana Raju KVS, Effect of dietary supplementation of Tulasi (*Ocimum sanctum*) and selenium on lipid peroxidation levels and growth rate in broiler chickens. Tamilnadu J. of Vet. & Anim. Sci, 3: 144-149, (2007).
 20. Surai PF, Effect of selenium and vitamin E content of the maternal diet on the antioxidant system of the yolk and the developing chick. British Poult. Sci, 41: 235-243, (2000).
 21. Payne RL and Southern LL, Comparison of inorganic and organic selenium sources for broilers. Poult. Sci, 84: 898-902, (2005).
 22. Sahin N, Tuzcu M., Orhan C, Onderci M, Eroksuz Y and Sahin K, The effects of vitamin C and E supplementation on heat shock protein 70 response of ovary and brain in heat-stressed quail. British Poult. Sci, 50: 259-265 (2009).
 23. Swain BK and Johri TS, Effect of supplementation of combinations of different levels of selenium and vitamin E on relative weights of some organs and serum enzyme level in broilers. Indian J. Poult. Sci, 35: 66-69, (2000).
 24. Siam SS, Mansour KM, El-Anwer EMM and El-Warith AA, Laying hens performance, hatchability, immune response and some blood constituents as affected by vitamin E and selenium supplementation under hot conditions. Egyptian Poult. Sci. J, 24: 483-496, (2004).
 25. Leshchinsky TV and Klasing KC, Relationship between the level of dietary vitamin E and the immune response of broiler chickens. Poult. Sci, 80: 1590-1599, (2001).
 26. Asli MM, Hosseini SA, Lotfollahian H and Shariatmadari F, Effects of probiotics, yeast, vitamin E and vitamin C supplements on performance and immune response of laying hen during high environmental temperature. Int. J. Poult. Sci, 6: 895-900, (2007).