

SPECIMEN FORMAT FOR THESES OF MONTH

Faculty : Science

Department : Zoology

Branch/ Area: : Environmental Biology

Sub Subject Heading: : Water pollution

Candidate's Name : M. Alaguprathana

Candidate's Address with email : #60, Ramalingam colony,
Mettupalayam Road,
Coimbatore- 641 043
Email: texuniverse.avinuty@gmail.com

Title of the thesis : Bioremediation of methyl orange from aqueous solutions using *Oedogonium subplagiostomum* AP1

(i) In Roman Script -

(ii) In roman Script

Nomenclature of Degree: : Ph.D

Month & Year of Enrolment: : December 2014

Month & Year of Registration: : January 2015

Month & Year of Submission: : January 2019

Month & Year of Award : July, 2019

Name of Supervisor : Dr. M. Poonkothai

Designation of Supervisor : Assistant Professor (SG) in Zoology

Centre/department/school in which research was conducted : Department of Zoology,
Avinashilingam Institute for Home Science and Higher Education for Women

University's Name & Address : Avinashilingam Institute for Home Science and Higher Education for Women

Abstract within 300 words:

Algae are extensively used as biosorbent for the sorption of dyes, however decolourisation and dye uptake of methyl orange by *Oedogonium subplagiostomum* AP1 has not been explored. Various process parameters namely dye concentration (100-900mg/L), biosorbent concentration (100-700mg/L), pH (2-11), temperature (20°C-60°C) and contact time (1-10 days) were optimized RSM. Desorption and regeneration efficiency of the alga was recorded and column study was conducted in the laboratory. The adsorption isotherm, kinetic and thermodynamic studies were conducted. The physico-chemical parameters of treated methyl orange were analysed. Bioassay studies such as phytotoxicity, zootoxicity, microbial toxicity and cytogenotoxicity were performed to evaluate the toxicity of the degraded metabolites of dye after decolourisation.

The alga was identified as *Oedogonium subplagiostomum* AP1. The maximum decolourisation and dye uptake of 97% and 120mg/g was observed under optimized conditions (Dye concentration: 500mg/L, biosorbent concentration: 400mg/L, pH: 6, temperature: 30°C and incubation time: 5 days). Adsorption isotherm, kinetics and thermodynamic studies showed that a monolayer adsorption of dye onto the biosorbent through chemisorption and exothermic process. The analytical studies confirmed the decolourisation of methyl orange by the accumulation of dyes onto the surface of the alga. The toxicity studies confirmed the treated methyl orange was non-toxic to flora, fauna and the micro flora. Algal compost provides an alternative for chemical fertilizer. The physical properties of the fabrics dyed using algae treated water proved its utilization in textile industries. The physicochemical parameters of treated effluent fall within the permissible limits. The molecular docking study confirms the possible interaction between the azoreductase and methyl orange.

i) Major objectives :

- To identify the alga for its potential in dye decolourisation.
- To study the kinetic behaviour of the dye using batch mode and Response Surface Methodology (RSM) to optimize the interacting process parameters to achieve maximum decolourisation.
- To assess the dye elution, regeneration and reuse efficiency of the biosorbent.
- To evaluate the biosorption mechanism of the dye onto the alga under controlled system by the computation of isotherm and kinetic parameters.

- To determine the adsorbent-adsorbate interaction using UV- vis spectrophotometer, Fourier transform infrared (FT-IR) spectrophotometry, Scanning electron microscopy (SEM) with Energy-dispersive X-ray spectroscopy (EDX) and X-ray diffraction (XRD).
- To assess the toxicity reduction potential of treated dye solution by means of phytotoxicity, zootoxicity, microbial toxicity and cytogenotoxicity assays.
- To analyze the reuse potential of dye desorbed algae and treated dye solution for composting and dyeing fabrics.
- To evaluate the physicochemical characterization of textile dyeing industrial effluent.
- To validate the relationship between protein-dye interaction through *in silico* approach.

ii) Hypothesis:

The present study clearly supported the rejection of null hypotheses and the acceptance of alternate hypotheses stating the efficiency of *Oedogonium subplagiostomum* AP1 in the removal of methyl orange from aqueous solutions.

iii) Methodology :

The alga was collected from the natural pond located near Perur, Coimbatore, Tamilnadu and identified based on its morphological and phylogenetic method.

The screening of textile dyes against the selected alga was carried out. The effectiveness of the *Oedogonium subplagiostomum* AP1 for the removal of methyl orange from aqueous solutions was executed in batch mode. Various process parameters like dye concentration, biosorbent concentration, pH, temperature and contact time were optimized and under optimal conditions the decolourisation and uptake of dye by *Oedogonium subplagiostomum* AP1 was assessed. The algal biomass and the process variables were optimised for the maximum dye removal in batch mode using Response Surface Methodology (RSM). The efficiency of different eluents to desorb the methyl orange from the algal biomass and their regeneration efficiency was recorded by adsorption-desorption experiments under optimised conditions. Column study was conducted to check the different flow rate on the removal of dye.

The capacity of an adsorbent is described by its equilibrium sorption isotherm, which is characterized by certain constant values that express the surface properties and affinity of the

biosorbent. Langmuir and Freundlich isotherm models were performed to analyse the validity of adsorption data. The mechanism of biosorption efficiency was investigated by testing the adsorption kinetic data using pseudo-first and pseudo-second order kinetic models. The nature of adsorption was evaluated by thermodynamic parameters such as Gibbs free energy, enthalpy and entropy.

The interaction of the adsorbent-adsorbate was demonstrated using UV-Vis spectral analysis, FT-IR analysis and the surface morphology of the biosorbent was documented by SEM with EDX (determine the chemical components) and XRD respectively.

Bioassay studies such as phytotoxicity, zootoxicity, microbial toxicity and cytogenotoxicity were performed to evaluate the toxicity of the degraded metabolites of dye after decolourisation.

The reuse efficiency of the dye desorbed algae and treated dye solution was subjected to composting and dyeing fabric. The composting of desorbed algae was carried out using *Eudrilus eugeniae* for 60 days. At the termination of the experiment, the algal compost was analysed for its pH, electrical conductivity, organic carbon, moisture content and NPK levels. The values obtained in the physicochemical analysis of algal compost can help to improve the soil fertility and plant growth. Thus, the algal compost can be used as an alternative for chemical fertilizer and thereby reduces the level of pollution.

The untreated and treated textile dyeing effluent was analysed for its physical parameters viz., colour, odour, pH, electrical conductivity, turbidity and chemical parameters such as alkalinity, total dissolved solids (TDS), total suspended solids (TSS), total solids (TS), total hardness, biological oxygen demand (BOD), chemical oxygen demand (COD), total alkalinity, chloride, sulphate and nitrate.

The methyl orange was subjected to *in silico* studies for their binding against the target protein (azoreductase) involved in bioremediation using the commercially available tool Glide. The docking study showed that the binding mode of methyl orange was stabilized by hydrogen bonds which might be involved in the effective interaction of the ligand (methyl orange) with the target protein (azoreductase) leading to undergo degradation.

iv) Findings:

The morphological identification indicated that the alga belongs to *Oedogonium* species. The 5.8S rRNA of *Oedogonium* species was sequenced and compared with 5.8S rDNA database of NCBI

for further identification. The strain showed 99% identity with *Oedogonium subplagiostomum*. Hence, the strain was identified as *Oedogonium subplagiostomum* AP1. The sequence of 5.8S rRNA of *Oedogonium subplagiostomum* AP1 was submitted to GenBank with the accession number KY575148.

The findings revealed that under optimal conditions (dye concentration: 500mg/L, biosorbent concentration: 400mg/L, pH: 6, temperature: 30°C and contact time: 5 days) 97% of dye was removed and 120mg/g of dye uptake was obtained from the aqueous solutions.

Optimisation carried out using Box-Behnken Design against the process variables showed maximum dye uptake and removal of 97% and 120mg/g respectively (Optimised conditions: 500mg/L dye concentration with 400mg/L biosorbent concentration, pH 6.5 and contact time 5.5 days). The values of the predicted response were in good agreement with experimental values indicating the suitability of the experimental model and the success of RSM. The biosorbent could be regenerated using 0.1N NaOH as a best desorbing agent with 83% recovery on 2nd day. A consistent reduction in the adsorption capacity of the dye was noticed after three cycles of adsorption-desorption. The column studies indicated that at low flow rate (10ml/h) maximum colour removal (97%) was noted and increased flow rate (120ml/h) reduces the decolourisation efficiency (2%). This suggests that the column packed with *Oedogonium subplagiostomum* AP1 can be developed for the removal of methyl orange in large scale level.

The results of equilibrium data well suited with the Langmuir isotherm than Freundlich isotherm model thus, proving monolayer adsorption of methyl orange on the algal surface. Analysis of data shows that the process involves pseudo-second order kinetic model and the thermodynamic study portrays the evidence for the feasibility, spontaneous and exothermic nature of the adsorption which is controlled by chemisorption process.

UV-spectral analysis revealed the disappearance of peak in treated dye solution indicating the decolourisation of methyl orange by biodegradation or biosorption. FT-IR spectra revealed the presence of functional groups namely hydroxyl, amino, N-N and C-N on the algal surface which were the main active binding sites responsible for the adsorption of methyl orange. Scanning electron micrograph revealed a smooth texture of the biosorbent surface indicating the entrapment of dyes. The EDX spectrum of *O. subplagiostomum* AP1 shows the major elements such as C, O, Mg²⁺, Si, Cl⁻, K⁺ and Ca²⁺ respectively. When the algae interacted with methyl orange the elements such as Mg²⁺, Cl⁻ and K⁺ were eliminated suggesting the possibility of the dye interaction with *O. subplagiostomum*

AP1. The XRD spectrum of the biosorbent before adsorption of methyl orange showed the crystalline nature and after dye adsorption reveals amorphous nature suggesting the alteration in the structure when the dye molecules were adsorbed.

Phytotoxicity studies were conducted to evaluate the impact of the T₁ (tap water), T₂ (untreated dye solution) and T₃ (treated dye solution) on *Tagetes erecta* by determining its biometric (germination percentage, seedling length, vigour index) and yield attributes (Height of the plant, number of flower, size and weight of the flower) on 7th and 60th day respectively. At the end of 60th day, the experimental soil samples were also analysed for their physicochemical parameters namely pH, electrical conductivity, total nitrogen, total phosphorus, total potassium, sodium, calcium, iron, copper, zinc, nickel and manganese respectively.

Seven days old seedlings of *Tagetes erecta* showed maximum germination percentage, seedling length and vigour index in T₃ plants when compared to T₂ plants indicating the non-toxic nature of the degraded metabolites whereas untreated dye solution showed inhibitory effect on the growth parameters. Similar result was recorded in the yield attributes (height of the plant, number of flower, size and weight of the flower) on 60th day old *Tagetes erecta*.

The treated dye solution might have enriched the soil nutrients which thereby enhance the growth and yield of *Tagetes erecta* whereas the untreated dye solution retarded the plant growth and yield.

Zootoxicity study was analysed to assess the impact of tap water (T₁), untreated dye (T₂) and treated dye (T₃) solutions on the behavioural response, mortality, haematological (RBC, PCV, Hb, WBC, MCV, MCH and MCHC), biochemical (plasma glucose and protein), enzymological (AST and ALT) and histological examination (gills, liver and kidney) of *Labeo rohita*.

The fishes exposed to tap water (T₁) and treated dye solution (T₃) showed normal behavioural response whereas T₂ fishes showed abnormal behaviour. The mortality rate was less in T₁ and T₃ fishes whereas T₂ fishes recorded maximum mortality. Fishes exposed to untreated dye solution showed reduction in RBC, PCV, Hb, MCHC, plasma glucose and plasma protein, increased level of WBC, MCV and MCH and also alteration in aspartate and alanine amino transaminases (AST and ALT) thereby indicating the toxicity of the dye. No such reduction and alteration was observed in haematological, biochemical and enzymological levels of fishes exposed to tap water (T₁) and treated dye solution (T₃) indicating the non-toxic nature of the degraded metabolites of dye.

Histological examinations in T₂ fishes revealed necrosis of secondary lamella, swelling in interlamella and severe haemorrhage in the gills, vacuolar degeneration, sinusoidal congestion, necrosis and haemorrhage of hepatocytes and degenerative and necrotic changes in the renal tubules, congested and shrunken glomeruli in kidney thereby indicating the toxicity of the dye. The histology of T₁ and T₃ fishes showed no structural changes indicating the non-toxic nature of the degraded metabolites of dye.

Microbial toxicity was assessed against the selected bacterial and fungal isolates by agar well diffusion technique. The toxic effect of the dye and the degraded metabolites of dye were observed by the presence of zone of inhibition against the tested microflora.

The results revealed that the degraded metabolites of dye did not show growth inhibition zone against the tested microflora suggesting its non-toxic nature. The microbial toxicity of methyl orange against the bacterial and fungal isolates revealed the zone of inhibition or stimulatory effect suggesting its toxicity.

The cytogenotoxicity studies were performed using *Allium cepa* to assess the effect of tap water (T₁), untreated (T₂) and treated dye (T₃) solutions on root growth, root length, mitotic index, mitotic inhibition/depression and chromosomal aberrations.

Allium cepa grown with tap water (T₁) exhibited maximum root growth, root length and mitotic index followed by treated dye solution (T₃) whereas minimum root growth, length and mitotic index were noticed in untreated dye solution. The mitotic inhibition and chromosomal aberrations were high in untreated dye solution (T₂) and meagre in treated dye solution and negligible in tap water. The chromosomal aberrations such as disoriented and abnormal grouping of chromosomes, vagrant and laggard chromosomes, chromosomal loss, sticky chain and disturbed metaphase, pulverized and disturbed anaphase, chromosomal displacement in anaphase, abnormal telophase, and chromosomal bridge at telophase, spindle disturbances and binucleate cells were observed in *Allium cepa* grown with untreated dye solution. No such aberrations were recorded in *Allium cepa* treated with tap water and meagre changes were noticed in *Allium cepa* exposed to treated dye solution (T₃) indicating the non-toxic nature of the degraded metabolites of dye.

In order to save water, recycling of wastewater for dyeing fabrics reduces the wastage of water. The fabrics were dyed with tap water and treated dye solution and evaluated for the physical properties namely fabric weight, thickness, strength, elongation, stiffness (warp and weft) and colour fastness.

(sunlight, pressing and crocking) respectively. The results revealed that the physical properties of the fabrics dyed with treated dye solution are comparable with that of tap water dyed fabric.

The levels of physicochemical parameters in untreated dyeing effluent does not fall within the tolerance limits prescribed by BIS for the discharge into inland surface whereas the treated effluent falls within the prescribed limits suggesting the potential of *Oedogonium subplagiostomum* AP1 in the treatment of textile dyeing effluent.

The hydrogen bond interaction from the docking studies indicates that the amino acid such as SER10, ARG12, SER18, SER18 and SER16 play an important role as catalytic residues in azoreductase. A good dock score obtained indicates the higher binding affinity towards the target binding protein.

Examiners

Internal Examiner : Dr. P.U. Mahalingam

Professor and Head, Department of Biology,
The Gandhigram Rural Institute, Gandhigram, Dindigul,

External Examiner:

Dr.S.Kamalakannan,
Professor, Division of Biotechnology,
College of Environmental and Bioresource Sciences,
Chonbuk National University,
Iksan Campus 194, 195, madong,
Iksan city, Jeonbuk
South Korea.

